

DAFTAR PUSTAKA

- Adnyani, Ayu Luih; Adiputra, Luh Made Indah Sri Handari. (2017). *Prevalensi Gangguan Fungsi Pendengaran Akibat Kebisingan Lingkungan Kerja pada Pekerja Kayu di Desa Mas Kecamatan Ubud Kabupaten Gianyar*. E-Jurnal Medika Udayana, [S.I.], v. 6, n. 12, p. 144-147, dec.. ISSN 2303-1395.
- Agrawal, S.K, Schindler, D. N., Javkler, RK, Robinson, S., (2008). *Occupational Hearing Loss in Current Diagnosis & Treatment Otolaryngology Head & Neck Surgery*, 2 nd Edition, Newyork : Mc Graww Hill Lange. pp. 732-43.
- Alberti, P.W., (1997). *Noise and the ear*. In: Stephens, D., ed. Scott- Brown"s Adult Audiology 6th Ed. Great Britain: Butterworth-Heinemann, pp.1-34.
- Aldini, G. et al. (2018). '*N-Asetilsistein as an antioxidant and disulphide breaking agent: the reasons why*', Free Radical Research.
- Arts, A.H. (1999). *Differential Diagnosis of Sensorineural Hearing Loss*. In:Cummings, C.W., Fredrickson, J.M. and Harker, L.A., eds. Otolaryngology Head and Neck Surgery 3rd Ed. St. Louis: Mosby, pp.2923-24.
- Asevedo E, et al. (2014). *Systematic review of N-Acetyl Cysteine in the treatment of addictions*. Revista Brasileira de Psiquitria ; 36 : 168-75
- Bansal M., (2013). *Diseases of Ear, Nose and Throat*, Edisi Pertama,.
- Bashiruddin J dan Indro S. (2012). *Gangguan Pendengaran Akibat Bising (Noise Induced Hearing Loss)*. Buku Ajar Ilmu Kesehatan Telinga, Hidung, Tenggorokan, Kepala dan Leher Edisi 7. Fakultas Kedokteran Universitas Indonesia.
- Bramhall, N., Konrad-Martin, D., McMillan, G., Griest, S. (2017). *Auditory Brainstem Response Altered In Humans With Noise Exposure Despite Normal Outer*

- Hair Cell Function. Ear and Hearing*, 1(38), e1-e12.
<https://doi.org/10.1097/aud.0000000000000370>
- Campbell, K.C.M., (2016). *Otoacoustic Emissions* Diakses melalui
[:http://www.emedicine.medscape.com/article/835943-overview](http://www.emedicine.medscape.com/article/835943-overview)
- Cappaert, N.L., Klis, S.F., Muijser, H., Kulig, B.M. and Smoorenburg, G.F. (2000).
Noise-induced hearing loss in rats. *Noise & Health*, 3, pp.23-32.
- Cerda M, and Pluth M. (2018). *Linking the Antioxidant Activity of N-Acetyl Cysteine to H₂S and Sulfane Sulfur Species*. *Cell Chemical Biology* ; 25 : 353-5
- Ciuman Raphael R, (2013). *Inner ear symptoms and disease: otorhinolaryngology head and neck surgery*. 17th ed. Connecticut: BC decker Inc; 2009. p. 273-6.
- Demirel, R. et al. (2009). *Noise Induces Oxidative Stress in Rat*, *Eur J Gen*
- Derekoy, F. S. et al., (2004). *Effects of Ascorbic Acid on Oxidative System and Transient Evoked Otoacoustic Emissions in Rabbits Exposed to Noise, Exposure*, (October), pp. 1775–1779.
- Doosti, A. et al., (2014). *Distortion Product Otoacoustic Emission (DPOAE) as an Appropriate Tool in Assessment of Otoprotective Effects of Antioxidants in Noise-Induced Hearing Loss (NIHL)*, *Indian Journal of Otolaryngology and Head and Neck Surgery*, 66(3), pp. 325–329. doi: 10.1007/s12070-014-0721-7.
- Emmerich, E., Richter, F., Meissner, W., Dieroff, H. (2000). *The Effect Of Impulse Noise Exposure On Distortion Product Otoacoustic Emissions In the Awake Guinea Pig*. *European Archives of Oto-Rhino-Laryngology*, 3(257), 128-132.
<https://doi.org/10.1007/s004050050208>
- Eryani YM, et al. (2017). *Faktor Risiko Terjadinya Gangguan Pendengaran Akibat Bisping*. Lampung : Medula, Volume 7, No. 4..

- Feldman L, et al. (2012). *N-Asetilsistein use for amelioration of aminoglycoside – induced ototoxicity in dialysis patients*
- Fernandez, K., Jeffers, P., Lall, K., Liberman, M., Kujawa, S. (2015). *Aging After Noise Exposure: Acceleration Of Cochlear Synaptopathy In “Recovered” Ears*. J. Neurosci., 19(35), 7509-7520. <https://doi.org/10.1523/jneurosci.5138-14.2015>
- Fetoni, A. R. et al., (2009). „*Protective effects of N-asetilsistein on noiseinduced hearing loss in guinea pigs.*“, Acta otorhinolaryngologica Italica : organo ufficiale della Società italiana di otorinolaringologia e chirurgia cervico-facciale, 29(2), pp. 70–5. Available at: [http://www.ncbi.nlm.nih.gov/pubmed/20111615%5Cnhttp://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2808688&tool=pmcentrez&ren](http://www.ncbi.nlm.nih.gov/pubmed/20111615%5Cnhttp://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2808688&tool=pmcentrez&rend)
- Gillespie, P.G. (2006). *Haircell function*. In : Otolaryngology Basic Science and Clinical Review. New York : Thieme : 332-338.
- Guyton, A.C., and Hall, J.E., (2006). *The sense of hearing*. In Guyton AC, Hall JE, eds. Textbook of medical physiology. 11th ed. Philadelphia : Elsevier Saunders, pp. 652-657.
- Habybabady, R., Mortazavi, S., Khavanin, A., Mirzaei, R., Arab, M., Mesbahzadeh, B., ... & Mohammadi, M. (2018). *Protective Effects Of N-acetyl-l-cysteine On the Density Of Spiral Ganglion Cells And Histological Changes Induced By Continuous Noise Exposure In Rats*. MJMS, 5(25), 48-58. <https://doi.org/10.21315/mjms2018.25.5.5>
- Hajavi N. (2019). *Efficacy of N-Asetilsistein in patients with dyspepsia with negative helicobacter pylori infection*. Erciyes Med ; 41(2) : 153-7

- Hall J. W. (2009). *A Guide to Otoacoustic Emissions (OAE) for Physicians*. Minnesota : Maico Diagnostics.
- Hamidian, S. M.-T., (2015). *An Additive Effect of Oral N-Acetyl Cysteine On eradication of Helicobacter Pylori*. Hindawi Publishing Corporation, Volume 2015.
- Harrianto, R. (2010). *Bahaya kerja fisik*. Dalam: Buku ajar kesehatan kerja.
- Haryuna, T.S.H., (2013). *Pengaruh Curcuminoid terhadap Paparan Bising yang Ditinjau dari Ekspresi HSP-70, NF- κ B, TLR-2, TLR-4, MMP-9 dan Kolagen Tipe IV pada Fibroblas Koklea Rattus Norvegicus (Studi Eksperimental Laboratorik Ex Vivo)*. [Disertasi] Program Studi Doktor (S3) Ilmu Kedokteran. Universitas Sumatera Utara.
- Henderson, D. et al., (2006). „*The role of oxidative stress in noise-induced hearing loss, in Ear and hearing*, pp. 1–19. doi: 10.1097/01.aud.0000191942.36672.f3.
- Hirose, K., Discolo, C., Keasler, J., Ransohoff, R. (2005). *Mononuclear Phagocytes Migrate Into the Murine Cochlea After Acoustic Trauma*. J. Comp. Neurol., 2(489), 180-194. <https://doi.org/10.1002/cne.20619>
- Jiang, Y.; Rumble, J.L.; Gleixner, A.M.; Unnithan, A.S.; Pulugulla, S.H.; Posimo, J.M.; Choi, H.J.H.; Crum, T.S.; Pant, D.B.; Leak, R.K. (2013). *N-Acetyl cysteine blunts proteotoxicity in a heat shock protein-dependent manner*. Neuroscience, 255(), 19–32. doi:10.1016/j.neuroscience.2013.09.049
- Kementerian Kesehatan RI. (2014). *Situasi Penyandang Disabilitas* : Buletin Jendela Data dan Informasi Kesehatan :. Jakarta;. 8-17 p.

- Kementrian Tenaga Kerja (1999). *Keputusan Menteri Tenaga Kerja Republik Indonesia Nomor 51 Tahun 1999 Tentang Nilai Ambang Batas Faktor Fisika di Tempat Kerja*, MEN, Jakarta.
- Kopke, R. D. et al. (2007). *NAC for noise: From the bench top to the clinic*, Hearing Research, 226(1–2), pp. 114–125. doi: 10.1016/j.heares.2006.10.008.
- Kramer, S. et al. (2006). *Efficacy of the antioxidant N-asetilsistein (NAC) in protecting ears exposed to loud music*. Journal of the American Academy of Audiology, 17(4), pp. 265–278. doi: 10.3766/jaaa.17.4.5.
- Kramer, S., Dreisbach, L., Lockwood, J., Baldwin, K., Kopke, R., Scranton, S., ... & O'Leary, M. (2006). *Efficacy Of the Antioxidant N-asetilsistein (Nac) In Protecting Ears Exposed To Loud Music*. J Am Acad Audiol, 04(17), 265-278. <https://doi.org/10.3766/jaaa.17.4.5>
- Kujawa, S.G. (2009). *Noise induce hearing loss*. In: Snow, Jr JB and Wackym, PA. Ballenger's Otorhinolaryngology Head and Neck Surgery. 17th ed. Centennial edition. People's Medical Publishing House. Philadelphia, pp.265-71.
- Laer, L., Carlsson, P.I., Ottschytsch, N., Bondeson, M.L., Konings, A., Vandeveld, A., Dieltjens, N., Fransen, E., Snyders, D., Borg, E., Raes, A. and Van Camp, G. (2006). *The contribution of genes involved potassium recycling in the inner ear to noise induced hearing loss*. Human Mutation, 27(80), pp.786-95.
- Lindblad, A., Rosenhall, U., Olofsson, Å., Hagerman, B. (2011). *The Efficacy Of N-asetilsistein To Protect the Human Cochlea From Subclinical Hearing Loss*

Caused By Impulse Noise: A Controlled Trial. Noise Health, 55(13), 392.

<https://doi.org/10.4103/1463-1741.90293>

Maison, S., Usubuchi, H., Liberman, M. (2013). *Efferent Feedback Minimizes Cochlear Neuropathy From Moderate Noise Exposure.* J. Neurosci., 13(33), 5542-5552. <https://doi.org/10.1523/jneurosci.5027-12.2013>

Mashaghi A, Bezrukavnikov S, Minde DP, Wentink AS, Kityk R, Zachmann-Brand B, et al. (2016). "Alternative modes of client binding enable functional plasticity of Hsp70". Nature. 539 (7629): 448–451. Bibcode:2016Natur.539..448M. doi:10.1038/nature20137.

Mathur, N.N., (2014). *Inner ear, noise induced hearing loss.* Available at: <http://emedicine.medscape.com/article/857813-overview#a0104>

May, L., Kramarenko, I., Brandon, C., Voelkel-Johnson, C., Roy, S., Truong, K., ... & Cunningham, L. (2013). *Inner Ear Supporting Cells Protect Hair Cells By Secreting Hsp70.* J. Clin. Invest., 8(123), 3577-3587. <https://doi.org/10.1172/jci68480>

Miller, J.M., Brown, J.N. and Schacht, J., (2003). *8-iso-prostaglandin F2 α , a product of noise exposure, reduce inner ear blood flow.* Audiol Neurotol, 8, pp.207-21.

Moller, A.R., (2006). *Physiology of the ear and the auditory nervous system.* In: Jackler, R.K., Brackmann, D.E., eds. Neurotology. 2nd ed. Philadelphia: Elsevier Mosby, pp. 52-57.

Nagashima, R., Sugiyama, C., Yoneyama, M. and Ogita, K., (2005). *Transcriptional factors in the kokhlea within the inner ear.* Journal of Pharmacological Science, 99, pp. 301-306.

- Nandi, Dhatrak. (2008). *Occupational noise-induced hearing loss in India*. Indian Journal of Occupational and environmental, 12:53-6. Pujol Remy Rebillard Guy, "Cochlear Fluid". 2014 <<http://www.cochlea.eu/en/cochlea>>
- Nassiri, P., Zare S., Esmail M.R.M., Pourbakhti A, Azam K, Golmohammadi T (2016). „*Assessment of the Effects of Different Sound Pressure Levels on Distortion Product Otoacoustic Emissions (DPOAEs) in Rats*, pp. 93–99.
- O Hong, et al. (2013). *Understanding and preventing noise-induced hearing loss*. Elsevier: Disease-a Month 59..
- O’leary S. Ototoxicity. In: Gleeson M, Browning G, Burha MJ, Clarke R, Hibbert J, Jones NS, et al, eds. (2008). *Scott-brown’s otorhinolaryngology head & neck surgery*. 7th ed. London: Edward Arnold Ltd; p. 3567-73.
- Öçalan, R., Akin, F., Yilmaz, Y., Ozlugedik, S., Gürgen, S. (2014). *Division Of the Stapedial Tendon Results In Noise-induced Damage To The Inner Ear*. Med Sci Monit, (20), 742-746. <https://doi.org/10.12659/msm.890158>
- Oghalai J.S. and Brownell W.E. (2008). *Anatomy and physiology of the ear*. In: Lalwani, A.K. Current Diagnosis dan Treatment in Otolaryngology-Head and Neck Surgery. New York: McGraw-Hill Company, pp.577-95.
- Okpala, N., (2012). *Management of blast injuries in mass casualty environments - A new algorithm - 1st place Ambroise Paré Award 2012*. Available at: http://www.mciforum.com/category/new_developments/230_management_of_blast_injuries_in_mass_casualty_environments_a_new_algorithm_-_1st_place_ambroise_par_award_2012.html.
- Pawlowsky, K.S., (2004). *Anatomy and physiology of the cochlea*. In: Roland PS, Rutka JA, eds. Ototoxicity. Hamilton: BC Decker Inc, pp. 1-15. Penerbit buku kedokteran. EGC. Jakarta, hlm.129-47.

- Probst, R., Grevers, G. and Iro, H., (2006). *Basic Otorhinolaryngology*. Newyork : Georg Thieme Verlag Stuttgart, pp.172-176 ; 260-262.
- Radomska-Leśniewska, D. M. and Skopiński, P. (2012). '*N-asetilsistein as an anti-oxidant and anti-inflammatory drug and its some clinical applications*', Central-European Journal of Immunology.
- Rewerska, A., Pawelczyk, M., Rajkowska, E., Politanski, P. and Sliwinska-Kowalska, M. (2013). *Evaluating d-methionine dose to attenuate oxidative stress-mediated hearing loss following overexposure to noise*. Eur Arch Otorhinolaryngol, 270(4), pp.1513-20.
- Roland PS, Pawlowski KS. Ototoxicity. In: Wackym PA, Snow JBS, eds (2009). *Ballenger's Otorhin olaryngology head and neck surgery*. 17th ed. Connecticut: BC decker Inc;. p. 273-6.
- Salamon, et al. (2019). *Medical and dietary uses of N-Acetilcysteine*. MDPI ; 8 : 1-16
- Salawati L. (2013). *Noise-Induced Hearing Loss*. Jurnal Kedokteran syiah kuala Volume 13 No 1. April
- Septiana, N. R., & Widowati, E. (2017). *Gangguan Pendengaran Akibat Bising*. HIGEIA (Journal of Public Health Research and Development), 1(1), 73-82.
- Sherwood L. (2006). *Telinga : Pendengaran dan Keseimbangan*. Dalam Fisiologi Manusia dari Sel ke Sistem Edisi 2. Jakarta : EGC.
- Tepel M. (2007). *N-Asetilsistein in the prevention of ototoxicity*. International Society of Nephrology ; 72 : 231-2
- Tortora J. (2017). *Principles of Anatomy and Physiology* 15th Edition. Wiley

- Ward S, ed. (2005). *Role of N-Acetylcystein in the Treatment of Acute Respiratory Disorders*. Wolters Kluwer Health,.
- Wei W, Linu A, Joshua O, Richard M, Glenn G, Ercal, et al. (2008). *Effects of N-asetilsistein amide (NACA), a thiol antioxidant on radiation-induced cytotoxicity in Chinese hamster ovary cells*. Life Sciences,; 82: 1122-30.
- Xu, F., Cheng, Y., Yan, W. (2021). *Up-regulation Of Autophagy and Apoptosis Of Cochlear Hair Cells In Mouse Models For Deafness*. Arch Med Sci, 2(17), 535-641. <https://doi.org/10.5114/aoms.2018.75348>
- Yang M, Tan H, Yang Q, Wang F, Yao H, Wei Q, Tanguay RM, Wu T. (2006). *Association of hsp70 polymorphisms with risk of noise-induced hearing loss in Chinese automobile workers*. Cell Stress Chaperones. Autumn;11(3):233-9. doi: 10.1379/csc-192r.1. PMID: 17009596; PMCID: PMC1576471.
- Yilmaz, H., Aydin, S., Sanli, A., Erdogan, B., Kibar, S., Sirvanci, S., ... & Kiran, D. (2015). *Evaluation Of the Effect Of Betahistine On Noise-induced Hearing Loss Using Distortion Product Otoacoustic Emission And Scanning Electron Microscopy*. Int Adv Otol, 1(11), 6-11. <https://doi.org/10.5152/iao.2015.368>
- Yuan, H., Wang, X., Hill, K., Chen, J., Lemasters, J., Yang, S., ... & Sha, S. (2015). *Autophagy Attenuates Noise-induced Hearing Loss By Reducing Oxidative Stress*. Antioxidants & Redox Signaling, 15(22), 1308-1324. <https://doi.org/10.1089/ars.2014.6004>
- Zuo, H., Cui, B., She, X., & Wu, M. (2008). *Changes in Guinea pig cochlear hair cells after sound conditioning and noise exposure*. Journal of Occupational Health, 50(5), 373-379.