

## DAFTAR PUSTAKA

1. Madkour AM, ELMaraghy AA, Elsayed MM. Prevalence and outcome of sepsis in respiratory intensive care unit. Egypt J Bronchol [Internet]. 2022;16(1). Tersedia pada: <https://doi.org/10.1186/s43168-022-00135-9>
2. Fathi M, Markazi-Moghaddam N, Ramezankhani A. A systematic review on risk factors associated with sepsis in patients admitted to intensive care units. Aust Crit Care [Internet]. 2019;32(2):155–64. Tersedia pada: <https://doi.org/10.1016/j.aucc.2018.02.005>
3. Lecronier M, Jung B, Molinari N, Pinot J, Similowski T, Jaber S, et al. Severe but reversible impaired diaphragm function in septic mechanically ventilated patients. Ann Intensive Care [Internet]. 2022;12(1). Tersedia pada: <https://doi.org/10.1186/s13613-022-01005-9>
4. Liu N, Ren J, Yu L, Xie J. Mechanical ventilation associated with worse survival in septic patients: a retrospective analysis of MIMIC-III. J Emerg Crit Care Med. 2020;4:14–14.
5. Toker AK, Kose S, Turken M. Comparison of sofa score, sirs, qsofa, and qsofa + l criteria in the diagnosis and prognosis of sepsis. Eurasian J Med. 2021;53(1):40–7.
6. Trzeciak S, Cinel I, Dellinger RP, Shapiro NI, Arnold RC, Parrillo JE, et al. Resuscitating the microcirculation in sepsis: The central role of nitric oxide, emerging concepts for novel therapies, and challenges for clinical trials. Acad Emerg Med. 2008;15(5):399–413.
7. Sá CSC de. Carbon Dioxide gap: a review of prognostic value and therapeutic effectiveness. Medicina (B Aires). 2022;
8. Patil VP. Mystery of PCO<sub>2</sub> gap in sepsis. Indian J Crit Care Med. 2019;23(10):443–4.
9. Gavelli F, Teboul JL, Monnet X. How can CO<sub>2</sub>-derived indices guide resuscitation in critically ill patients? J Thorac Dis. 2019;11(6):1528–37.
10. Zief Z, Schneider AG, Liaudet L. Pathophysiology and clinical applications of the veno-arterial PCO<sub>2</sub> gap. Crit Care [Internet].



- 2021;25(1). Tersedia pada: <https://doi.org/10.1186/s13054-021-03671-w>
11. Abdalazeem ES, Abdelgawad B, El-Rahman AHA. Role of central venous-arterial PCO<sub>2</sub> difference in prediction of clinical outcome in ICU septic patients. *Egypt J Anaesth* [Internet]. 2021;37(1):189–95. Tersedia pada: <https://doi.org/10.1080/11101849.2021.1910180>
  12. Bitar ZI, Maadarani OS, El-Shably AM, Elshabasy RD, Zaalouk TM. The Forgotten Hemodynamic (PCO<sub>2</sub> Gap) in Severe Sepsis. *Crit Care Res Pract.* 2020;2020:10–4.
  13. Guo Z, Wang Y, Xie C, Hua G, Ge S, Li Y. Effects of respiratory rate on venous-to-arterial CO<sub>2</sub> tension difference in septic shock patients undergoing volume mechanical ventilation. *Eur J Med Res* [Internet]. 2020;25(1):1–6. Tersedia pada: <https://doi.org/10.1186/s40001-020-00402-9>
  14. Kriswidyatomo P, Kloping YP, Jaya MG, Nugraha RA, Putri CP, Putra DH, et al. Prognostic Value of PCO<sub>2</sub> Gap in Adult Septic Shock Patients: A Systematic Review and Meta-Analysis. *Turkish J Anaesthesiol Reanim.* 2022;50(5):324–31.
  15. Barış E, Arıcı MA, Hamurtekin E. The Role of Nicotinic Anti-Inflammatory Pathway in Prostaglandin Mediated Inflammatory Response in Sepsis: A Short Review. *Clin Exp Heal Sci.* 2019;
  16. Argent AC, Kissoon N “Tex.” Mechanical Ventilation in Infection, Sepsis and Organ Failure. *Pediatr Neonatal Mech Vent From Basics to Clin Pract.* 2015;1–1642.
  17. Sakr Y, Jaschinski U, Wittebole X, Szakmany T, Lipman J, Ñamendys-Silva SA, et al. Sepsis in intensive care unit patients: Worldwide data from the intensive care over nations audit. *Open Forum Infect Dis.* 2018;5(12):1–9.
  18. Pant A, Mackraj I, Govender T. Advances in sepsis diagnosis and management: a paradigm shift towards nanotechnology. *J Biomed Sci* [Internet]. 2021;28(1):1–30. Tersedia pada: <https://doi.org/10.1186/s12929-0-00702-6>



19. Vincent JL. The Clinical Challenge of Sepsis Identification and Monitoring. *PLoS Med.* 2016;13(5):1–10.
20. Kabi A, Mohanty A, Kumar SK, Singh V, Jha MK, Gupta P. Clinical spectrum and risk factors for hospital-acquired septicemia in a tertiary care centre of North-East India Ankita. *J Fam Med Prim Care* [Internet]. 2020;6(2):169–70. Tersedia pada:  
<http://www.jfmprc.com/article.asp?issn=2249-4863;year=2017;volume=6;issue=1;spage=169;epage=170;aulast=Faizi>
21. Marasinghe MSP, Fonseka WIT, Wanishri PC, Nissanka NKSM, Silva BSS De. An Exploration of Patients' Experiences of Mechanical Ventilation. *OUSL J.* 2015;9(0):83–96.
22. Karakurt S. Noninvasive mechanical ventilation. *Marmara Med J.* 2016;29(2):32–5.
23. Purnawan I, Sutrisna E, Hidayat AI. Gambaran Respon Pasien ICU Terhadap Pemasangan Ventilator Mekanik di ICU RSUD RSUD Prof. Dr. Margono Soekarjo. *J Bionursing.* 2020;2(2):120–5.
24. Schönhofer B, Kuhlen R, Neumann P, Westhoff M, Berndt C, Sitter H. Klinische leitlinie: Nicht invasive beatmung bei akuter respiratorischer insuffizienz. *Dtsch Arztebl.* 2008;105(24):424–33.
25. Dellaca RL, Veneroni C, Farre R. Trends in mechanical ventilation: Are we ventilating our patients in the best possible way? *Breathe.* 2017;13(2):84–98.
26. Spieth PM, Koch T, Gama de Abreu M. Approaches to ventilation in intensive care. *Dtsch Arztebl Int.* 2014;111(42):714–20.
27. Zhou X-F, Yu R-G, Chen Q, Xue Y-M, Chen H. Performance of Lactate and CO<sub>2</sub>-Derived Parameters in Predicting Major Postoperative Complications After Cardiac Surgery With Cardiopulmonary Bypass: Protocol of a Diagnostic Accuracy Study. *Front Cardiovasc Med.* 21;8(October):1–6.
28. Said A, Alansary A, Al-Gendy M, El-Ozairy H, Abdulla M, Salmoun. Combination of Central Venous-Arterial Carbon Dioxide Gap With



Arterio-Venous Oxygen Content Difference During Resuscitation As a Predictor of Multi-Organ Dysfunction in Septic Patients. *Ain Shams Med J.* 2019;70(10):735–45.

29. Sayed G. The Impact of Early Goal Directed Therapy and PCO<sub>2</sub> Gap Protocol on Outcomes of High-Risk Surgical Patients Admitted to ICU: A Prospect Study *J Anesth Clin Res.* 2021;12(10001015):1012.
30. Yuan S, He H, Long Y. Interpretation of venous-to-arterial carbon dioxide difference in the resuscitation of septic shock patients. *J Thorac Dis.* 2019;11(CI):S1538–43.
31. Sugiyono. *Metode Penelitian Kuantitatif Kualitatif & RND.* Bandung: Alfabeta; 2016.
32. Loiacono LA, Shapiro DS. Detection of Hypoxia at the Cellular Level. *Crit Care Clin.* 2010;26(2):409–21.
33. Robin E, Futier E, Pires O, Fleyfel M, Tavernier B, Lebuffe G, et al. Central venous-to-arterial carbon dioxide difference as a prognostic tool in high-risk surgical patients. *Crit Care [Internet].* 2015;19(1):1–10. Tersedia pada: <http://dx.doi.org/10.1186/s13054-015-0917-6>
34. Zeng Z, Huang Q, Mao L, Wu J, An S, Chen Z, et al. The Pyruvate Dehydrogenase Complex in Sepsis: Metabolic Regulation and Targeted Therapy. *Front Nutr.* 2021;8(December):1–9.
35. Nassar B, Badr M, Van Grunderbeeck N, Temime J, Pepy F, Gasan G, et al. Central venous-to-arterial PCO<sub>2</sub> difference as a marker to identify fluid responsiveness in septic shock. *Sci Rep [Internet].* 2021;11(1):1–10. Tersedia pada: <https://doi.org/10.1038/s41598-021-96806-6>

