

Length-weight relationship and condition factor of the Celebes rainbowfish *Marosatherina ladigesii*, endemic to the Maros karst region, South Sulawesi, Indonesia

¹Sharifuddin B. A. Omar, ²Kariyanti, ¹Dewi Yanuarita, ¹Mohammad T. Umar, ²Yeni S. A. Lawi

¹ Aquatic Resources Management Study Program, Hasanuddin University, Makassar, South Sulawesi, Indonesia; ² Balik Diwa College of Marine Technology, Makassar, South Sulawesi, Indonesia. Corresponding author: S. B. A. Omar, sharifuddin@unhas.ac.id

Abstract. Locally called *beseng-beseng*, *Marosatherina ladigesii* (Ahl 1936) is a rainbowfish endemic to the Maros karst region in South Sulawesi, Indonesia. The colorful > *ladigesii* male has become popular within ornamental fish hobbyists worldwide. However there is a lack of published information on the basic biological parameters of this species. This research aimed to determine the length-weight relationship and condition factors of two wild *M. ladigesii* populations. Sampling was carried out in two studies. In the first study, samples were collected 12 times at fortnightly intervals from October 2013 to March 2014. The fish were caught using a rectangular net (3 x 1m), 0.5 cm mesh size, from Bantimurung River (N = 338) and Pattunuang River (N = 331), both in Maros District. The samples were examined in the Fisheries Biology Laboratory, Hasanuddin University. The fish collected from Bantimurung river comprised 69 males and 269 females, while those from Pattunuang River comprised 88 males and 243 females. In the second study, samples were collected 8 times from July to October 2020. The number of samples obtained from the Bantimurung River was 206 individuals, while in the Pattunuang River no fish samples were obtained. The growth pattern of both male and female *M. ladigesii* from both rivers was isometric. In both populations the female *M. ladigesii* tended to have a higher condition factor than male individuals. In the second study, the growth type of *M. ladigesii* was hypoallometric.

Key Words: endemic species, ornamental fish, biological parameters, isometric growth.

Introduction. In the field of ecology, endemism is a phenomenon where one or more species have distributions limited to a particular (restricted) geographical area. To be called endemic, a taxon must occur naturally in that particular area and nowhere else (Omar 2012). Endemism is extremely important in systematics and biogeography, as well as in conservation biology; areas with a high level of endemism and species biodiversity will generally be highly rated in terms of conservation priorities (Parenti 2011). Sulawesi Island has the highest level of freshwater fish endemism in Indonesia (Kottelat et al 1993). According to Hadiaty (2018), 68 endemic freshwater fish species have been described from Sulawesi.

The Maros karst region in South Sulawesi Province is spreaded over three Districts: Maros, Pangkajene and Kepulauan, and Barru. This area has a high level of endemism and is home to many unique species. Biospeleological experts have labelled this area as a biodiversity hot spot (Suhardjono & Ubaidillah 2012). The endemic ichthyofauna of the Maros karst area includes *Bostrychus microphthalmus*, *Dermogenys orientalis*, *Lagusia micracanthus*, *Nomorhamphus brembachi*, *Nomorhamphus liemi*, *Oryzias celebensis*, and *Marosatherina ladigesii* (Hadiaty 2012; Hadiaty 2018).

Marosatherina ladigesii (Ahl 1936), known locally as *beseng-beseng*, is a member of the Family Telmatherinidae, which includes the sailfin silversides and rainbowfishes. The Celebes rainbowfish was originally described as *Telmatherina ladigesii* (Ahl 1936), and Kottelat et al (1993) reported this fish from the Bantimurung River in the Maros karst region, in South Sulawesi. A phylogenetic analysis of the Telmatherinidae by Aarn et al (1998) found significant differences between this species and other members of the

genus *Telmatherina*, warranting classification under a separate genus, named *Marosatherina* in recognition of the region from which the type species was first described. To date, *M. ladiges* is not only the type species for the genus *Marosatherina* (Aarn et al 1998) but also the sole known species in this genus (Froese & Pauly 2018). Since this change in valid taxonomy, *M. ladiges* has been found in several other rivers within the Maros karst region, i.e. Kassikebo, Ta'deang, Pattunuang, Leang-leang, Tampala, Manrepo, and Rumbia Rivers (Hadiaty 2007, 2012).

Like other rainbowfishes, *M. ladiges* exhibits sexual dimorphism; the colorful and showy males are popular with the aquarium hobby, giving this species an economic value in the international freshwater ornamental fish trade. Demand from aquarium traders and hobbyists around the world has prompted fishermen to catch *M. ladiges* to the point where populations have declined and the species is considered to be overfished. If this level of exploitation continues, it is feared that the ornamental fishery could threaten the survival of *M. ladiges* in the wild. Together with habitat degradation, uncontrolled exploitation of this endemic species has already resulted in reduced reproductive capacity (Omar 2012). Listed as Vulnerable in the IUCN Red List of Endangered Species since 1996, the assessment of *M. ladiges* was recently updated to reflect the change in taxonomy from the genus *Telmatherina* to the currently accepted genus *Marosatherina* (Kottelat 2018).

Research on *M. ladiges* has covered several aspects of reproductive biology (Andriani 2000; Nasution et al 2006; Said & Mayasari 2007; Omar et al 2014; Kariyanti et al 2014; Kariyanti 2015; Jayadi et al 2016a; Jayadi et al 2018; Kariyanti et al 2019), bio-ecology and morphology (Andriani 2000), karyotype (Andriani 2000; Andriani et al 2004), habitat (Hadiaty 2007), and genetics (Jayadi et al 2015) as well as domestication (Said et al 2007; Jayadi et al 2016b; Kariyanti & Andi Lawi 2020), including the effect of live and artificial feed on growth (Triyanto & Said 2006). However there is still a lack of data on many aspects of the biology and ecology of this species, as evidenced by the paucity of data and information in both the IUCN Red List assessment (Kottelat 2018) and FishBase, the global database of fishes (Froese & Pauly 2018). In particular, there appear to be no published data on the growth pattern of this species in the wild. Therefore, the aim of the present research was the study of the length-weight relationship and condition factor of *M. ladiges* populations in two rivers (Bantimurung and Pattunuang) within the endemic distribution of this species.

Material and Method

Period and location. The present research was conducted in two periods. The first study was conducted from October 2013 to March 2014, while the second study from July to October 2020. Samples of the endemic fish *M. ladiges* were collected from the Bantimurung River and Pattunuang River in Maros District, South Sulawesi Province, Indonesia (Figure 1).

Sampling and sample handling. Samples were collected 12 times at fortnightly intervals in the first study, while in the second study collection was performed 8 times. The fish were captured using a rectangular multifilament net (4.7 x 1 m) with mesh size 0.3 cm. The net was set on the river bed by two fishermen, each holding one end of the net. The net was then swept to the left and the right and then lifted to the surface. The fish were carefully removed from the net, euthanized, and placed in a coolbox with ice blocks to preserve the fish cell structure during transport from the field to the laboratory.

The fish samples were analyzed at the Fish Biology Laboratory in the Fisheries Department, Faculty of Marine Science and Fisheries, Hasanuddin University, Makassar, Indonesia. The total length (TL) of each specimen was measured using calipers (0.1 mm precision). The specimens were then weighed using a digital balance (0.01 g precision). Each specimen was then dissected; the gonads were removed and observed to determine sexes.




Figure 1. *Marosatherina ladigesii* sample collection sites on the Bantimurung River (A) and Pattunuang River (B), in Maros District, South Sulawesi Province, Indonesia.

Data analysis. The length-weight relationship was analyzed overall, by sexes, by site, and by sampling period. The parameters a and b in the length-weight relationship equation $W = a L^b$ were calculated as the intercept (a) and regression coefficient (b) using the least mean squares regression method on the log-transformed data using the linear equation $\log W = \log a + b \log L$ (W = body weight, in g; L = TL, in mm) (Le Cren 1951; Hamid et al 2015).

The growth pattern was determined based on the value of b as isometric ($b = 3$); hypoallometric or negative allometric ($b < 3$); and hyperallometric or allometric positive ($b > 3$) (Froese et al 2011). A t-test was used to determine whether the mean value of b was significantly different from 3 at the 95% confidence level (Hossain et al 2013). Differences in the mean values of the regression coefficient b by sex were also evaluated using a t-test with 95% confidence interval (Fowler et al 1998).

The condition factor was analyzed by sexes. For fish populations with an isometric growth pattern, the condition factor (PI) can be determined using the equation $K = 10^5 W / L^3$ (Barnham & Baxter 1998) where W = mean body weight (in g) of the fish in a given size class and L = mean total length (in mm) of the fish in the same size class. For fish populations with an allometric growth (hypoallometric or hyperallometric), the relative condition factor (K_n) was used with the equation $K_n = W / W^*$ (Le Cren 1951) where W = the observed mass (in g) of an individual and W^* = predicted mass, which is obtained from the linear regression of the weight-length relationship of the respective population sample (aL^b).

Results

Sample number and fish size. During the first research a total of 338 *M. ladiges* were collected from Bantimurung River, comprised of 69 males and 269 females, while a total of 331 *M. ladiges* were collected from Pattunuang River, and comprised of 88 males and 243 females. The total number of *M. ladiges* in the second study collected from Bantimurung River was 206 individuals, without differentiating between male and female fish. On the other hand, in Pattunuang River there were no fish in the four months of sampling (Table 1).

Table 1

Total length and body weight of *Marosatherina ladiges* specimens collected during first and second study by sexes and site

Site: River	Sexes	Total length (mm)			Body weight (g)		
		Range	Mean	SE	Range	Mean	SE
First study (October 2013 – March 2014)							
Bantimurung	Male	34-65	48.71	1.06	0.30-3.11	1.29	0.09
Bantimurung	Female	20-70	42.74	1.19	0.06-3.06	0.94	0.08
Pattunuang	Male	30-60	48.19	0.77	0.19-2.19	1.15	0.05
Pattunuang	Female	20-55	35.33	1.10	0.05-1.74	0.54	0.05
Second study (July – October 2020)							
Bantimurung	Unknown	19-69	44.80	0.63	0.06-2.42	0.89	0.09

Length-weight relationship. Based on the t-tests performed on the values of *b* obtained (through linear regression of log-transformed data), the growth pattern of *M. ladiges* collected during the first study from both Bantimurung River and Pattunuang River was isometric ($p > 0.05$). Isometric growth indicates that all body parts grow at approximately the same rate, and thus that body proportions do not vary significantly over the size range studied. There was no significant difference in the value of *b* for male and female specimens from either the Bantimurung River or the Pattunuang River ($p > 0.05$), so that the length-weight data on males and females could be analyzed together. The length-weight curves for male and female *M. ladiges* by site and for all specimens combined, together with the regression equations obtained, are shown in Figure 2. In contrast, the growth type of *M. ladiges* during the second study from Bantimurung River was hypoallometric. The curve of the length-weight relationship of *M. ladiges* and its correlation coefficient can be seen in Figure 3.

In addition to the lack of significant difference in the *b* values by site or by sexes, the high correlation coefficient (*r*) values in Figure 2 and Figure 3 (0.8553 to 0.9642) show a very strong correlation between growth in length and increase in weight for both male and female *M. ladiges* individuals, in the Bantimurung River and the Pattunuang River.




Figure 2. Length-weight curves and equations for *Marosatherina ladigesii* in the first study. A- Males from Bantimurung River; B - Females from Bantimurung River; C - Males from Pattunuang River; D - Females from Pattunuang River; E - All specimens (male + female) from Bantimurung River; F - All specimens (male + female) from Pattunuang River.




Figure 3. Length-weight curve and equation for *Marosatherina ladiges* from Bantimurung River in the second study.

Condition factor. The range and mean values obtained for *M. ladiges* during this study are shown in Table 2.

Table 2

Condition factor of *Marosatherina ladiges* from the two study sites during first and second study

Site: River	Sexes	n	Range	Mean	SE
First study (October 2013 – March 2014)					
Bantimurung	Male	69	0.5938-2.1399	1.0107	0.0270
Bantimurung	Female	269	0.4938-2.1224	1.0192	0.0127
Bantimurung	Male + Female	338	0.4938-2.1399	1.0174	0.0115
Pattunuang	Male	88	0.1830-1.5781	0.9850	0.0180
Pattunuang	Female	243	0.1132-2.4219	1.0044	0.0140
Pattunuang	Male + Female	331	0.1132-2.4219	0.9993	0.0113
Second study (July – October 2020)					
Bantimurung	Unknown	206	0.2192-3.1671	1.0587	0.0231

Discussion

Size distribution. Nasution et al (2006) reported that body lengths of *M. ladiges* caught in several rivers in the Maros karst region ranged from 35.8 to 43.3 mm with a body weight of 0.46-2.90 g. Hadiaty (2007) reported that the total length of the fish found ranged from 15.9 to 58.0 mm. The body length and body weight of the fish were also reported by Said et al (2007). They found fish lengths in Bantimurung River ranged from 37-55 mm and body weights from 0.61 to 2.89 g, while in Pattunuang River 44-52 mm and 0.95-1.72 g. The latest data on *M. ladiges* was presented by Jayadi et al (2016a) who found body lengths ranging from 36.4-56.4 mm and body weights of 1.04-2.40 g. The size range of both male and female specimens collected during the first and second studies (Table 1) was relatively large compared to samples reported from several other studies on *M. ladiges* within the Maros karst region. On average, male *M. ladiges* were larger (both longer and heavier) than females in both Bantimurung River and Pattunuang River.

Length-weight relationship. The body weight of fishes varies exponentially with body length. This is reflected in the form of the length-weight equation $W = aL^b$. The exponent b can be used to determine the growth pattern. In fish with an isometric growth pattern ($b \approx 3$), weight increases in proportion with length. A hypoallometric or negative allometric growth pattern ($b < 3$) means that length increase proportionately faster than weight; conversely, a hyperallometric or positive allometric growth pattern ($b > 3$) means that weight increase proportionately faster than length (Froese et al 2011).

Furthermore, Froese et al (2011) consider the length-weight relationship to be extremely important for fisheries management. The equation describing the relationship can be used to convert length data into biomass data, to determine the condition factor, and to compare fish growth between fisheries management areas, as well as in studies on reproduction food habits.

There are no previous studies on the growth pattern of this fish in its native habitat, the Maros karst region. Therefore, the data obtained are compared with data on other, related, endemic species from Sulawesi. The species include *Adrianichthys oophorus*, *Glossogobius matanensis*, *Paratherina striata*, *Telmatherina antoniae*, *Telmatherina celebensis*, *Telmatherina prognatha*, and *Oryzias matanensis*. In general, these endemic fishes also tend to have isometric and hypoallometric growth patterns (Table 3).

Table 3

Regression coefficients and growth patterns of some fishes endemic to Sulawesi

Species	Sexes	n	Regression parameters			Growth type	Source
			a	b	r		
<i>Telmatherina celebensis</i>	Male	141	0.0093	3.1124	0.8958	Isometric	Furkon (2003)
	Female	132	0.0036	3.1556	0.9602	Isometric	
<i>Telmatherina celebensis</i>	Male	812	0.00078	3.0729	0.9000	Isometric	Nasution (2007)
	Female	713	0.00005	3.0476	0.9487	Isometric	
<i>Paratherina striata</i>	Male	168	0.5209	2.9740	0.9251	Isometric	Wardani (2007)
	Female	235	0.6615	3.1024	0.9566	Isometric	
<i>Glossogobius matanensis</i>	Male	1401	0.0089	2.98	0.92	Hypoallometric	Mamangkey (2010)
	Female	794	0.0851	2.30	0.88	Hypoallometric	
<i>Telmatherina antoniae</i>	Male	1437	0.00008	3.210	0.979	Hyperallometric	Tantu (2012)
	Female	1270	0.0003	2.915	0.958	Hyperallometric	
<i>Adrianichthys oophorus</i>	Male	169	0.000006	3.074	0.9701	Isometric	Gundo et al (2014)
	Female	566	0.000008	3.011	0.9487	Isometric	
	Pooled	735	0.000006	3.089	0.9607	Isometric	
<i>Lagusia micracanthus</i>	Male	307	0.00002	2.9489	0.9430	Isometric	Nur (2015)
	Female	292	0.00001	3.0600	0.9284	Isometric	
	Pooled	599	0.00002	3.0041	0.9376	Isometric	
<i>Telmatherina prognatha</i>	Male	483	0.0004	2.5995	0.8544	Hypoallometric	Chadijah et al (2019)
	Female	370	0.0006	2.4875	0.8660	Hypoallometric	
<i>Oryzias matanensis</i>	Male	410	0.0014	2.0367	0.8695	Hypoallometric	Rinandha et al (2020)
	Female	470	0.00072	2.2444	0.8523	Hypoallometric	

The regression coefficients can vary both temporally and spatially, as they are affected by many internal and external factors. These include sexes, gonad development, spawning seasons, gut fullness, health condition, habitat type and condition, the food available (quantity, quality, and size), seasonal effect, as well as capture and preservation techniques (Le Cren 1951; Wootton 1998; Ahmed et al 2011; Alavi-Yeganeh et al 2011; Froese et al 2011; Ahmed et al 2012; Hossain et al 2012; Alam et al 2013; Hamid et al 2015; Jafari et al 2016; Ahamed et al 2017; Ahmed et al 2017; Azevedo et al 2017; Shalloof & El-Far 2017; Olopade et al 2018; Mitu et al 2019; Hanif et al 2020). The value of b can vary not only between species, but also between stocks of the same species (Amin et al 2005). Kharat et al (2008) and Asadi et al (2017) both report variations in b depending on the number and size range of fish observed, while Patimar et al (2009) consider that variations in b can occur in response to changes in habitat. Harrison (2001), Acosta et al (2004), and Hamid et al (2015) consider that life-stage can

affect the length-weight relationship. None of these factors appear to have had a significant influence on the results of this study.

Condition factor. Knowing the length-weight relationship for a fish species or population enables the condition factor to be estimated. The condition factor can be used to evaluate and compare the plumpness of an individual fish or group of fishes with the average value for the population or species. The condition factor can be influenced by both biotic and abiotic environmental factors so that it can be used as an index representative of the status of the aquatic ecosystem in which the fish are living. The condition factor can also be useful in monitoring fishing intensity, ontogenetic population structure, and growth rates (Ahmed et al 2011).

Overall, in both the Bantimurung River and the Pattunuang River in the first study, females tended to have a slightly higher condition factor than males. Furthermore, for both males and females, the condition factor of fish from the Bantimurung River tended to be higher than that of fish from the Pattunuang River. The analysis of variance (ANOVA) and Student's t-test showed that in the Bantimurung River, the difference between the condition factor of male and female *M. ladiges* was not significant ($p>0.05$). Conversely, in the Pattunuang River, there was a significant difference between the condition factor of male and female *M. ladiges* ($p<0.05$). There was no significant difference in condition factor ($p>0.05$) between males from Bantimurung River and males from Pattunuang River; between females from Bantimurung River and females from Pattunuang River, or between the combined (male + female) samples from the Bantimurung River and the Pattunuang River. The condition factor in the second study in Bantimurung River shows a value that is relatively greater than the value of the condition factor in the first study

Barnham & Baxter (1998) consider that fish with a condition factor (K) of 1.00 are generally in poor condition, with a long and thin body, while fish in good condition should have a condition factor around 1.40. Values of $K>1$ can be considered indicative of environmental conditions favorable for fish growth, while values of $K<1$ indicate poor environmental conditions (Ahmed et al 2017; Singh & Serajuddin 2017). Based on these criteria, the higher mean condition factor of *M. ladiges* from Bantimurung River indicates that conditions may be more favorable than in the Pattunuang River.

As for the length-weight relationship, there are no previous studies on the condition factor of *M. ladiges* from the Maros karst region. Table 4 provides information on the condition factor of some other fishes endemic to Sulawesi. Based on the data in Tables 2 and 4, the condition factor values for *M. ladiges* in this study are similar to those found for other Sulawesi endemic fishes. Similar to the results of this study, Manangkalangi (2009) also found that in the rainbowfish *Melanotaenia arfakensis* the condition factor of females tends to be higher than that of males.

Table 4
Condition factor values reported for some fishes endemic to Sulawesi

Species	Sexes	Sample size	Range	Source
<i>Telmatherina antoniae</i>	Male	556	0.85-1.19	Sumassetiyadi (2003)
	Female	228	0.90-1.20	
<i>Telmatherina celebensis</i>	Male	812	0.93-1.21	Nasution (2007)
	Female	713	1.09-1.26	
<i>Paratherina striata</i>	Male	168	0.41-2.50	Wardani (2007)
	Female	235	0.48-2.08	
<i>Telmatherina antoniae</i>	Male	1437	0.944-1.117	Tantu (2012)
	Female	1270	0.0820-1.481	
<i>Adrianichthys oophorus</i>	Male	169	0.76-2.47	Gundo et al (2014)
	Female	566	1.04-2.12	
<i>Lagusia micracanthus</i>	Male	307	1.2710-4.0294	Nur (2015)
	Female	292	1.1264-3.5095	
<i>Oryzias</i>	Male	156	0.61-1.58	Rinandha et al (2020)
<i>matanensis</i>	Female	200	0.62-1.83	

As Baby et al (2011) pointed out, spatial and temporal variations in the condition factor of both male and female fishes can be linked to biotic and abiotic environmental conditions. Furthermore, the sex and abundance of organisms present in a given aquatic environment can affect the condition factor (Zargar et al 2012; Awasthi et al 2015; Ahmed et al 2017). The condition factor is also related to the availability of food and gonad weight (Le Cren 1951; Asadi et al 2017). According to Hossain et al (2013), the condition factor tends to remain fairly constant during the pre-spawning period, falls during spawning, and is lowest for a period just after spawning. Other factors which can result in fluctuations of fish condition factor include the size class or life-stage of the fish (Encina & Granado-Lorencio 1997; Asadi et al 2017), stress (Neff & Cargnelli 2004; Ahmed et al 2017), and other water quality parameters (Khallaf et al 2003; Olopade et al 2018). None of these factors were apparent during this study.

Conclusions. In the first study, an isometric growth pattern was observed in both male and female *M. ladiges* individuals from the Bantimurung River and the Pattunuang River, indicating weight gain is proportional to the increase in length. In contrast, a hypoallometric growth type was found in *M. ladiges* in Bantimurung River in the second study. Hypoallometric or allometric negative growth pattern ($b < 3$) means that length increase proportionately faster than weight. On average, the condition factor of the second study tended to be higher than that of the first study in the Bantimurung River. The mean condition factor of *M. ladiges* from Bantimurung River was higher than that of fish from the Pattunuang River.

Acknowledgements. We thank members of the Maros Karst Region for their participation in the field: Dewi Rahmasari Afrilia Rahim, Dian Julitha, Dwi Sabriyadi Aarsal, Mahjati Zatil Ilmi, Nadia Alimah, and Nurwahida. This study was carried out with the assistance of Unhas Basic Research grants accorded to the corresponding author under contract number 1585/UN4.22/PT.01.03/2020.

References

- Aarn, Ivantsoff W., Kottelat M., 1998 Phylogenetic analysis of Telmatherinidae (Teleostei: Atherinomorpha) with description of *Marosatherina*, a new genus from Sulawesi. Ichthyological Exploration of Freshwaters 9(3):311-323.
- Acosta A. F. G., De La Cruz Agüero G., De La Cruz Agüero J., 2004 Length-weight relationships of fish species caught in mangrove swamp in the Gulf of California (Mexico). Journal of Applied Ichthyology 20:154-155.
- Ahamed F., Saha N., Rasel M., Ahmed Z. F., Hossain M. Y., 2017 Sex ratio, length-weight and length-length relationships of *Amblypharyngodon mola* (Cyprinidae) in the Payra River, southern Bangladesh. International Journal of Fisheries and Aquatic Studies 5(4):359-362.
- Ahmed E. O., Ali M. E., Aziz A. A., 2011 Length-weight relationships and condition factors of six fish species in Atbara River and Khashm El- Girba Reservoir, Sudan. International Journal of Agriculture Sciences 3(1):65-70.
- Ahmed E. O., Ali M. E., Aziz A. A., Rafi E. M. K., 2017 Length-weight relationships and condition factors of five freshwater fish species in Roseires reservoir, Sudan. European Journal of Physical and Agricultural Sciences 5(2):26-33.
- Ahmed Z. F., Hossain M. Y., Ohtomi J., 2012 Condition, length-weight and length-length relationships of the silver hatchet Chela, *Chela cachius* (Hamilton, 1822) in the Old Brahmaputra River of Bangladesh. Journal of Freshwater Ecology 27(1):123-130.
- Alam M. M., Jahan S. N., Hussain M. A., De M., Goutham-Bharathi M. P., Magalhães A. L. B., Mazlan A. G., Simon K. D., 2013 Length-length relationship, length-weight relationship and condition factor of freshwater fish species of Bangladesh. AACL Bioflux 6(5):498-509.
- Alavi-Yeganeh M. S., Seifabadi S. J., Keivany Y., Kazemi B., Wallis G. P., 2011 Comparison of length-weight relationships in different populations and sexes of Iranian thoothcarps. Journal of Applied Ichthyology 27:1401-1403.

- Amin S. M. N., Arshad A., Haldar G. C., Shohaimi S., Ara R., 2005 Estimation of size frequency distribution, sex ratio and length-weight relationship of *Hilsa (Tenulosa ilisha)* in the Bangladesh Water. Research Journal of Agriculture and Biological Sciences 1(1):61-66.
- Andriani I., 2000 [Bioecology, morphology, caryotype and reproduction of ornamental Celebes rainbowfish (*Telmatherina ladigesii*) in the River Maros, South Sulawesi]. MSc Thesis, Post Graduate Program, Bogor Agricultural University, Bogor, Indonesia. [In Indonesian].
- Andriani I., Sugiri N., Carman O., Sjafei D. S., 2004 [The caryotype of endemic ornamental Celebes rainbowfish (*Telmatherina ladigesii*) from Maros River, South Sulawesi]. Jurnal Akuakultur Indonesia 3(2):13-18. [In Indonesian].
- Asadi H., Sattari M., Motalebi Y., Zamani-Faradonbeh M., Gheytsi A., 2017 Length-weight relationship and condition factor of seven fish species from Shahrbiyar River, Southern Caspian Sea basin, Iran. Iranian Journal of Fisheries Sciences 16(2):733-741.
- Awasthi M., Kashyap A., Serajuddin M., 2015 Length-weight relationship and condition factor of five sub-populations of *Trichogaster lalius* (Osphronemidae) of central and eastern regions of India. Journal of Ichthyology 55(6):849-853.
- Azevedo J. W. J., Castro A. C. L., Silva M. H. L., 2017 Length-weight relation, condition factor and gonadosomatic index of the whitemouth croaker, *Micropogonias furnieri* (Desmarest, 1823) (Actinopterygii: Sciaenidae), caught in Lençóis Bay, state of Maranhão, eastern Amazon, Brazil. Brazilian Journal of Oceanography 65(1):1-8.
- Baby F., Tharian J., Abraham K. M., Ramprasanth M. R., Ali A., Raghavan R., 2011 Length-weight relationship and condition factor of an endemic stone sucker, *Garra gotyla stenorhynchus* (Jerdon, 1849) from two opposite flowing rivers in southern Western Ghats. Journal of Threatened Taxa 3(6):1851-1855.
- Barnham C., Baxter A., 1998 Condition factor, K, for salmonid fish. Fisheries Notes, pp. 1-3.
- Chadijah A., Sulistiono, Haryani G. S., Affandi R., Mashar A., 2019 [Size distribution, growth pattern, and condition factor of endemic Opudi fish (*Telmatherina prognatha*) in Lake Matano, South Sulawesi]. Jurnal Ilmu Pertanian Indonesia (JIPI) 24(4):295-303. [In Indonesian].
- Encina L., Granado-Lorencio C., 1997 Seasonal changes in condition, nutrition, gonad maturation and energy content in barbel, *Barbus sclateri*, inhabiting a fluctuating river. Environmental Biology of Fishes 50(1):75-84.
- Fowler J., Cohen L., Jarvis P., 1998 Practical statistics for field biology. Second edition. John Wiley & Sons Ltd, Chichester, England, 296 p.
- Froese R., Pauly D. (eds), 2018 FishBase. World Wide Web electronic publication. www.fishbase.org, version (06/2018) [Accessed 20 December 2018].
- Froese R., Tsikliras A. C., Stergiou K. I., 2011 Editorial note on weight-length relations of fishes. Acta Ichthyologica et Piscatoria 41(4):261-263.
- Furkon A., 2003 [Food habit and growth of opudi fish *Telmatherina celebensis* in Lake Towuti, South Sulawesi]. BSc Thesis, Graduate Program, Bogor Agricultural University, Bogor, Indonesia. [In Indonesian].
- Gundo M. T., Rahardjo M. F., Lumban Batu D. T. F., Hadie W., 2014 [Length-weight relationship and condition factor of eggcarrying buntinge, *Adrianichthys oophorus* Kottelat, 1990 (Beloniformes: Adrianichthyidae) in Lake Poso, Central Sulawesi]. Jurnal Iktiologi Indonesia 14(3):225-234. [In Indonesian].
- Hadiaty R. K., 2007 [Scientific review of a rainbowfish {*Marosatherina ladigesii* (Ahl 1936)} an endemic fauna of Sulawesi]. Berita Biologi 8(6):473-479. [In Indonesian].
- Hadiaty R. K., 2012 [Fish. In: Fauna of karst and cave of Maros, South Sulawesi]. Suhardjono Y. R., Ubaidillah R. (eds), pp. 89-113, Lembaga Ilmu Pengetahuan Indonesia, Cibinong. [In Indonesian].
- Hadiaty R. K., 2018 [Taxonomical status of endemic freshwater ichthyofauna of Sulawesi]. Jurnal Iktiologi Indonesia 18(2):175-190. [In Indonesian].

- Hamid M. A., Mansor M., Nor S. A. M., 2015 Length-weight relationship and condition factor of fish populations in Temengor Reservoir: indication of environmental health. *Sains Malaysiana* 44(1):61–66.
- Hanif M. A., Siddik M. A. B., Ali M. M., 2020 Length-weight relationships of seven cyprinid fish species from the Kaptai Lake, Bangladesh. *Journal of Applied Ichthyology* 36(2):261–264.
- Harrison T. D., 2001 Length-weight relationships of fishes from South African estuaries. *Journal of Applied Ichthyology* 17:46–48.
- Hossain M. Y., Rahman M. M., Abdallah E. M., Ohtomi J., 2013 Biometric relationships of the pool barb *Puntius sophore* (Hamilton 1822) (Cyprinidae) from three major rivers of Bangladesh. *Sains Malaysiana* 42(11):1571–1580.
- Hossain M. Y., Rahman M. M., Fulanda B., Jewel M. A. S., Ahamed F., Ohtomi J., 2012 Length-weight and length-length relationships of five threatened fish species from the Jamuna (Brahmaputra River tributary) River, northern Bangladesh. *Journal of Applied Ichthyology* 28:275–277.
- Jafari O., Hedayati A. A., Keivany Y., 2016 Length-weight relationships and condition factors of *Alburnus zagrosensis* Coad, 2009, from three rivers of Tigris basin in Iran (Teleostei: Cyprinidae). *Iranian Journal of Ichthyology* 3(4):316–319.
- Jayadi, Hadijah S., Tang B., Husma A., 2016a [Reproductive biology of Celebes rainbowfish (*Marosatherina ladiges* Ahl, 1936) in some rivers of South Sulawesi]. *Jurnal Iktiologi Indonesia* 16(1):185–198. [In Indonesian].
- Jayadi, Husma A., Nursahran, Ardiansyah, Sriwahidah, 2016b Domestication of Celebes rainbowfish (*Marosatherina ladiges*). *AACL Bioflux* 9:1067–1077.
- Jayadi, Hadijah S., Harlina, Rustam, Nursahran, 2018 Embryonic and larvae of endemic Celebes rainbowfish *Marosatherina ladiges* (Ahl, 1936) (Atheriniformes: Telmatherinidae). *Pakistan Journal of Biological Sciences* 21:78–86.
- Jayadi, Tamsil A., Hadijah S., 2015 [Study of genetic variation of beseng-beseng fish (*Telmatherina ladiges*) from South Sulawesi using the Random Amplified Polymorphism DNA method (RAPD)] Prosiding Seminar Nasional Mikrobiologi Kesehatan dan Lingkungan; Januari 2015; UIN Alaudin Makassar, Indonesia. [In Indonesian].
- Kariyanti, 2015 [Reproductive biology of the endemic beseng-beseng fish (*Marosatherina ladiges* Ahl, 1936) in the Bantimurung River and the Pattunuang Asue River, Maros Regency, South Sulawesi]. MSc Thesis, Hasanuddin University, Makassar, Indonesia. [In Indonesian].
- Kariyanti, Andi Lawi Y. S., 2020 [The effect of different types of feeds on eggs on the beseng-beseng fish (*Marosatherina ladiges* Ahl, 1936)]. *Jurnal Airaha* 9(01):024–028. [In Indonesian].
- Kariyanti, Omar S. B. A., Tresnati J., 2014 [Analysis of fecundity and egg diameter of beseng-beseng fish (*Marosatherina ladiges* Ahl, 1936) in the Pattunuang Asue River and Bantimurung River, Maros Regency, South Sulawesi]. Prosiding Simposium Nasional I Kelautan dan Perikanan. MSP-09. 3 Mei 2014, Makassar, Indonesia. [In Indonesian].
- Kariyanti, Omar S. B. A., Tresnati J., 2019 [Macroscopic and microscopic identification of the gonad maturity level of the endemic beseng-beseng fish (*Marosatherina ladiges* Ahl, 1936)]. *Agrokompleks* 19(20):45–50. [In Indonesian].
- Khallaf E. A., Galal M., Authman M., 2003 The biology of *Oreochromis niloticus* in a polluted canal. *Ecotoxicology* 12:405–416.
- Kharat S. S., Khillare Y. K., Dahanukar N., 2008 Allometric scaling in growth and reproduction of a freshwater loach *Nemacheilus mooreh* (Sykes, 1839). *Electronic Journal of Ichthyology* 4(1):8–17.
- Kottelat M., 2018 *Marosatherina ladiges* (amended version of 1996 assessment). The IUCN Red List of Threatened Species 2018: e.T21574A126023332. Downloaded on 22 September 2018.
- Kottelat M., Whitten A. J., Kartikasari S. N., Wirjoatmodjo S., 1993 Freshwater Fishes of Western Indonesia and Sulawesi. Periplus Editions (HK) Ltd, Hong Kong.

- Le Cren C. D., 1951 The length-weight relationship and seasonal cycle in gonad weight and condition in the perch (*Perca fluviatilis*). *Journal of Animal Ecology* 20(2):201-219.
- Mamangkey J. J., 2010 [Biopopulation of butini (*Glossogobius matanensis*), endemic fishes in Lake Towuti, South Sulawesi]. PhD Dissertation, Post Graduate Program, Bogor Agricultural University, Bogor, Indonesia. [In Indonesian].
- Manangkalangi E., 2009 [Food, growth, and reproduction of Rainbow Arfak (*Melanotaenia arfakensis* Allen) in River Nimbai and River Aimasi, Manokwari]. MSc Thesis, Post Graduate Program, Bogor Agricultural University, Bogor, Indonesia. [In Indonesian].
- Mitu N. R., Alam M. M., Hussain M. A., Hasan M. R., Singha A. C., 2019 Length-weight and length-length relationships, sex ratio and condition factors of the Asian striped dwarf catfish *Mystus tengara* (Hamilton, 1822) (Siluriformes: Bagridae) in the Ganges River, Northwestern Bangladesh. *Iranian Journal of Ichthyology* 6(1):21-30.
- Nasution S. H., 2007 Growth and condition factor of rainbow selebensis (*Telmatherina celebensis* Boulenger) in Lake Towuti, South Celebes. *Indonesian Fisheries Research Journal* 13(2):117-123.
- Nasution S. H., Said D. S., Lukman, Triyanto, Fauzi H., 2006 [Reproductive aspects of beseng-beseng (*Telmatherina ladiges* Ahl) of various rivers in South Sulawesi]. *Proceedings of the National Seminar on Fish IV*, August 29-30, 2006, Jatiluhur, Ichthyology Society of Indonesia, pp. 83-93. [In Indonesian].
- Neff B. D., Cargnelli L. M., 2004 Relationships between condition factors, parasite load and paternity in blue gill sunfish, *Lepomis macrochirus*. *Environmental Biology of Fishes* 71(1):297-304.
- Nur M., 2015 [Reproductive biology of the endemic pirik fish (*Lagusia micracanthus* Bleeker, 1860) in South Sulawesi]. MSc Thesis, Hasanuddin University, Makassar, Indonesia. [In Indonesian].
- Olopade O. A., Dienne H. E., Eyekpeghe A., 2018 Length frequency distribution, length-weight relationship and condition factor of cichlid fishes (Teleostei: Cichlidae) from the New Calabar River, Nigeria. *Iranian Journal of Ichthyology* 5(1):74-80.
- Omar S. B. A., 2012 [The world of fish]. Gadjah Mada University Press, Yogyakarta, Indonesia. [In Indonesian].
- Omar S. B. A., Kariyanti, Tresnati J., Umar M. T., Kune S., 2014 [Sex ratios and size at sexual maturity of beseng-beseng *Marosatherina ladiges* (Ahl, 1936) endemic fish in Pattunung Asue River and Bantimurung River, Maros District, South Sulawesi]. *Prosiding Seminar Nasional Tahunan XI Hasil Penelitian Perikanan dan Kelautan Tahun 2014:BP-08*. [In Indonesian].
- Parenti L. R., 2011 Endemism and conservation of the native freshwater fish fauna of Sulawesi, Indonesia. *Prosiding Seminar Nasional Ikan VI*:1-10.
- Patimar R., Yousefi M., Hosieni S. M., 2009 Age, growth and reproduction of the sand smelt *Atherina boyeri* Risso, 1810 in the Gomishan wetland – southeast Caspian Sea. *Estuarine, Coastal and Shelf Science* 81(4):457-462.
- Rinandha A., Omar S. B. A., Tresnati J., Yanuarita D., Umar M. T., 2020 Length-weight relationship and condition factors of Matano medaka (*Oryzias matanensis* Aurich, 1935) in Towuti Lake, South Sulawesi, Indonesia. *AACL Bioflux* 13(4):1946-1954.
- Said D. S., Mayasari N., 2007 [Reproduction and growth of rainbowfish *Telmatherina ladiges* with a different sex ratio in ex-situ habitat]. *Aquacultura Indonesiana* 8(1):41-47. [In Indonesian].
- Said D. S., Lukman, Triyanto, Mayasari N., Nasution S. H., Sulaeman, Supranoto, 2007 [Domestication of Sulawesi rainbowfish *Telmatherina ladiges* through the application of artificial habitat]. *Pusat Penelitian Limnologi, LIPI*. [In Indonesian].
- Shalloof K. A. Sh., El-Far A. M., 2017 Length-weight relationship and condition factor of some fishes from the River Nile in Egypt with special reference to four *Tilapia* species. *Egyptian Journal of Aquatic Biology & Fisheries* 21(2):33-46.
- Singh M., Serajuddin M., 2017 Length-weight, length-length relationship and condition factor of *Channa punctatus* collected from three different rivers of India. *Journal of Entomology and Zoology Studies* 5(1):191-197.

- Suhardjono Y. R., Ubaidillah R. (eds), 2012 [Fauna of karst and cave of Maros, South Sulawesi]. Lembaga Ilmu Pengetahuan Indonesia, Cibinong. [In Indonesian].
- Sumassetiyadi M. A., 2003 [Reproductive aspects of opudi (*Telmatherina antoniae*) in Lake Matano, South Sulawesi]. BSc Thesis, Graduate Program, Bogor Agricultural University, Bogor, Indonesia. [In Indonesian].
- Tantu F. Y., 2012 [Reproductive ecobiology of *Telmatherina antoniae* (Kottelat, 1991) as the base of fish endemic conservation in Lake Matano, South Sulawesi]. PhD Dissertation, Post Graduate Program, Bogor Agricultural University, Bogor, Indonesia. [In Indonesian].
- Triyanto, Said D. S., 2006 [Effect of feed types on growth of rainbowfish, *Marosatherina ladigesii*]. Jurnal Iktiologi Indonesia 6(2):85-92. [In Indonesian].
- Wardani D. S. B., 2007 [Growth and morphometric-meristic characteristics of bonti-bonti (*Paratherina striata*) in Lake Towuti, South Sulawesi]. BSc Thesis, Graduate Program, Bogor Agricultural University, Bogor, Indonesia. [In Indonesian].
- Wootton R. J., 1998 Ecology of Teleost fishes. Second edition, Kluwer Academic Publishers, Fish and Fisheries Series 24, Dordrecht, 386 p.
- Zargar U. R., Yousuf A. R., Mustaq B., Jain D., 2012 Length-weight relationship of the crucian carp, *Carassius carassius* in relation to water quality, sex and season in some lentic water bodies of Kashmir Himalayas. Turkish Journal of Fisheries and Aquatic Sciences 12:683-689.

Received: 05 October 2020. Accepted: 06 November 2020. Published online: 13 November 2020.

Authors:

Sharifuddin Bin Andy Omar, Hasanuddin University, Faculty of Marine Science and Fisheries, Fisheries Department, Aquatic Resources Management Study Program, Indonesia, South Sulawesi, 90245 Makassar, Tamalanrea, e-mail: sharifuddin@unhas.ac.id

Kariyanti, Balik Diwa College of Marine Technology, Aquaculture Study Program, Indonesia, South Sulawesi, 90245 Makassar, Tamalanrea, e-mail: kariyanti@stitek-balikdiwa.ac.id

Dewi Yanuarita, Hasanuddin University, Faculty of Marine Science and Fisheries, Fisheries Department, Aquatic Resources Management Study Program, Indonesia, South Sulawesi, 90245 Makassar, Tamalanrea, e-mail: dy.satari2@gmail.com

Mohammad Tauhid Umar, Hasanuddin University, Faculty of Marine Science and Fisheries, Fisheries Department, Aquatic Resources Management Study Program, Indonesia, South Sulawesi, 90245 Makassar, Tamalanrea, e-mail: tauhid.umar72@gmail.com

Yeni Savitri Andi Lawi, Balik Diwa College of Marine Technology, Aquaculture Study Program, Indonesia, South Sulawesi, 90245 Makassar, Tamalanrea, email: yeni@stitek-balikdiwa.ac.id

This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

How to cite this article:

Omar S. B. A., Kariyanti, Yanuarita D., Umar M. T., Lawi Y. S. A., 2020 Length-weight relationship and condition factor of the Celebes rainbowfish *Marosatherina ladigesii*, endemic to the Maros karst region, South Sulawesi, Indonesia. AACL Bioflux 13(6):3384-3396.