

PAPER • OPEN ACCESS

Flow cytometry analysis on colchicine induced polyploid of Katokkon peppers (*Capsicum chinense* Jacq.)

To cite this article: R Sjahril *et al* 2021 *IOP Conf. Ser.: Earth Environ. Sci.* **807** 032024

View the [article online](#) for updates and enhancements.

A promotional banner for the 240th ECS Meeting. The banner features a colorful striped border at the top. On the left, the ECS logo is displayed in a green circle. To its right, the text reads "240th ECS Meeting" in large blue font, followed by "Oct 10-14, 2021, Orlando, Florida" in a smaller black font. Below this, it says "Register early and save up to 20% on registration costs" in bold black text, and "Early registration deadline Sep 13" in a smaller black font. At the bottom left, there is a red "REGISTER NOW" button. On the right side of the banner, there is a photograph of a diverse group of people in professional attire, smiling and clapping, suggesting a successful conference event.

ECS **240th ECS Meeting**
Oct 10-14, 2021, Orlando, Florida
**Register early and save
up to 20% on registration costs**
Early registration deadline Sep 13
REGISTER NOW

Flow cytometry analysis on colchicine induced polyploid of Katokkon peppers (*Capsicum chinense* Jacq.)

R Sjahril¹, Kasmianti^{1,4}, M Riadi², I Ridwan³, I Jamaluddin¹ and N J Panga⁵

¹Laboratory of Plant Bio-science and Reproduction Biotechnology, Faculty of Agriculture, Universitas Hasanuddin, Makassar 90245, Indonesia.

²Laboratory of Plant Breeding and Seed Science Faculty of Agriculture, Universitas Hasanuddin, Makassar 90245, Indonesia.

³Agrotechnology Study Program, Faculty of Agriculture, Universitas Hasanuddin.

⁴Master Student, Agrotechnology, Universitas Hasanuddin, Makassar, 90245, Indonesia.

⁵Department of Agrotechnology, Faculty of Agriculture, Universitas Musamus, Merauke 99611, Indonesia.

E-mail: rinaldi.sjahril@gmail.com

Abstract. A preliminary study was conducted to determine the ploidy level of Katokkon pepper after colchicine induction using flow cytometry analysis. Compared to squash method, flow cytometry analysis can provide effectively faster results of the ploidy levels of the cells. The research was conducted at the Laboratory of Plant Bioscience and Reproduction Biotechnology, Department of Agronomy, Faculty of Agriculture, Hasanuddin University Makassar. Polyploidy of the Katokkon pepper (*Capsicum chinense* Jacq.) were induced by immersing the seeds in colchicine solution at concentrations of 0.00%, 0.0125%, 0.025%, 0.050% (w/v). For each concentration, an immersion time of 1.5, 3.0 and 4.5 hours were employed, respectively. Ploidy level was analyzed using the flow cytometer machine (Partec® Cy-Flow Space). The resulting histogram of the analysis shows differences between control (0.00%) and the rest of Colchicine treatment. Based on the peak position on the histogram, the colchicine concentration and their various immersion time given did not produce tetraploid (4n) plants. Nevertheless, at 0.10% colchicine concentration with all immersion times, it was obtained mixoploid plants with 2n and 4n pairs of chromosomes.

1. Introduction

Polyloidization is one of modern methods in plant breeding to obtain superior plants with better characters by changing the genetic makeup of the plants [1]. Polyloid plants are attributed to plants that have three or more sets of chromosomes [2]. The polyloid chromosomes represent the number of more than two paired chromosome sets that can be induced chemically using colchicine. Amanah et al. [3] showed in their study that polyloid plants with three and four sets of chromosomes were obtained from the application of colchicine on cayenne pepper (*Capsicum frutescens* L.).

The successful of polyloidization can be determined by observing the number of mutated chromosomes using a simple way such as in squash method or using flow cytometry. Although the squash technique is quite simple, but it does not allow for large sample analysis because it requires a



relatively long time. Compared to the squash technique, which calculates chromosomes with the aid of a microscope [4], flow cytometry is known as a method to rapidly characterize the optical properties of cells and cell components in an individual by measuring the light emitted or scattered from cells or cell components. The method allows a lot of genetic material to be evaluated simultaneously, and the resulting information can be processed quickly. As a result, this technique allows a large-scale comparative analysis to determine polyploid levels [5]. Plant ploidy analysis can be performed quickly and accurately using flow cytometry with the advantages, among others, of ease and speed of sample preparation, as well as allowing analysis of larger numbers of samples [6].

Flow cytometry (FCM) has been widely used to detect polyploidy in plants such as *Catharanthus roseus* L. G. Don [7], *Brassica napus* L. [8], *Cajanus cajan* L. [9], and *Arabidopsis lyrata* [6]. Ploidy from plants is determined by observing data in the form of a peak curve or histogram peak shown on the monitor screen which is obtained based on the glowing rays captured by the detector on the flow cytometer [6]. Examples of flow cytometry results from haploid, diploid, and mixoploid plants is shown in Figure 1.




Figure 1. Histogram flow cytometry of *Brassica napus* L. [9].

A preliminary study carried out by the use of high concentrations of colchicine (0.0%, 0.25%, 0.50%, 0.75% and 1.0%) with varied immersion time (1.5 hours, 3 hours and 4.5 hours) in vitro in katokkon pepper (*Capsicum chinense* Jacq.) had been unsuccessful. In this previous study, the results showed that the katokkon shoots failed to develop. The failure was thought due to the colchicine concentration was too high so it was toxic [10]. Therefore, further research was carried out by lowering the colchicine concentration to get the best concentration that could produce tetraploid plants based on plant morphological characters in culture bottles and the results of ploidy level analysis using flow cytometry.


2. Methodology


The research was conducted at the Laboratory of Plant Bioscience and Reproduction Biotechnology, Faculty of Agriculture, Hasanuddin University, Makassar. The method used in planting katokkon pepper seeds (*Capsicum chinense* Jacq.) and colchicine induction is based on the results of preliminary study that have been conducted [10]. Lower concentration of colchicine (0.00%, 0.0125%, 0.025%, 0.050% and 0.10%) and immersion time of 1.5, 3.0 and 4.5 hours were used in the recent study compared to those used in the preliminary studies.

Analysis of the degree of ploidy was carried out using the Partec® Cy-Flow Space flow cytometer. Young leaves sample with a size of 0.5 x 0.5 cm was placed separately in a Petri dishes. The leaves were given 250 µl of Cystain Pi extracting buffer. Leaves chopped with a razor blade until smooth for ± 60 seconds. The extract was then filtered on a sample tube filter to obtain about 0.2 µl of phytrate. A total of 800 µl of propidium iodide dye was inserted into the sample tube. Then, the sample tube was placed into the FCM machine for analysis and the results can be obtained directly observed on a computer screen in the form of a histogram.

3. Results and discussion

Colchicine induction is a mechanism that is often used to encourage mutations, resulting in changes in the shape, size and number of chromosomes. In this study, the changes that occurred were marked visually by the size of the plantlets and histograms resulting from the flow cytometry analysis. Colchicine concentration has a significant effect on katokkon pepper (*Capsicum chinense* Jacq.) When viewed from the plantlet morphological characters (Figure 2). Colchicine-induced plantlets were characterized with shorter plants, narrower leaf width and fewer leaves than controls. Polyploid individuals have different morphological characters from diploid plants [3]. Plants treated with colchicine would have shorter plants than diploid plants as control [11].








Figure 2. Katokkon pepper plantlets (*Capsicum chinense* Jacq.) treated with several colchicine concentration and immersion time, and the flow cytometric histogram : a (0.00% colchicine, 1.5 hours), b (0.00% colchicine, 3.0 hours), c (0.00% colchicine, 4.5 hours), d (0.0125% colchicine, 1.5 hours), e (0.0125% colchicine, 3 hours), f (0.0125% colchicine, 4.5 hours), g (0.025% colchicine, 1.5 hours), h (0.025% colchicine, 3 hours), i (0.025% colchicine, 4.5 hours), j (0.05% colchicine, 1.5 hours), k (0.05% colchicine, 3 hours), l (0.05% colchicine, 4.5 hours), m (0.10% colchicine, 1.5 hours), n (0.10% colchicine, 3 hours), o (0.10% colchicine, 4.5 hours).

The histogram of the analysis using Partec® Cy-Flow Space shows the difference between the control and the colchicine treatment results. Ploidy from plants is determined by observing data in the form of a peak curve or histogram peak shown on the monitor screen which is obtained based on the glowing rays captured by the detector on the flow cytometer [6]. The peak on channel 200 shows diploid plants while on channel 400 shows tetraploid plants. Based on the peak position on the histogram, the colchicine concentration and immersion time given did not produce tetraploid plants. However, at 0.1% colchicine concentration with various immersion times, mixoploid plants were obtained, this can be seen from the results of the histogram which showed a high enough peak on channel 400 (Figure 2: m, n, and o). The peaks seen on channels 200 and 400 at 0.1% colchicine concentration indicate that the plantlets are mixoploid plants with two and four pairs of chromosomes.

4. Conclusion

Use of the flow cytometer in the analysis of polyploidization of colchicine induced katokkon pepper using a colchicine concentration of 0.0125% - 0.1% with 1.5 - 4.5 hours of immersion, respectively resulted in mixoploid plants with 2 and 4 pairs of chromosomes.

Acknowledgment

This research was supported by the Ministry of Research, Technology and Higher Education of Republic of Indonesia. The authors would like to show appreciation for the support given in funding the research.

References

- [1] Pradana D A and S Hartatik 2019 Pengaruh colchicine terhadap karakter morfologi tanaman terung (*Solanum melongena* L.) *Berk. Ilm. Pertan.* **2** (4): 155–158.
- [2] Escandón A S and J C Hagiwara 2006 A new variety of *Bacopa monneri* obtained by in vitro polyploidization *Electron. J. Biotechnol.* **9** (3).
- [3] Amanah H A, E L Arumingtyas, and S Indriyani 2016 Chromosome analysis of cayenne (*Capsicum frutescens* L.) in colchicine induced mutation *J. Appl. Hortic.* **18** (3) 217–220.
- [4] Guo X 2012 Polyploidy levels of Chinese large-flower chrysanthemum determined by flow cytometry *African J. Biotechnol.* **11** (31): 7789–7794.
- [5] Kron P, J Suda, and B C Husband 2007 Applications of flow cytometry to evolutionary and population biology *Annu. Rev. Ecol. Evol. Syst.* **38** (1): 847–876.
- [6] Dart S, P Kron, and B K Mable 2004 Characterizing polyploidy in *Arabidopsis lyrata* using chromosome counts and flow cytometry *Can. J. Bot.* **82** (2): 185–197.
- [7] Xing S H *et al.*, 2011 Induction and flow cytometry identification of tetraploids from seed-derived explants through colchicine treatments in *Catharanthus roseus* (L.) G. Don *J. Biomed. Biotechnol.* **2011**.
- [8] Takahira J, A Cousin, M N Nelson, and W A Cowling 2011 Improvement in efficiency of microspore culture to produce doubled haploid canola (*Brassica napus* L.) by flow cytometry *Plant Cell. Tissue Organ Cult.* **104** (1): 51–59.
- [9] Udensi O U and V Ontui 2013 Determination by flow cytometry polyploidy inducing-capacity of colchicine in *Cajanus cajan* (L.) mill sp, *Pakistan Journal of Biological Sciences* **16** (13): 630–635.
- [10] Kasmianti, R Sjahril, M Riadi, I Ridwan, and T AR The effects of colchicine concentration and soaking time on formation of leaves and roots of katokkon (*Capsicum chinense* Jacq.) in vitro *IOP Conf. Ser. Earth Environ. Sci.* **486** 2020.
- [11] Murni D 2010 Pengaruh perlakuan colchicine terhadap jumlah kromosom dan fenotip tanaman cabe keriting (*Capsicum annum* L.) *J. Agroekotek* **2** (1): 43–48.