

## DAFTAR PUSTAKA

- Abhishek Singh, Awadhesh Yadav and Ashish Singh. 2012. *Utilization of postnatal care for newborns and its association with neonatal mortality in India*: BMC Pregnancy and Childbirth 2012, 12:33 Page 3 of 6 /www.biomedcentral.com/1 http://471-2393/12/33
- Ahmad Rizani, Mohammad hakimi, Djauhar Ismail. 2009. *Hubungan Pengetahuan, Sikap dan Perilaku Ibu dalam pemberian Imunisasi Hepatitis B 0-7 hari di Kota Banjarmasin*, Berita kedokteran Masyarakat Vol 25, Maret 2009. Hal 12-20
- Anna M van Eijk. 2006. *Use of antenatal services and delivery care among women in rural western Kenya: a community based survey*, BioMed Central Ltd.
- Anwar. 2003. *Decision making Pattern of Maternal Health Care in East nusa Tenggara Province*. Jurnal Ekologi Kesehatan Vol 2 No 1, April 2003: 200 - 208
- Alminar Lai Dakhi. 2012. *Contribution of Delivery assistance by Skilled Attended Out Risk Redaction Of neonatal Mortality In Indonesia. Maternal and Child Health-Reproductive Health Study Program of Public Health Science*.
- Alfredo el Fort, monika T Kothari, Nouredine Abderrahim. 2008. *Association between Maternal, Birth and Newborn Characteristics and Neonatal Mortality In Five Asian Countrie*, Demographic and Health Division, Macro International Inc. 11785 Beltsville drive, calverton, MD, 20705, USA.
- Arif tiro dan Ardianingsih. 2011. *Tehnik Pengambilan Sampel*. Andira Publisher
- Azucena Bardaji. 2012. *Impact of Malaria at the End of Pregnancy on Infant Mortality and Morbidity Malaria in Pregnancy and Infant Survival*, <http://jid.oxfordjournals.org> download September 2012

- Bates AS, ET.AL. 2009. *Risk factors for underimmunization in poor urban infants*. The Journal of the American Medical Association 269:3140-3145
- Brigitte Walther. 2010. *Placental Malaria is associated with reduced early life weight development of affected children independent of low birth weight*, Walther et al. *Malaria Journal* 2010, 9:16 <http://www.malariajournal.com>
- BJ Stoll. 1997. *The global impact of infection*. *Clin Perinatol* 1997;24:1-2  
Barbara Stoll, MD Grady Memorial Hospital Division of Neonatology Atlanta,
- Bale, J., B. Stoll, A. Lucas. 2003. *And Committee on Improving Birth Outcomes. Improving birth outcomes: Meeting the challenge in the developing world*. Washington, DC: The National Academy Press
- Bang, A., R. Bang, S. Baitule, M. Reddy, and M. Deshmukh. 1999. *Effect of home-based neonatal care and management of sepsis on neonatal mortality: Field trial in rural India*. *Lancet* 354: 1955–1961.
- Bhutta, Z. A., T. Ahmed, R.E. Black, S. Cousen, K. Dewey, E. Giugliani, B.A. Haider, B. Kirkwood, S.S. Morris, H.P. Sachdev, M. Shekar . 2008. *Maternal and Child Undernutrition Study Group. 2008. What works? Interventions for maternal and child undernutrition and survival*. *Lancet* 371: 41-64.
- Bhutta, Z. A., G.L. Darmstadt, B.S. Hasan, and R.A. Haws. 2005. *Community-based interventions for improving perinatal and neonatal health outcomes in developing countries: A review of the evidence*. *Pediatrics* 115: 519-617.
- Bobak, M. Irene. 2005. *Buku Ajar Keperawatan Maternitas*, edisi 4. Alih Bahasa : Maria A. Wijayarini. Jakarta : EGC
- Bryant AS, Haas JS, McElrath TF, McCormick MC. 2006. *Predictors of compliance with the postpartum visit among women living in healthy start project areas*. *Matern Child Health J*, **10** : 511-516

- Christiana R Titaley, Cynthia L Hunter, Michael J Dibley, Peter Heywood. 2010. *Why do some women still prefer traditional birth attendants and home delivery?: a qualitative study on delivery care services in West Java Province, Indonesia*. Titaley et al. BMC Pregnancy and Childbirth 2010, 10:43 <http://www.biomedcentral.com/1471-2393/10/43>
- Christine Luxemburger, Rose McGready. 2001. *Effects of Malaria during Pregnancy on Infant Mortality in an Area of Low Malaria Transmission*, American Journal of Epidemiology Vol. 154, No. 5
- Chia-Lin Tseng, Olivia Oxlade, Dick Menzies, Anne Aspler, Kevin Schwartzman. 2011. *Cost-Effectiveness of Novel Vaccines for Tuberculosis Control: A Decision Analysis Study*, [www.biomedcentral.com/1471-2458/11/56](http://www.biomedcentral.com/1471-2458/11/56).
- Delong, G., P. Leslie, S.H. Wang, X.M. Jiang, M.L. Zhang, M. Rakeman, J.Y. Jiang, T. Ma, and X.Y. Cao. 1997. *Effect on infant mortality of iodination of irrigation water in the severely iodine deficient areas of China*. Lancet 350: 771–773.
- Donnay, F. 2000. *Maternal survival in developing countries: What has been done, what can be achieved in the next decade*. International Journal of Gynaecology and Obstetrics 70: 89-97.
- Enas Daher .2008. *Factors associated with lack of postnatal care among Palestinian women: A cross-sectional study of three clinics in the West Bank* BMC Pregnancy and Childbirth 2008, **8**:26 doi:10.1186/1471-2393-8-26
- Confidential Enquiry Into Maternal and Child Health. 2005. *"Stillbirth, Neonatal and Post-Neonatal Mortality"* . RCOG. London.
- Duffy, S. 2007. *Obstetric Hemorrhage In Gambi, Ethiopia: The Obstetrician & Gynecologist* 2007;9(2) P.121-126

- Efriza. 2007. *Determinan Kematian Neonatal di RSUD Dr. Achmad Mochtar Bukittinggi, Kesehatan Masyarakat Nasional*, Vol.2 No. 3 (diakses oktober 2011)
- Falgunee K. Parekh. 2007. *Prevalence and Risk of Plasmodium falciparum and P. vivax Malaria among Pregnant Women Living in the Hypoendemic Communities of the Peruvian Amazon*, *Am. J. Trop. Med. Hyg.*, 77(3), 2007, pp. 451–457
- Faheem G . 2010. *Latent tuberculosis among pregnant mothers in a resource poor setting in Northern Tanzania: a cross-sectional study*, *MC Infectious Diseases 2010*, **10**:52 , BMC doi:10.1186/1471-2334-10-52.
- Global Health Council report*. 2010. Available at <http://www.globalhealth.org/reports/report.php3> last accessed on August 5, 2010
- Glover, V., and O’Connor, T.G. 2002. “Effects of Antenatal Stress and Anxiety” . *Br J of Psychiatry* 180: 389-391
- Gordon D, Milberg J, Daling J, Hickok D. 1991. *Advanced Maternal Age As A Risk Factor For Cesarean Delivery*. *Obstet Gynecol*. 1991 Apr;77(4):493-497
- Graham, W., Bell, J.S. and Bullough, C.H.W. 2001. *Can skilled attendance at delivery reduce maternal mortality in developing countries? Studies in Health Services Organisation & Policy* 17:97-129
- Graham, J., Zhang, L., & Schwalberg, R. 2007. Association between maternal chronic conditions and birth outcomes in a non-Hispanic black-white Mississippi birth cohort. *Public Health Nursing* 24(4), pp 311-317.
- GC Khilnani. 2004. *Tuberculosis and Pregnancy*, *The Indian Journal of Chest Diseases & Allied Sciences* vol 46

- Hill K dan Coy Y, 2006, *Obstetri William* edisi 21, Alih bahasa: dr Andri Hartono et al. Jakarta :EGC
- Housne Ara Begum dkk. 2012. *Differentials in Place of Delivery and Delivery Assistance in Urban Slum areas, Bangladesh, Journal of Family and reproductive Health*, Vol, 6. No, 2
- Hill, Dr. Huffman, SL, Zehner, ER, Victora, C.,1999. *Breastfeeding and neonatal mortality*. LINKAGES, Academy for Educational Development. Washington, DC 1999
- Hannah Blencowe, Joy Lawn, Josh Vandelaer, Martha Roper, Simon Cousens. 2010. *Tetanus toxoid immunization to reduce mortality from neonatal tetanus*, Oxford Journal, International Journal of Epidemiology. Volume 39
- Hannah Blencowe, Simon Cousens, Luke C Mullany, Anne CC Lee Kate Kerber, Steve Wall, Gary L Darmstadt **and** Joy E Lawn. 2011. *Clean birth and postnatal care practices to reduce neonatal deaths from sepsis and tetanus: a systematic review and Delphi estimation of mortality effect BMC Public Health* 2011, **11**(Suppl 3):S11
- Ibrahim,. 2004. *Imunisasi dan Kematian Anak Balita*, Medika, Nomor 6 Tahun 17, Jakarta.
- Jean-Pierre Van geertruyden, Florence Thomas, Annette Erhart, Umberto D' Alessandro. 2004. *THE CONTRIBUTION OF MALARIA IN PREGNANCY TO PERINATAL MORTALITY*, *Am. J. Trop. Med. Hyg.*, 71(Suppl 2), 2004, pp. 35–40 Copyright © 2004 by The American Society of Tropical Medicine and Hygiene
- Iyabo T Runsewe-Abiodun, Olusoga B Ogunfowora and Bolanle M Fetuga. 2006. *Neonatal malaria in Nigeria -a 2 year review BMC Pediatrics* 2006, **6**:19 doi:10.1186/1471-2431-6-19
- Imtiaz Jehan. 2009. *Neonatal mortality, risk factors and causes: a prospective population-based cohort study in urban Pakistan Bulletin of the World Health Organization* 2009;**87**:130-138. doi: 10.2471/BLT.08.050963

- Jumiarni .2002. *Asuhan Perawatan Neonatal*. Jakarta :Penerbit Buku kedokteran EGO
- J. E. Lawn .2009. *Two million intrapartum-related stillbirths and neonatal deaths: Where, why, and what can be done?* International Journal of Gynecology and Obstetrics 107 (2009) S5–S19
- Jeanne Rini Poespoprodjo, Wendy Fobia, dkk. 2008. *Adverse Pregnancy Outcomes in an Area Where Multidrug-Resistant Plasmodium vivax and Plasmodium falciparum Infections Are Endemic*, akses dari <http://cid.oxfordjournals.org/> september 2012
- Kamran Bagheri Lankarani. 2011. *The necessity of booster vaccination after neonatal hepatitis B Vaccination*, Hapat Mon. 2011;11(6):419-421
- Kamwendo, LA. & Bullough, C. 2005. *Insight On Skilled Attendance At Birth In Malawi-The Findings Of A Structured Document And Literature Review*. Malawi Medjournal 16(2):40-42
- Katz, J., K.J. West, S. Khatri S, E.K. Pradhan, S.C. LeClerq, P. Christian, L.S. Wu, R.K. Adhikari, S.R. Shrestha, and A. Sommer. 2000. *Maternal low-dose vitamin A or beta-carotene supplementation has no effect on fetal loss and early infant mortality: A randomized cluster trial in Nepal*. American Journal of Clinical Nutrition 71: 1570–1576.
- Katz J., Keith P., Khatri S., Christian P., LeClerq S., Pradhan E. K., and Shrestha S. R. 2003. *"Risk Factors for Early Mortality in Sarlahi District Nepal"*. Bulletin WHO;81:717-725.
- Kasmiyati, Asih.et.al. 2011. *Factors Influencing Infant Mortality in Indonesia*. Journal of Population, Vol. 3, No 1, June:pp.156.
- Knippenberg, R., L.E. Lawn, G.L. Darmstadt, G. Bekyorian, H. Fogstad, N. Waleign, V. Paul, and Lancet Neonatal Survival Steering Team. 2005. *Systematic scaling up newborn care in countries*. Lancet 365: 1087-98.

- Kraner, Michael S, et.al. 2009. *Analysis of perinatal mortality and its components : time for a change*, American Journal of Epidemiology, volume 156, number 6 (493-497).
- Kristiyanasari. Weni. 2009. *Neonatus dan Asuhan Keperawatan Anak*. Yogyakarta : Nuha Medika
- Lawn, J.E., S. Cousens, and J. Zupan .2005. *4 million neonatal deaths: When? Where? Why?* Lancet 365: 891-899
- Lawn, J.E., B. McCarthy, and S.R. Ross .2001. *The healthy newborn: A reference guide for program managers*. Atlanta, Georgia: Centers for Disease Control and Prevention and Care. Akses dari: [http://www.cdc.gov/reproductivehealth/health\\_newborn.htm](http://www.cdc.gov/reproductivehealth/health_newborn.htm). September 2012
- Lawn, J., McCarthy, B.J., Ross, S.R. 2001. *"The Healthy Newborn: A Reference Manual for Program Managers "*. Geneva. New York.
- Lawn, J.E., J. Zupan, G. Begkoyian, and R. Knippenberg, Newborn survival. In D.T. Jamison, J.G. Breman, A.R. Measham, G. Alleyne, M. Claeson, D.B. Evans, P. Jha, A. Mills, and P. Musgrove (eds.). 2006. *Disease control priorities in developing countries*. 2nd edition. New York: Oxford University Press, pp. 531-549. Available at <http://files.dcp2.org/pdf/DCP/DCP27.pdf>
- Leon-Cava, N., C. Lutter, J. Ross, and L. Martin L . 2002. *Quantifying the benefits of breastfeeding: A summary of the evidence*. Washington, DC: Pan American Health Organization and LINKAGES. Akses dari: <http://www.linkagesproject.org/media/publications/Technical> Agustus 2012.
- Macdonagh, S. 2005. *Achieving Skilled Attendance For all: A Synthesis Of Current Knowledge And Recommended Actions For Scaling Up*. DFID Health Resource Centre. [Online] [Http://Www.Expandnet.Net](http://www.expandnet.net)
- Mrisho M, Obrist B, Armstrong-Schellenberg J, Haws RA, Mushi AK, Mshinda H, Tanner M, Schellenberg D. 2009. *The use of antenatal and postnatal care: perspectives and experiences of women and*

*health care providers in rural southern Tanzania. BMC pregnancy and childbirth* 2009, 9(1):10

Manish K. Singh. 2010. *Factors Influencing Utilization of ASHA Services under NRHM in Relation to Maternal Health in Rural Lucknow, Indian J Community Med.* 2010 July; 35(3): 414–419. doi: 10.4103/0970-0218.69272

Murray, J., Adeyi, G.N., Graeff, J., Fields, R., Ramuson, M., Salgad, R., Sanghvi, T. 1997. *"Emphasis Behaviors in Maternal and Child Health"*. BASICS. USA

Manandhar, D.S., D. Osrin, B.P. Shrestha, N. Mesko, J. Mossision, K.M. Tumbahangphe, S. Tamang, S. Thapa, D. Shrestha, B. Thapa, J.R. Shrestha, A. Wade, J. Borghi, H. Standing, M. Manandhar, A.M. Costello, and members of the MIRA Makwanpur trial team. 2004. *Effect of a participatory intervention with women's groups on birth outcomes in Nepal: Clusterrandomised controlled trial.* Lancet 364: 970-79.

Masanja, H., D. de Savigny, P. Smithson, J. Schellenberg, T. John, C. Mbuya, G. Upunda, T. Boerma, C. Victora, T. Smith, and H. Mshinda .2008. *Child survival gains in Tanzania: Analysis of data from Demographic and Health Surveys.* Lancet 371: 1276-1283.

Nurmiati, Besral., 2008. *Pengaruh Durasi Pemberian ASI Terhadap Ketahanan Hidup Bayi di Indonesia, Makalah Kesehatan, Vol. 12, No. 2, (diakses 4 Oktober 2011).*

Ngoc, N.T.N., M. Meriardi, H. Abdel-Aleem, G. Carroli, M. Purwar, N. Zavaleta, L. Campódonico, M.M. Ali, G.J. Hofmeyr, M. Mathai, O. Lincetto, and J. Villar. 2006. *Causes of stillbirths and early neonatal deaths: Data from 7993 pregnancies in six developing countries.* Bulletin of the World Health Organization 84: 699-705.

Othman Kakaire, Dan K Kaye and Michael O Osinde. 2011. *Male involvement in birth preparedness and complication readiness for emergency obstetric referrals in rural Uganda Kakaire et al.*



- Reproductive Health 2011, 8:12 <http://www.reproductive-health-journal.com/content/8/1/12>, akses agustus 2012
- Ozren Mamula dkk. 2009. *Complication During Pregnancy, Labor and Puerperium In Women with Increased BMI at Pregnancy Term*, Central European Journal of Medic. 4(1). 2009.71-75
- OlabisiM. Loto and Ibraheem Awowole, 2012, *Tuberculosis in Pregnancy: A Review*, Hindawi Publishing Corporation Journal of Pregnancy Volume 2012, Article ID 379271, 7 pages doi:10.1155/2012/379271
- Pattinson RC, De Jong G, Theron GB .2009. *Primary causes of total perinatally related wastage at Tygerberg Hospital. S Afr Med J* 1989; 75: 50-4 pmid: 2643836.
- Priyadi N, dkk.2008. *Analisis faktor Resiko Kematian Neonatal Studi kasus Kontrol di Kecamatan Losari Kabupaten Brebes Tahun 2006*, Jurnal Promosi Kesehatan Indonesia Vol 3/ No.1/Januari 2008.UNDIP
- Regional Committee for South-East Asia. 2003. *"Health Of The Newborn"*. World Health Assembly. Indonesia
- Rahman M, Chen LC, M Yhunos dkk. 2000. *Reduction of Neonatal Mortality by Immunization of Non Pregnant Women in Rural Bangladesh*. Pub Med 1982, 60 (2): 261 -7.
- Save the Children .2006. *State of the world's mothers 2006: Saving the lives of mothers and newborns*. Westport, Connecticut: Save the Children. Available at <http://www.savethechildren.org>. akses maret 2013
- Singh, Abhishek · Yadav, Awadhesh · Singh, Ashish. 2012. *Utilization of postnatal care for newborns and its association with neonatal mortality in India: An analytical appraisal*. MEDLINE®/PubMed® on July 29, 2012, 8:01 PM UTC.
- Shulman, C., E. Dorman, F. Cutts, K. Kawuondo, J.N. Bulmer, N. Peshu, and K. Marsh .1999. *Intermittent sulphadoxine-pyrimethimine to prevent severe anaemia secondary to malaria in pregnancy: A randomized, placebo-controlled trial*. Lancet 353: 632-636.

- Stephansson O., Dickman P. W., Johansson A. L. V., Kieler H., and Cnattingius S. 2003. *"Time of Birth and Risk of Intrapartum and Early Neonatal Death"*. *Epidemiol J*;14:218-222.
- Sudarti, Endang Khoirunnisa. 2010. *Asuhan Kebidanan Neonatus, Bayi, dan Anak Balita*. Yogyakarta ; Nuha Medika.
- Sulochana Dhakal dkk.2007. Utilisation of postnatal care among rural women in Nepal. *BMC Pregnancy and Childbirth* 2007, **7**:19 doi:10.1186/1471-2393-7-19
- Sulistiyowati, Ning., Ronoatmodjo, S., Tarigan, L.H. 2001. *Kematian Perinatal Hubungannya dengan Faktor Praktek Kesehatan Ibu Selama Kehamilan di Kota Bekasi Tahun 2001*.
- Suroyo. 2009. *Kehidupan Neonatus Di Indonesia Meninggal*. Jakarta:www.google.com
- Tinker, A.G., V.K. Paul, and J.D. Ruben. 2006. *Maternal and newborn care: The right to a healthy newborn. International Journal of Gynecology and Obstetrics* 94: 269-276.
- UNFA Country UNFPA Country Technical Services Team for East and South-East Asia, Bangkok, Thailand March 2006
- Ukwuma, Michael Chijioke. 2012. *Multiparity and Childbirth Complications in Rural Women of Northeastern Nigerian Origin IOSR. Journal of Pharmacy and Biological Sciences (IOSRJPBS) ISSN: 2278-3008 Volume 2, Issue 3 (July-August 2012), PP 01-04 www.iosrjournals.org*
- Wiknjosastro, H. 2002. *"Ilmu Kebidanan"*. Yayasan Bina Pustaka. Jakarta.
- WHO. 2003. Recommended standards for surveillance of selected vaccine-preventable diseases. Geneva, World Health Organization, 2003 (Documen WHO/V&B/03.01) documents/DocsPDF06/843.pdf, akses agustus 2012.
- World Health Organization. 2006. *"Neonatal and perinatal Mortality : Country, Region and Global Estimates "*. Geneva.

WHO .1999. *Progress towards the global elimination of neonatal tetanus, 1990-1998. Weekly Epidemiological Record* 1999;74(10):73–80.

World Health Report. 2005. *Make Every Mother and Child Count*. Geneva, World Health Organization,

Zupan J. 2005. ” *Perinatal Mortality in Developing Countries*”. N Engl J Med 352;20

Zubairu, I dkk. 2010. *Birth Preparedness, Complication Readiness and Fathers’ Participation in Maternity Care in a Northern Nigerian Community*. African Journal of Reproductive Health Mar 2010; 14(1): 2 1



## KUESIONER PENELITIAN

### I. KARAKTERISTIK RESPONDEN

#### A. Identitas orang tua

1. Nomor Responden : .....
2. Nama (Inisial Ibu) : .....
3. Nama Suami : .....
4. Umur Ibu : .....
5. Agama : .....
6. Pendidikan Terakhir : 1. Tidak sekolah      2. SD  
3. SLTP                      4. SLTA  
5. Diploma/PT
7. Jenis Pekerjaan (ibu) : 1. PNS/Peg. Swasta    2. Wiraswasta  
3. Buruh                      4. Tdk Bekerja  
5. Lainnya,.....
8. Jenis Pekerjaan (ayah) : 1. PNS/Peg. Swasta    2. Wiraswasta  
3. Buruh                      4. Tdk Bekerja  
5. Lainnya,.....
9. Alamat : .....
10. No Telp/ HP : .....

#### B. Identitas Bayi

11. Nama Bayi : .....
12. Tgl lahir : .....
13. Anak ke : .....
14. Jenis Kelamin Bayi : 1. Laki-laki              2. Perempuan

### I. KEMATIAN NEONATAL

1. Keadaan bayi waktu di lahirkan :.....
- a. Hidup                      b. Mati
2. Jika mati, usia berapa hari
- a. 0-7 hari                      b. 8-28 hari

3. Apakah usia kandungan ibu cukup bulan
  - a. Ya
  - b. Tidak
4. Berat bayi ibu sewaktu dilahirkan
  - a. <2500 g
  - b. ≥2500 g
5. Apakah ada riwayat bayi merintih waktu lahir
  - a. Ya
  - b. Tidak
6. Apakah ada riwayat kejang sewaktu bayi baru lahir
  - a. Ya
  - b. Tidak
7. Apakah bayi ibu langsung di berikan ASI setelah lahir
  - a. Ya
  - b. Tidak
8. Apakah bayi tidak mau menyusu atau memuntahkan semua atau bahkan tidak mau minum sesaat setelah lahir
  - a. Ya
  - b. Tidak
9. Apakah timbul warna kuning pada tubuh bayi atau faeses berwarna kuning pucat setelah beberapa hari kelahirannya
  - a. Ya
  - b. Tidak

## II. KUALITAS PERTOLONGAN PERSALINAN

<b>DIMENSI VARIABEL KUALITAS PERTOLONGAN PERSALINAN</b>		
<b>No</b>	<b>DIMENSI VARIABEL</b>	<b>SKOR</b>
<b>1</b>	<b>TANGIBLE ( <i>Penampilan fisik</i> )</b>	
<b>2</b>	<b>EMPATHY ( <i>Kemampu pahaman</i> )</b>	
<b>3</b>	<b>RELIABILITY ( <i>Kehandalan</i> )</b>	
<b>4</b>	<b>RESPONSIVENESS ( <i>Ketanggapan</i> )</b>	
<b>5</b>	<b>ASSURANCE ( <i>Jaminan kepastian</i> )</b>	
<b>TOTAL SKOR</b>		

KUALITAS PERTOLONGAN PERSALINAN YANG DIBERIKAN OLEH DOKTER/BIDAN		<b>Keterangan :</b> <b>STB = Sangat Tidak Baik</b> <b>TB = Tidak Baik</b> <b>CKP = Cukup</b> <b>B = Baik</b> <b>SB = Sangat Baik</b>				
Berikut ini adalah pernyataan tentang kualitas pertolongan persalinan yang telah diberikan oleh dokter/bidan pada sdr. Mohon sdr memberi tanda silang pada angka yang tertera disebelah kanan pernyataan ini.						
Dimensi TANGIBLE (Penampilan fisik)		STB	TB	CKP	B	SB
		1	2	3	4	5
1	Kebersihan ruangan bersalin, Kerapihan, dan kenyamanan ruangan					
2	Penataan eksterior, dan interior ruangan bersalin					
3	Kelengkapan partus set, kesiapan penolong persalinan, dan kebersihan					
5	Alat-alat yang dipakai					
6	Kerapihan, dan kebersihan penampilan petugas penolong persalinan					
<b>TOTAL SCORE</b>						

<b>KUALITAS PERTOLONGAN PERSALINAN YANG DIBERIKAN OLEH DOKTER/BIDAN</b>		<b>Keterangan :</b> <b>STB = Sangat Tidak Baik</b> <b>TB = Tidak Baik</b> <b>CKP = Cukup</b> <b>B = Baik</b> <b>SB = Sangat Baik</b>				
<b>Berikut ini adalah pernyataan tentang kualitas pertolongan persalinan yang telah diberikan oleh dokter/bidan pada sdr. Mohon sdr memberi tanda silang pada angka yang tertera disebelah kanan pernyataan ini.</b>						
<b>Dimensi EMPATHY (Kemampu Pahaman)</b>		<b>STB</b>	<b>TB</b>	<b>CKP</b>	<b>B</b>	<b>SB</b>
		<b>1</b>	<b>2</b>	<b>3</b>	<b>4</b>	<b>5</b>
<b>1</b>	<b>Pemberian perhatian kepada ibu selama proses persalianan</b>					
<b>2</b>	<b>Perhatian terhadap kebutuhan ibu selama proses persalinan dan keluarganya</b>					
<b>3</b>	<b>Kelengkapan partus set, kesiapan penolong persalinan, dan kebersihan</b>					
<b>4</b>	<b>Pelayanan kepada semua ibu tanpa memandang status sosialnya</b>					
<b>TOTAL SCORE</b>						



KUALITAS PERTOLONGAN PERSALINAN YANG DIBERIKAN OLEH DOKTER/BIDAN		Keterangan : STB = Sangat Tidak Baik TB = Tidak Baik CKP = Cukup B = Baik SB = Sangat Baik				
Berikut ini adalah pernyataan tentang kualitas pertolongan persalinan yang telah diberikan oleh dokter/bidan pada sdr. Mohon sdr memberi tanda silang pada angka yang tertera disebelah kanan pernyataan ini.						
Dimensi REALIBILITY (Kehandalan)		STB	TB	CKP	B	SB
		1	2	3	4	5
1	Prosedur penerimaan pasien yang cepat dan tepat					
2	Perhatian terhadap ibu dalam proses persalinan dan kepada keluarganya					
3	Pelayanan pemriksaan, dan pertolongan persalinan yang yang disesuaikan dengan kondisi ibu yang akan melahirkan					
4	Pelayanan pertolongan persalinan di kamar bersalin sesuai dengan protap					
<b>TOTAL SCORE</b>						

KUALITAS PERTOLONGAN PERSALINAN YANG DIBERIKAN OLEH DOKTER/BIDAN		Keterangan : STB = Sangat Tidak Baik TB = Tidak Baik CKP = Cukup B = Baik SB = Sangat Baik				
Berikut ini adalah pernyataan tentang kualitas pertolongan persalinan yang telah diberikan oleh dokter/bidan pada sdr. Mohon sdr memberi tanda silang pada angka yang tertera disebelah kanan pernyataan ini.						
Dimensi RESPONSIVENEES (Ketanggapan)		STB	TB	CKP	B	SB
		1	2	3	4	5
1	Kemampua penolong persalinan yang tanggap menyelesaikan penolong persalinan					
2	Penolong persalinan memberikan informasi yang jelas dan mudah dimengerti					
3	Tindakan yang cepat pada saat ibu membutuhkan					
4	Penolong persalinan bersedia membantu ibu ketika akan BAB/BAK khususnya ketika ibu baru selesai melahirkan					
<b>TOTAL SCORE</b>						

KUALITAS PERTOLONGAN PERSALINAN YANG DIBERIKAN OLEH DOKTER/BIDAN		Keterangan : STB = Sangat Tidak Baik TB = Tidak Baik CKP = Cukup B = Baik SB = Sangat Baik				
Berikut ini adalah pernyataan tentang kualitas pertolongan persalinan yang telah diberikan oleh dokter/bidan pada sdr. Mohon sdr memberi tanda silang pada angka yang tertera disebelah kanan pernyataan ini.						
Dimensi ASSURANCE (Jaminan Kepastian)		STB	TB	CKP	B	SB
		1	2	3	4	5
1	Pengetahuan dan kemampuan penolong persalinan dalam melakukan pertolongan persalinan					
2	Ketrampilan penolong persalinan dalam membantu ibu dalam proses persalinan					
3	Pelayanan yang sopan dan ramah					
5	Jaminan keamanan pelayanan dan kepercayaan terhadap penolong persalinan					
6						
<b>TOTAL SCORE</b>						

### III. KOMPLIKASI PERSALINAN

1. Apakah waktu ibu melahirkan ada komplikasi yang menyertai
  - a. Ya
  - b. TidakJika ya, lanjut ke pertanyaan berikut
2. Apakah ibu mengalami keracunan kehamilan (preeklamsi dan eklamsi)
  - a. Ya
  - b. Tidak
3. Apakah ibu juga mengalami robekan rahim (uterus)
  - a. Ya
  - b. Tidak
4. Apakah ibu mengalami robekan jalan lahir sewaktu melahirkan
  - a. Ya
  - b. Tidak
5. Apakah ibu mengalami perdarahan sewaktu melahirkan
  - a. Ya
  - b. Tidak
6. Apakah ibu mengalami persalinan macet sewaktu persalinan
  - a. Ya
  - b. TidakJika Ya
7. Apakah ibu dirangsang dengan obat-obatan tertentu (induksi persalinan) untuk merangsang nyeri atau kontraksi uterus
  - a. Ya
  - b. Tidak
8. Apakah ibu mengalami ketuban pecah dini
  - a. Ya
  - b. Tidak
9. Apakah ibu mengalami infeksi pasca persalinan (infeksi postpartum)
  - a. Ya
  - b. Tidak
10. Apakah sewaktu melahirkan ibu melahirkan bayi besar (makrosomia) sehingga bahunya tertahan di jalan lahir dan susah untuk dilahirkan (distosia bahu)
  - a. Ya
  - b. Tidak

### IV. PEMANFAATAN PERAWATAN NEONATAL

1. Apakah ibu pernah kontak dengan ke petugas kesehatan pada periode neonatal?
  - a. Ya
  - b. Tidak pernah
2. Berapa kali ibu kontak dengan petugas kesehatan
  - a. <2 kali
  - b. ≥2 kali

3. Saat pemeriksaan apakah petugas menimbang berat badan bayi ibu
  - a. Ya
  - b. Tidak
4. Apakah petugas juga mengukur panjang badan bayi ibu
  - a. Ya
  - b. Tidak
5. Apakah petugas memeriksa kondisi tali pusat bayi ibu
  - a. Ya
  - b. Tidak
6. Apakah petugas mengukur lingkar kepala bayi
  - a. Ya
  - b. Tidak
7. Apakah petugas juga mengukur tanda-tanda vital pada bayi ibu
  - a. Ya
  - b. Tidak
8. Apakah petugas menanyakan tentang pola BAB/BAK bayi
  - a. Ya
  - b. Tidak
  - b. Tidak

#### **V. PEMBERIAN IMUNISASI**

1. Apakah bayi ibu diberikan imunisasi HB0 segera setelah lahir ?
  - a. Ya
  - b. Tidak
2. Apakah bayi ibu diberikan imunisasi BCG pada masa neonatal ?
  - a. Ya
  - b. Tidak
3. Apakah bayi ibu diberikan imunisasi Polio pada masa neonatal ?
  - a. Ya
  - b. Tidak
4. Apakah petugas memberitahukan kapan pemberian imunisasi ulangan?
  - a. Ya
  - b. Tidak

#### **VI. RIWAYAT PENYAKIT IBU**

1. Apakah ada riwayat penyakit yang menyertai ibu selama proses kehamilan
  - a. Ya
  - b. Tidak
2. Apakah selama hamil ibu menderita penyakit malaria
  - a. Ya
  - b. Tidak

Jika ya, lanjut ke pertanyaan berikut:
3. Apakah ibu mendapat pengobatan malaria dari dokter atau bidan ?
  - a. Ya
  - b. Tidak



## HASIL ANALISIS DATA BIVARIAT

### Crosstabs

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
kualitaspertolongan * kematian neonatal	138	100.0%	0	.0%	138	100.0%
komplikasipersalinan * kematian neonatal	138	100.0%	0	.0%	138	100.0%
pemanfaatanperawatan neonatal * kematian neonatal	138	100.0%	0	.0%	138	100.0%
pemberianimunisasi * kematian neonatal	138	100.0%	0	.0%	138	100.0%
riwayatpenyakit * kematian neonatal	138	100.0%	0	.0%	138	100.0%

### kualitaspertolongan \* kematian neonatal

Crosstab

			kematian neonatal		Total
			kontrol	kasus	
kualitaspertolongan	baik	Count	53	17	70
		% within kematian neonatal	75.7%	25.0%	50.7%
	tidakbaik	Count	17	51	68
		% within kematian neonatal	24.3%	75.0%	49.3%
Total		Count	70	68	138
		% within kematian neonatal	100.0%	100.0%	100.0%

### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	35.493 <sup>a</sup>	1	.000	.000	.000
Continuity Correction <sup>b</sup>	33.493	1	.000		
Likelihood Ratio	37.193	1	.000		
Fisher's Exact Test					
Linear-by-Linear Association	35.236	1	.000		
N of Valid Cases	138				

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 33.51.

b. Computed only for a 2x2 table

### Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for kualitaspertolongan (baik / tidakbaik)	9.353	4.311	20.290
For cohort kematian neonatal = control	3.029	1.965	4.667
For cohort kematian neonatal = kasus	.324	.209	.501
N of Valid Cases	138		

### komplikasipersalinan \* kematian neonatal

#### Crosstab

			kematian neonatal		Total
			kontrol	kasus	
komplikasipersalinan	tidakadakomplikasi	Count	46	6	52
		% within kematian neonatal	65.7%	8.8%	37.7%
	adakomplikasi	Count	24	62	86
		% within kematian neonatal	34.3%	91.2%	62.3%
Total	Count	70	68	138	
	% within kematian neonatal	100.0%	100.0%	100.0%	



### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	47.541 <sup>a</sup>	1	.000	.000	.000
Continuity Correction <sup>b</sup>	45.149	1	.000		
Likelihood Ratio	52.250	1	.000		
Fisher's Exact Test					
Linear-by-Linear Association	47.196	1	.000		
N of Valid Cases	138				

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 25.62.

b. Computed only for a 2x2 table

### Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for komplikasipersalinan (tidakadakomplikasi / adakomplikasi)	19.806	7.489	52.377
For cohort kematian neonatal = kontrol	3.170	2.226	4.514
For cohort kematian neonatal = kasus	.160	.075	.344
N of Valid Cases	138		

### pemanfaatanperawatan neonatal \* kematian neonatal

#### Crosstab

			kematian neonatal		Total
			kontrol	kasus	
pemanfaatanperawatan neonatal	lengkap	Count	60	38	98
		% within kematian neonatal	85.7%	55.9%	71.0%
	tidaklengkap	Count	10	30	40
		% within kematian neonatal	14.3%	44.1%	29.0%

Total	Count	70	68	138
	% within kematian neonatal	100.0%	100.0%	100.0%

#### Chi-Square Tests

	Value	Df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	14.913 <sup>a</sup>	1	.000		
Continuity Correction <sup>b</sup>	13.499	1	.000		
Likelihood Ratio	15.417	1	.000		
Fisher's Exact Test				.000	.000
Linear-by-Linear Association	14.805	1	.000		
N of Valid Cases	138				

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 19.71.

b. Computed only for a 2x2 table

#### Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for pemanfaatanperawatan neonatal (lengkap / tidaklengkap)	4.737	2.080	10.787
For cohort kematian neonatal = kontrol	2.449	1.400	4.285
For cohort kematian neonatal = kasus	.517	.381	.702
N of Valid Cases	138		

## pemberian imunisasi \* kematian neonatal

Crosstab

			kematian neonatal		Total
			kontrol	kasus	
pemberian imunisasi	lengkap	Count	70	49	119
		% within kematian neonatal	100.0%	72.1%	86.2%
	tidak lengkap	Count	0	19	19
		% within kematian neonatal	.0%	27.9%	13.8%
Total		Count	70	68	138
		% within kematian neonatal	100.0%	100.0%	100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	22.682 <sup>a</sup>	1	.000		
Continuity Correction <sup>b</sup>	20.389	1	.000		
Likelihood Ratio	30.036	1	.000		
Fisher's Exact Test				.000	.000
Linear-by-Linear Association	22.517	1	.000		
N of Valid Cases	138				

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 9.36.

b. Computed only for a 2x2 table

Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
For cohort kematian neonatal = kasus	.412	.332	.510
N of Valid Cases	138		

## riwayatpenyakit \* kematian neonatal

Crosstab

			kematian neonatal		Total
			kontrol	kasus	
riwayatpenyakit tidakadariwayat	Count	69	57	126	
	% within kematian neonatal	98.6%	83.8%	91.3%	
adariwayat	Count	1	11	12	
	% within kematian neonatal	1.4%	16.2%	8.7%	
Total	Count	70	68	138	
	% within kematian neonatal	100.0%	100.0%	100.0%	

Chi-Square Tests

	Value	Df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	9.449 <sup>a</sup>	1	.002	.002	.002
Continuity Correction <sup>b</sup>	7.683	1	.006		
Likelihood Ratio	10.867	1	.001		
Fisher's Exact Test					
Linear-by-Linear Association	9.381	1	.002		
N of Valid Cases	138				

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 5.91.

b. Computed only for a 2x2 table

Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for riwayatpenyakit (tidakadariwayat / adariwayat)	13.316	1.669	106.262
For cohort kematian neonatal = kontrol	6.571	1.000	43.205

For cohort kematian neonatal = kasus	.494	.382	.638
N of Valid Cases	138		

## HASIL ANALISIS DATA MULTIVARIAT

### Logistic Regression

#### Block 0: Beginning Block

**Classification Table<sup>a,b</sup>**

Observed			Predicted		
			kematian neonatal		Percentage Correct
			kontrol	kasus	
Step 0	kematian neonatal	kontrol	70	0	100.0
		kasus	68	0	
Overall Percentage					50.7

a. Constant is included in the model.

b. The cut value is .500

**Variables in the Equation**

		B	S.E.	Wald	df	Sig.	Exp(B)
Step 0	Constant	-.029	.170	.029	1	.865	.971

**Variables not in the Equation**

			Score	df	Sig.
Step 0	Variables	kualitas_x(1)	35.493	1	.000
		kmplks_x(1)	47.541	1	.000
		pmnftn_x(1)	14.913	1	.000
		imun_x(1)	22.682	1	.000
		rwyt_x(1)	9.449	1	.002
Overall Statistics			76.414	5	.000

## Block 1: Method = Backward Stepwise (Wald)

**Omnibus Tests of Model Coefficients**

		Chi-square	df	Sig.
Step 1	Step	101.386	5	.000
	Block	101.386	5	.000
	Model	101.386	5	.000
Step 2 <sup>a</sup>	Step	-6.874	1	.009
	Block	94.512	4	.000
	Model	94.512	4	.000

a. A negative Chi-squares value indicates that the Chi-squares value has decreased from the previous step.

**Model Summary**

Step	-2 Log likelihood	Cox & Snell R Square	Nagelkerke R Square
1	89.894 <sup>a</sup>	.520	.694
2	96.767 <sup>b</sup>	.496	.661

a. Estimation terminated at iteration number 20 because maximum iterations has been reached. Final solution cannot be found.

b. Estimation terminated at iteration number 6 because parameter estimates changed by less than .001.

**Hosmer and Lemeshow Test**

Step	Chi-square	df	Sig.
1	11.875	6	.065
2	12.825	5	.025

**Classification Table<sup>a</sup>**

Observed			Predicted		
			kematian neonatal		Percentage Correct
			kontrol	kasus	
Step 1	kematian neonatal	kontrol	63	7	90.0
		kasus	12	56	82.4
Overall Percentage					86.2
Step 2	kematian neonatal	kontrol	63	7	90.0
		kasus	12	56	82.4
Overall Percentage					86.2

a. The cut value is .500

**Variables in the Equation**

	B	S.E.	Wald	df	Sig.	Exp(B)	95% C.I. for EXP(B)		
							Lower	Upper	
Step 1 <sup>a</sup>	kualitas_x(1)	1.945	.544	12.768	1	.000	6.993	2.406	20.324
	kmplks_x(1)	3.171	.685	21.450	1	.000	23.828	6.228	91.173
	pmnftn_x(1)	1.453	.672	4.671	1	.031	4.274	1.145	15.958
	imun_x(1)	19.765	8625.578	.000	1	.998	3.836E8	.000	.
	rwyt_x(1)	2.233	1.328	2.826	1	.093	9.327	.690	126.009
	Constant	-3.911	.736	28.225	1	.000	.020		
Step 2 <sup>a</sup>	kualitas_x(1)	2.158	.536	16.230	1	.000	8.657	3.029	24.739
	kmplks_x(1)	3.410	.678	25.279	1	.000	30.251	8.008	114.277
	pmnftn_x(1)	1.783	.643	7.704	1	.006	5.950	1.689	20.962
	rwyt_x(1)	2.576	1.203	4.582	1	.032	13.143	1.243	138.984
	Constant	-4.094	.748	29.967	1	.000	.017		

a. Variable(s) entered on step 1: kualitas\_x, kmplks\_x, pmnftn\_x, imun\_x, rwyt\_x.



**Variables not in the Equation<sup>b</sup>**

			Score	df	Sig.
Step 2 <sup>a</sup>	Variables	imun_x(1)	4.451	1	.035
Overall Statistics			4.451	1	.035

a. Variable(s) removed on step 2: imun\_x.

b. Adding the most significant variable will result in a model which duplicates a prior model.

## HASIL ANALISIS DATA UNIVARIAT

### Crosstabs

**kelompok umur \* kematian neonatal Crosstabulation**

			kematian neonatal		Total
			kontrol	kasus	
kelompok umur	17-27 th	Count	33	30	63
		% within kematian neonatal	47.1%	44.1%	45.7%
	28-38 th	Count	32	32	64
		% within kematian neonatal	45.7%	47.1%	46.4%
	39-49 th	Count	5	6	11
		% within kematian neonatal	7.1%	8.8%	8.0%
Total		Count	70	68	138
		% within kematian neonatal	100.0%	100.0%	100.0%

**pendidikan responden (ibu) \* kematian neonatal Crosstabulation**

			kematian neonatal		Total
			kontrol	kasus	
pendidikan responden (ibu)	tidak sekolah	Count	0	6	6
		% within kematian neonatal	.0%	8.8%	4.3%
	SD	Count	1	4	5
		% within kematian neonatal	1.4%	5.9%	3.6%
	SLTP	Count	8	16	24
		% within kematian neonatal	11.4%	23.5%	17.4%
	SLTA	Count	34	32	66
		% within kematian neonatal	48.6%	47.1%	47.8%
	Diploma/PT	Count	27	10	37
		% within kematian neonatal	38.6%	14.7%	26.8%
Total		Count	70	68	138
		% within kematian neonatal	100.0%	100.0%	100.0%

**pekerjaan responden (ibu) \* kematian neonatal Crosstabulation**

			kematian neonatal		Total
			kontrol	kasus	
pekerjaan responden (ibu)	PNS / Peg.Swasta	Count	27	15	42
		% within kematian neonatal	38.6%	22.1%	30.4%
	wiraswasta	Count	15	11	26
		% within kematian neonatal	21.4%	16.2%	18.8%
	tidak bekerja	Count	28	42	70
		% within kematian neonatal	40.0%	61.8%	50.7%
Total		Count	70	68	138
		% within kematian neonatal	100.0%	100.0%	100.0%

**urutan lahir anak \* kematian neonatal Crosstabulation**

			kematian neonatal		Total
			kontrol	kasus	
urutan lahir anak	1-3	Count	64	60	124
		% within kematian neonatal	91.4%	88.2%	89.9%
	4-6	Count	6	8	14
		% within kematian neonatal	8.6%	11.8%	10.1%
Total		Count	70	68	138
		% within kematian neonatal	100.0%	100.0%	100.0%