

DAFTAR PUSTAKA

- Abbas A.K., Lichtman A.H. (2005). *Cellular and Molecular Immunology*. 5th Edition.
- Bisri T. (2012) neuroanesthesia dan critical care cedera otak traumatik. Bandung: *Saga Olahcitra*, 229-40. Terapi nutrisi pada pasien cedera kepala berat.
- Campos. Machado F.S. (2012). Nutrition therapy in severe head trauma patients. *Bras Ter intensive*, 24(1): 97-105.
- Cely C.M., Arora P., Quartin A.A., Kett D.H., Schein R.M. (2004). Relationship of baseline glucose homeostasis to hyperglycemia during medical critical illness. *Chest*, 126:879-87.
- Cook A.M., Peppard A., Magnuson B. (2008). Nutrition consideration in traumatic injury. *Nutrition in clinical practice*, 23(6):608-17
- Debora Y., Villyastuti Y.W., Harahap M.S. (2009). Nutrisi pada pasien cedera kepala. *J Anestesiologi Indonesia*, 1(1):56-64.
- Egi M., Bellomo R. (2009). Reducing glycemic variability in intensive care unit patients : a new therapeutic target?. *Journal of Diabetes Science and Technology*, 3(6):1302-8.
- Gupta A.K., Summors A. (2001). Nutrition in the neurocritical care unit. In: Notes in neuroanesthesia and critical care. *Green wich medical media London*, 212-3.

- Godoy D.A., Napoli M.D., Biestro A., Leinhardt R. (2011). Perioperative glukosa control in neurosurgical patients. *Anesthesiology Research and Practice*; 31 Juli 2011 [diunduh 21 Oktober 2011] + [sekitar 13 halaman]. Tersedia dari Hindawi Publishing Corporation.
- Herman M.A., Kahn B.B. (2006). Glucose transport and sensing in the maintenance of glucose hemoostasis and metabolic harmony. *J Clin Invest*, 116:1767-75.
- Harrison, M.J.G.. (1984). *Head injury*. In *Contemporary Neurology*, Butterwoths, London, 453-462.
- Hsu C.W. (2012). Glycemic control in critically ill patients. *World J Crit Care Med*, 1(1):31-39.
- Inzucchi S.E. (2006). Clinical practise. Management of hyperglycaemia in the hospital setting. *N Engl J Med*, 355:1903-11.
- Jeremitsky E., Omer L.A., Dunham C.M., Wilberger J., Rodriguez A. (2005). *The Impact of Hyperglycemia on Patient with Severe Brain Injury*. Departement of Surgery, Allegheny Hospital Pittsburgh USA, 47-50.
- Khosravani H, et al. (2009). Occurrence and Adverse Effect on Outcome of Hyperlactatemia in the Critically Ill. *Critical Care*, 13:R90.
- Lucena A.F., Tiburcio R.V., Vasconselos G.C., Ximenes J.D.A., Filho G.C. (2011). Influence of acute brain injuries on gut motility. *Bras ter Intensiva*, 23(1):96-103.

- Maurya I., Pawar M., Garg R., Kaur M., Sood R. (2011). Comparison of respiratory quotient and resting energy expenditure in two regimens of enteral feeding-continuous vs. *intermittent in head-injured critically ill patients*. *Saudi J Anaesth*, 5(2):195-201.
- Magistretti P.J., Pellerin L. (1999). Astrocytes couple synaptic activity to glucose utilization in the brain. *News Physiol Sci*, 14 (10) : 177-82.
- Prins A. (2010). Glucose : the worst of all evils. *S Afr J Clin Nutr*, 23:50-4.
- Pasternak J.J., McGregor D.G., Schroeder D.R., Lanier W.L., Shi Q., Hindman B.J., et al. (2008). Hyperglycemia in patients undergoing cerebral aneurysm surgery: its association with longterm gross neurologic and neuropsychological function. *Mayo Clin Proc*, 83:406-17.
- Rovliast A., Kotsou S. (2000). *The Influence of Hyperglycemia on Neurological Outcome in Patients with Severe Head Injury*. *Neurosurgery*, 46(2):335.
- Roberts P.R. (2001). Approach to nutritional support. In: Apostolakos MJ, Papadakos PJ, editor. *The Intensive Care Manual*, New York: McGraw-Hill Medical Publishing Division: 169-87.
- Slone D.S. (2004). Nutritional support of the critically ill and injured patient. *Critical care clinics*, 20:135-57.
- Serpa L.F., Kimura M., Faintuch J., Coconello I. (2003). Effects of continuous versus bolus infusion of enteral nutrition in critical patients. *HOSP. CLIN.FAC.MED.S.PAULO*, 58(1):9-14.

- Stahel P.F., Flierl M.A., Moore E.E. (2010). "Metabolic staging" after major trauma-a guide for clinical decision making ?. *Scandinavian journal of trauma, resuscitation and emergency medicine*, 18:34.
- Signorini D.F., Andrews P., J.D., Jone P.A., Mardlaw J.M., Miller J.D. (1999). *Predicting Survival Using Simple Clinical Variables: a Case Study in Traumatic Brain Injury*. *Neurology Neurosurgery Psychiatry*, vol 66, 20-5.
- Stroud M., Duncan H., Nightingale J. (2003). Guidelines for enteral feeding in adult hospital patients. *Gut*, 52(7): 1-12 [diunduh February 2013]. Tersedia dari www.gutjnl.com.
- Sengupta G., Guha A., Rudra A., Maitra G., Kumar P., Roy K. (2008). *Indian Journal of Anaesthesia*, 52(1):23-27.
- Turina M., Miller F.N., Tucker C.F., Polk H.C. (2006). Short-term hyperglycemia in surgical patients and a study of related cellular mechanism. *Ann Surg*, 243:845-51.
- The DAA (2011). Nutrition Support Interest Group. Enteral nutrition manual for adults in health care facilities, 5-18.
- Vespa P., et al. (2005). *Metabolic Crisis without Brain Ischemia is Common after Traumatic Brain Injury: a Combine microdialysis and PET Study*. *Journal of Cerebral Blood Flow & Metabolism*, 27:763-74.
- Van den Bergh G., Schetz M., Vlaaselaers D., Hermans G., Wlimer A., Bouillon R., et al. (2009). Intensive insulin therapy in critically ill

patients : nice sugar or leuon blood glucose target. *J Clin Endocrinol Metab*, 94:3163-70.

Weisman C. (2009). Nutrition and metabolic control. In: Miller RD. Eriksson LI. Fleisher LA. Wiener JP. Young WL, editors. *Miller's anesthesia*. 7th ed. Philadelphia: Elsevier, 390-3.

Young B., Ott L., Dempsey R., Haack D., Tibbs P. (1989). *Relationship Between Admission Hyperglycemia and Neurologic Outcome of Severely brain-injured Patient*. Division of Neurosurgery, University of Kentucky Medical Center, 466-72.

LAMPIRAN 1**PERSETUJUAN SEBELUM PERSIAPAN DIMULAI****PENGARUH PEMBERIAN NUTRISI ENTERAL INTERMITTEN TERHADAP KADAR GULA DARAH SEWAKTU PADA PASIEN CEDERA OTAK BERAT PASCABEDAH**

Yang bertandatangan di bawah ini :

Nama/Umur :

Ala m a t :

No. Rekam Medis :

Menyatakan dengan sesungguhnya bahwa saya telah mendapatkan penjelasan dan kesempatan bertanya hal-hal yang belum saya mengerti tentang penelitian ini. Penjelasan tersebut meliputi manfaat dan keuntungan serta efek samping dari pemberian nutrisi enteral intermiten terhadap kadar gula darah sewaktu pada pasien cedera otak berat pascabedah.

Efek samping yang mungkin ditimbulkan dari pemberian nutrisi ini adalah peningkatan kadar gula darah sewaktu. Bila terjadi hal demikian peneliti akan memberikan obat-obatan dan melakukan tindakan untuk menangani efek samping tersebut. Namun, secara teoritis, perlakuan pemberian obat ini memiliki efek samping yang minimal.

Setelah mendapat penjelasan tersebut, dengan ini saya menyatakan secara sukarela ikut serta dalam penelitian ini dan saya/keluarga saya berhak mengundurkan diri, bila ada alasan sehubungan dengan kesehatan saya.

Demikian pernyataan ini saya buat dengan sebenarnya, dengan penuh kesadaran dan tanpa paksaan

Makassar, 2013

Saksi

Yang menyatakan

.....
**Penanggung Jawab Medik,
dr. Hisbullah SpAn, KIC-KKAV**
Alamat : Jl. Perdos Makassar
HP : 085255958512

.....
**Penanggung Jawab Penelitian,
dr. Julia Hasir**
Alamat : Jl Letnan Jenderal Andi
Mappaoudang No. 81 Makassar
HP : 081342191718

LAMPIRAN 2**LEMBAR PENGAMBILAN DATA PENELITIAN**

| | |
|-----------------|-------------------------|
| Nama pasien : | No Rekam Medis : |
| Jenis Kelamin : | BB : Kg |
| Umur : thn | TB : Cm |
| Pendidikan : | BMI : Kg/m ² |
| Alamat : | No.Sampel : |
| Pekerjaan : | |

Data Klinis

1. Diagnosis Prabedah :
2. ASA PS :
3. Tanda Vital : TD = mmHg N = x/mnt
P = x/mnt S = °C
VAS =
4. Tindakan :
5. Mulai Anestesi :
6. Mulai Operasi :
7. Jumlah Perdarahan :
8. Jumlah Kalori :

GULA DARAH SEWAKTU 50%

| | 4 Jam | 8 Jam | 12 Jam | 16 Jam | 20 Jam | 24 Jam |
|-------------|-------|-------|--------|--------|--------|--------|
| GDS | | | | | | |
| Hemodinamik | | | | | | |

GULA DARAH SEWAKTU 75%

| | 4 Jam | 8 Jam | 12 Jam | 16 Jam | 20 Jam | 24 Jam |
|-------------|-------|-------|--------|--------|--------|--------|
| GDS | | | | | | |
| Hemodinamik | | | | | | |

GULA DARAH SEWAKTU 100%

| | 4 Jam | 8 Jam | 12 Jam | 16 Jam | 20 Jam | 24 Jam |
|-------------|-------|-------|--------|--------|--------|--------|
| GDS | | | | | | |
| Hemodinamik | | | | | | |

LAMPIRAN 3**CASE REPORT FORM**

Tanggal :

No. Urut :

I. Identitas

Nama/Umur : (Inisial)/tahun

Jenis Kelamin :

BB/TB : kg/.....cm.

Diagnosis :

II. Anamnesis

.....

III. Pemeriksaan Fisis

Tanda Vital :

Tekanan Darah : mmHg Pernapasan :x/mnt BB :.....kg

Denyut Nadi :x/mnt Suhu :⁰C TB :.....cm

Kepala :

Leher :

Thoraks :

Abdomen :

Ekstremitas :

IV. Pemeriksaan Penunjang

Laboratorium :

- Darah rutin
- Urine rutin
- Kimia darah

Pemeriksaan penunjang lainnya :

- Foto Thorax
- EKG

V. Diagnosis

.....

VI. Kesimpulan

LAMPIRAN 4**ADVERSE EVENT FORM****Identitas**

Nama (Inisial) / Umur :

No. MR :

Diagnosis :

Adverse event**KEBUTUHAN KALORI 50%**

| | 4 Jam | 8 Jam | 12 Jam | 16 Jam | 20 Jam | 24 Jam |
|-----------------------|--------------|--------------|---------------|---------------|---------------|---------------|
| Perubahan GDS | | | | | | |
| Perubahan Hemodinamik | | | | | | |