

## DAFTAR PUSTAKA

- Adams, W.J. dan Anramovic, J. 1991. Wound infiltration with 0,25% bupivacaine not effective for post operative analgesia after cholecystectomy. *Aust N Z J Surg*, 61(8):626-30.
- Alexander, J.L. 1997. Pain after laparoscopy. *Br J Anaesth*, 79:369-378.
- Biswas, B.K. dan Bithal, P.K. 2003. Preincision 0.25% bupivacaine scalp infiltration and postcraniotomy pain: a randomized double-blind, placebo-controlled study. *J Neurosurg Anesthesiol*.15(3):234-9.
- Benitez, M.A. et al. 2011. A strategy for conversion from subcutaneous to oral ketamine in cancer pain patients : Effect of a 1:1 ratio. *J Pain Symptom manage*, 41(6) :1098-105.
- Bourne, E. et al. 2010. A review of local anesthetic cardiotoxicity and treatment with lipid emulsion. *Local Reg Anesth*, 3:11-9.
- Carlton, S.M. and Coggeshall, R.E. 1999. Inflammation-induced changes in peripheral glutamate receptor populations. *Brain Res*, 820:63–70.
- Carlton, S.M. et al. 1995. Localization and activation of glutamate receptors in unmyelinated axons of rat glabrous skin. *Neurosci Lett*, 197:25–8.
- Cousins, M.J. et al. 2000. Labat lecture: Pain-a persistent problem. *Reg Anesth Pain Med*, 25:6–21.
- Cobby, T.F. dan Read, M.F. 1997. Wound infiltration with local anaesthetic after abdominal hysterectomy. *Br J Anaesth*, 78(4):431-2.
- Dickenson, A.H. 1995. Spinal cord pharmacology of pain. *Br J Anaesth*. 75:193–200.
- Du, J. et al. 2001. Glutamate-induced excitation and sensitization of nociceptors in rat glabrous skin. *J Pain*, 89:187–98.
- Desjardins P.J, et al. 2001. The injectable cyclooxygenase-2–specific inhibitor parecoxib sodium has analgesic efficacy when administered preoperatively. *Int Anesth Res Soc*, 93:721–7.

- Ejlerson, E. et al. 1992. A comparison between preincisional and postincisional lidocaine infiltration and postoperative pain. *Anaesth analg*, 74(4):495-8.
- Gottschalk, A. dan Smith, D.S. 2001. New concept in acute pain therapy : preemptive analgesia. *Am Fam Phycian*, 63(10):1979-85.
- Hanibal, K. et al. 1996. Preoperative wound infiltration with bupivacain reduces early and late opioid requirement after hysterectomy. *Anaesth Analg*, 83:376-81
- Hudspith, M.J.M. 1997. Glutamate: A role in normal brain function, anaesthesia, analgesia and CNS injury. *Br J Anaesth*, 78:731–47.
- Heavner, J.E. 2008. Pharmacology of local anesthetics. In: Longnecker DE, Brown DL, Newman MF, Zapol WM, editors. *Anesthesiology*. New York: McGraw-Hill, p. 981-98.
- Honarmand, A. Et al. 2012. Preinsisional administration of intravenous or subcutanteus infiltration low dose ketamin supresses post operative pain after appendictomy. *J pain Res*, 5:1-6.
- Javid, M. et al. 2012. Evaluation of a low dose ketamin in post tonsilectomy pain relief : A randomized trial company intravenous and subcutaneous ketamin in pediatrics. *Anesthesiology and pain medicine*, 2:85-89.
- Jackson, S.A. et al. 1996. Does postlaparoscopy pain relate to residual carbon dioxide? *Anaesthesia*, 51:485–7.
- Jin, Y.H. et al. 2012. Peripheral glutamat receptors are required for hyperalgesia induced by capsaicin. *Pain Res Treat*, 12:91
- Joris, J, et al. 1992. Metabolic and respiratory changes after cholecystectomyperformed via laparotomy or laparoscopy. *Br J Anaesth*, 69:341–5.
- Katzung, B.G. 1997. *Farmakologi dasar dan klinik*, edisi 6. Jakarta: EGC, p. 56-60.
- Kelly, D.J. et al. 2001. Preemptive Analgesia I : physiological pathwys and pharmacological modalities. *Can J Anesth*, 48;10:1000-10
- Kongara, K. 2008. *Studies on Renal Safety and Preventive Analgesic Efficacy of Tramadol and Parecoxib in Dogs*. Doctoral thesis. Massey University. Palmerston North, New Zealand.

- Kleinman,W. dan Mikhail, M. 2008. Regional Anesthesia & Pain Management in Morgan GE, Mikhail MS, Murray MJ editor. Clinical Anesthesiology 4<sup>th</sup>ed. McGraw-Hill, p.289-323.
- Klein, J.R. et al. 2000. Infiltration of the abdominal wall with local anaesthetic after total abdominal hysterectomy has no opioid sparing effect. *Br J Anaesth*, 84(2):248-9.
- Lawand, N.B. et al. 1997. Excitatory amino acid receptor involvement in peripheral nociceptive transmission in rats. *Eur J Pharm*, 324:169–77.
- Lalenoh, H. 2009. Pregabalin dan Gabapentin sebagai analgesia preemptif. *Majalah Anestesia dan Critical Care*, 27: p.185-9.
- Lin, C. dan Durieux, M.E. 2005. Ketamine and kids: an update. *Pediatric Anesthesia*, 15: p. 91-97.
- Morgan, C.J.A. dan Curran, H.V. 2012. Ketamin use. *Clinical Health Psychology*, University College London, London, vol 10, p 27-38.
- Moiniche, S. et al. 1998. A Qualitative systematic review of incisional local anesthesia for postoperative pain relief after abdominal operations. *Br J Anaesth*, 81:377-83
- Moiniche, S. et al. 2000. Local anesthetic infiltration for postoperative pain relief after laparoscopy: A qualitative and quantitative systematic review of intraperitoneal, port-site infiltration and mesosalpinx block. *J Anesth Analg*, 90:899–912
- Omote, K. et al. 1998. Formalin-induced release of excitatory amino acids in the skin of the rat hindpaw. *Brain Res*, 787:161–4.
- Priya, V. et al. 2002 Efficacy of intravenous ketoprofen for pre-emptive analgesia. *J Post Grad Med*, 48:109–12.
- Portera, S.G. et al. 1998. A randomized, double- blinded trial of preemptive analgesia in laparoscopy. *Obstet Gynecol*. 92(6):972-5.
- Rogawski, M.A. 1993. Therapeutic potential of excitatory aminoacid antagonists, channel blocks and 2, 3 benzodiazepines. *Trends Pharmacol Sci*, 14:325–31.

- Sajjad, A. et al. 2004. Preemptive analgesia: effect of low dose ketamine as preemptive analgesia in post operative pain management after lower abdominal surgery. *Journal of BSA*, 17(1):17-22.
- Safavi, M. et al. 2011. Preincisional analgesia with intravenous infiltration of ketamin reduces post operative pain in patients after open cholecystectomy : a randomized, double blind, placebo-controlled study. *Pain Med*, 12:1418-26.
- Safavi, M. et al. 1979. Pentazocine and morphine. *Br J Anaesth*, 51:497–502.
- Stubhaug, A. 1997. A new method to evaluate central sensitization to pain following surgery. Effect of ketamine. *Acta Anaesthesiol Scand Suppl* 110:154–5.
- Salermo, A. dan Herman, R. 2006. Efficacy and safety of steroid use for post operative pain relief. *J Bone Joint Surg Am*, 88:1361-72.
- Sonbatty, E.L. et al. 2006. Preemptive peritonsillar ketamine infiltration: postoperative analgesic efficacy versus meperidine. *Middle East J Anesthesiol*, 21(1):43-51.
- Tan, P.H. et al. 2007. Preincisional subcutaneous infiltration of ketamine suppresses postoperative pain after circumcision surgery. *Clin J Pain*, 23(3):214–8.
- Tverskoy, M. et al. 1996. Ketamin enhances lokal anesthetic and analgesic effects of bupivacaine by peripheral mechanism: a study in postoperative patients. *Neurosci Lett*, 215(1):5-8
- Vasan, N.R. et al. 2002. Preincisional bupivacaine in posttonsilectomy pain relief: a randomized prospectif study. *Arch Otolaryngeal Head Neck Surg*, 128(2):145-9.
- Woolf, C.J. dan Chong, M. 1993. Preemptive analgesia-treating postoperative pain by preventing the establishment of central sensitization. *Anesth Analg*, 77:352–79.
- Wikipedia, 2008. The Free Encyclopedia, Ketamine. Available at: <http://en.wikipedia.org>. December.
- Zahn, P.K. dan Brennan, T.J. 1998. Intrathecal metabotropic glutamate receptor antagonists do not decrease mechanical hyperalgesia in a rat model of postoperative pain. *Anesth Analg*, 6:1354–9.

## Lampiran 1

**TABEL PENENTUAN JUMLAH SAMPEL DARI POPULASI  
TERTENTU DENGAN TARAF KESALAHAN 1, 5, DAN 10%**

N	S			N	S			N	S		
	1%	5%	10%		1%	5%	10%		1%	5%	10%
10	10	10	10	280	197	115	138	2800	537	310	247
15	15	14	14	290	202	158	140	3000	543	312	248
20	19	19	19	300	207	161	143	3500	558	317	251
25	24	23	23	320	216	167	147	4000	569	320	254
30	29	28	27	340	225	172	151	4500	578	323	255
35	33	32	31	360	234	177	155	5000	586	326	257
40	38	36	35	380	242	182	158	6000	598	329	259
45	42	40	39	400	250	186	162	7000	606	332	261
50	47	44	42	420	257	191	165	8000	613	334	263
55	51	48	46	440	265	195	168	9000	618	335	263
60	55	51	49	460	272	198	171	10000	622	336	263
65	59	55	53	480	279	202	173	15000	635	340	266
70	63	58	56	500	285	205	176	20000	642	342	267
80	71	65	62	600	315	221	187	40000	563	345	269
85	75	68	65	650	329	227	191	50000	655	346	269
90	79	72	68	700	341	233	195	75000	658	346	270
95	83	75	71	750	352	238	199	100000	659	347	270
100	87	78	73	800	363	243	202	150000	661	347	270
110	94	84	78	850	373	247	205	200000	661	347	270
120	102	89	83	900	382	251	208	250000	662	348	270
130	109	95	88	950	391	255	211	300000	662	348	270
140	116	100	92	1000	399	258	213	350000	662	348	270
150	122	105	97	1050	414	265	217	400000	662	348	270
160	129	110	101	1100	427	270	221	450000	663	348	270
170	135	114	105	1200	440	275	224	500000	663	348	270
180	142	119	108	1300	450	279	227	550000	663	348	270
190	148	123	112	1400	460	283	229	600000	663	348	270
200	154	127	115	1500	469	286	232	650000	663	348	270
210	160	131	118	1600	477	289	234	700000	663	348	270
220	165	135	122	1700	485	292	235	750000	663	348	271
230	171	139	125	1800	492	294	237	800000	663	348	271
240	176	142	127	1900	498	297	238	850000	663	348	271
250	182	146	130	2000	510	301	241	900000	663	348	271
260	187	149	133	2200	520	304	243	950000	663	348	271
270	192	152	135	2600	529	307	245	1000000	664	349	272

## LAMPIRAN 2

### PERNYATAAN PERSETUJUAN PASIEN

Yang bertandatangan dibawah ini :

Nama/Umur :

A l a m a t :

No. Rekam Medis :

Menyatakan dengan sesungguhnya bahwa saya telah mendapatkan penjelasan dan kesempatan bertanya hal-hal yang belum saya mengerti tentang penelitian ini. Penjelasan tersebut meliputi manfaat dan keuntungan tehnik infiltrasi preinsisi dengan ketamin dan bupivakain serta efek sampingnyayang akan saya dapatkan selama penelitian ini. Efek samping yang paling mungkin ditimbulkan adalah nyeri. Peneliti akan melakukan tindakan terhadap efek samping tersebut dan memberikan obat sebagai penanganannya.

Setelah mendapat penjelasan tersebut, dengan ini saya menyatakan secara sukarela ikut serta dalam penelitian ini dan saya berhak mengundurkan diri bila ada alasan sehubungan dengan kesehatan saya. Demikian pula jika terjadi perselisihan saya akan melakukan musyawarah dengan peneliti untuk mencari jalan keluar yang terbaik tentang perselisihan tersebut.

Demikian pernyataan ini saya buat dengan sebenarnya, dengan penuh kesadaran dan tanpa paksaan.

Makassar, Maret 2013

Saksi,

Tanda Tangan

Yang Menyatakan,

1. \_\_\_\_\_

2. \_\_\_\_\_

(\_\_\_\_\_)

**Penanggung Jawab Medik,**

**Penanggung Jawab Penelitian,**

**dr. A. Salahuddin, SpAn**

**dr. Lismasari**

Telp.04115026635

Telp. 082189183067

Perumahan Griya prima tonasa

Jl. NTI FG. 5, Makassar.

Blok B II/1, Makassar.

### Lampiran 3

#### LEMBAR PENGAMBILAN DATA PENELITIAN

---

Nama pasien	:		BB	:	Kg
Umur	:	thn	TB	:	cm
Pendidikan	:		BMI	:	Kg/m <sup>2</sup>
Alamat	:		No.Sampel	:	
Pekerjaan	:				
No Rekam Medis	:				

---

#### Data Klinis

1. Diagnosis MRS :
2. ASA PS :
3. Mulai Anestesi infiltrasi :
4. Mulai operasi/insisi :
5. Mulai insersi trokar :
6. Mulai insuflasi :
7. Selesai Operasi :





## Lampiran 4

### LEMBAR PENGAMATAN

#### I. Pengamatan Respon Hemodinamik

Penilaian	TAR				LJ			
	T0	T1	T2	T3	T0	T1	T2	T3
KI <sub>p</sub>								
Ketamin 0,3 mg/kgBB								
Bupivacain 0,25%								
NaCl 0,9%								

Ket : T0 : nilai basal, T1 : insisi, T2 : insersi trokar, T3 : insuflasi CO<sub>2</sub>,

TAR : tekanan arteri rerata, LJ : Laju jantung



## II. Pengamatan Waktu *Rescue Analgesia*

Kelompok
WRA(Jam)
$\leq 1$
$\leq 2$
$\leq 3$
$\leq 4$
$\leq 5$
$\leq 6$
$> 6$

Ket : WRA : Waktu *Rescue Analgesia*

