

## DAFTAR PUSTAKA

1. KDIGO 2024 Clinical Practice Guideline for the Evaluation and Management of Chronic Kidney Disease. *Kidney Int* 2024;105:A1.
2. Bikbov B, Purcell C, Levey AS, Smith M, Abdoli A, Abebe M, et al. Global, regional, and national burden of chronic kidney disease, 1990–2017: a systematic analysis for the Global Burden of Disease Study 2017. *Lancet* 2020;395:709–33.
3. World Health Organization. The top 10 causes of death 2023. <https://www.who.int/news-room/fact-sheets/detail/the-top-10-causes-of-death>
4. Jager KJ, Kovesdy C, Langham R, Rosenberg M, Jha V, Zoccali C. A single number for advocacy and communication—worldwide more than 850 million individuals have kidney diseases. *Nephrol Dial Transplant*. 2019;34(11):1803-1805.
5. Goldstein-Fuchs J, LaPierre AF. Nutrition and Kidney Disease. In: Gilbert SJ, Weiner DE, editors. *National Kidney Foundation Primer on Kidney Diseases*. 6th ed. Philadelphia: W.B. Saunders; 2014. p. 467-75.
6. Baragetti I, De Simone I, Biazzi C, Buzzzi L, Ferrario F, Luise MC, et al. The low-protein diet for chronic kidney disease: 8 years of clinical experience in a nephrology ward. *Clin Kidney J*. 2020;13(5):253-260.
7. Nallu A, Sharma S, Ramezani A, Muralidharan J, Raj D. Gut microbiome in chronic kidney disease: challenges and opportunities. *Transl Res* 2017;179:24–37.
8. Mocanu C-A, Simionescu TP, Mocanu AE, Garneata L. Plant-Based versus Animal-Based Low Protein Diets in the Management of Chronic Kidney Disease. *Nutrients* 2021;13(11):3721.
9. Wang S, Lv D, Jiang S, Jiang J, Liang M, Hou F, et al. Quantitative reduction in short-chain fatty acids, especially butyrate, contributes to the progression of chronic kidney disease. *Clin Sci* 2019;133:1857–70.
10. Li Y, Chen X, Kwan TK, Loh Y, Singer J, Liu Y, et al. Dietary Fiber Protects against Diabetic Nephropathy through Short-Chain Fatty Acid-Mediated Activation of G Protein-Coupled Receptors GPR43 and GPR109A. *J Am Soc Nephrol*. 2020;31(6):1267-1281
11. Adachi K, Wakino S, Itoh H. PS-C23-12: Dietary intervention with the processed low protein genmai (brown rice) against chronic kidney disease. *J Hypertens* 2023;41:e407.
12. Watanabe S, Ohtsubo K. Low-Protein Diet: History and Use of Processed Low-Protein Rice for the Treatment of Chronic Kidney Disease. *Foods* 2021;10(10):2255
13. Watanabe S, Adachi K, Wakino S, Hisada T, Mochizuki J. Dietary therapy with low protein genmai (brown rice) to improve the gut-kidney axis and reduce CKD progression. *Asia Pac J Clin Nutr*. 2022;31(3):341–7.



Kesehatan RI. Pedoman Nasional Pelayanan Kedokteran Tata Laksana njal Kronik 2023. <https://p2ptm.kemkes.go.id>

Wilmer WA, Falkenhain ME, Ladson-Wofford SE, Nahman NS, Rovin BH. Ion: One or many therapies? *Kidney Int* 2001;59:1211–26.

16. Ko GJ, Obi Y, Tortorici AR, Kalantar-Zadeh K. Dietary protein intake and chronic kidney disease. *Curr Opin Clin Nutr Metab Care* 2017;20:77–85.
17. Bellizzi V. Low-protein diet or nutritional therapy in chronic kidney disease? *Blood Purif* (2013) 36 (1): 41–46.
18. Garneata L, Stancu A, Dragomir D, Stefan G, Mircescu G. Ketoanalogue-Supplemented Vegetarian Very Low-Protein Diet and CKD Progression. *J Am Soc Nephrol*. 2016;27(7):2164-2176.
19. Hahn D, Hodson EM, Fouque D. Low protein diets for non-diabetic adults with chronic kidney disease. *Cochrane Database Syst Rev*. 2020;10:CD001892.
20. Goraya N, Wesson DE. Dietary interventions to improve outcomes in chronic kidney disease: *Curr Opin Nephrol Hypertens*. 2015;24(6):505-510.
21. St-Jules DE, Woolf K, Pompeii ML, Kalantar-Zadeh K, Sevick MA. Reexamining the Phosphorus-Protein Dilemma: Does Phosphorus Restriction Compromise Protein Status? *J Ren Nutr*. 2016;26(3):136-140.
22. Tong Y, Guo S, Li T, Yang K, Gao W, Peng F, et al. Gut microbiota and renal fibrosis. *Life Sci*. 2024;357:123072.
23. Corte-Iglesias V, Saiz ML, Andrade-Lopez AC, Salazar N, Bernet CR, Martin-Martin C, et al. Propionate and butyrate counteract renal damage and progression to chronic kidney disease. *Nephrol Dial Transplant*. 2025;40:133-150.
24. Huang W, Man Y, Gao C, Zhou L, Gu J, Xu H, et al. Short-Chain Fatty Acids Ameliorate Diabetic Nephropathy via GPR43-Mediated Inhibition of Oxidative Stress and NF-κB Signaling. *Oxid Med Cell Longev*. 2020;4074832.
25. Lu P-C, Hsu C-N, Lin I, Lo M, Yang M-Y, Tain Y. The Association Between Changes in Plasma Short-Chain Fatty Acid Concentrations and Hypertension in Children With Chronic Kidney Disease. *Front Pediatr*. 2021;8:613641
26. Sun Y, Zhou C, Chen Y, He X, Gao F, Xue D. Quantitative increase in short-chain fatty acids, especially butyrate protects kidney from ischemia/reperfusion injury. *J Investig Med*. 2021;70:29-35.
27. Li Y, Ma J, Loh Y, Chadban S, Wu H. Short-chain fatty acids directly exert anti-inflammatory responses in podocytes and tubular epithelial cells exposed to high glucose. *Front Cell Dev Biol*. 2023;11:1182570.
28. Tilves C, Yeh H, Maruthur N, Juraschek S, Miller ER, White K, et al. Increases in Circulating and Fecal Butyrate are Associated With Reduced Blood Pressure and Hypertension: Results From the SPIRIT Trial. *J Am Heart Assoc*. 2022;11(8):e024763.
29. Onyszkiewicz M, Gawrys-Kopczynska M, Sałagaj M, Aleksandrowicz M, Sawicka A, Koźniewska E, et al. Valeric acid lowers arterial blood pressure in rats. *Eur J Pharmacol* 2020;877:173086.
30. Chen L, He F, Dong Y, Huang Y, Wang C, Harshfield G, et al. Modest Sodium Reduction Increases Circulating Short-Chain Fatty Acids in Untreated Hypertensives. *Hypertension*. 2020;76:73-79.
31. Wang Y, Ma L, Fu P. Roles of short-chain fatty acids in kidney diseases. *Chin Med* (10):1228-1232.



32. Chang Y, Chen Y, Zhou Q, Wang C, Chen L, Di W, et al. Short-chain fatty acids accompanying changes in the gut microbiome contribute to the development of hypertension in patients with preeclampsia. *Clin Sci*. 2020;134(2):289-302.
33. Koppe L, Fouque D, Soulage CO. The Role of Gut Microbiota and Diet on Uremic Retention Solutes Production in the Context of Chronic Kidney Disease. *Toxins*. 2018;10(4):155.
34. Kim SM, Song IH. The clinical impact of gut microbiota in chronic kidney disease. *Korean J Intern Med* 2020;35:1305–16.
35. Vanholder R, Glorieux G. The intestine and the kidneys: a bad marriage can be hazardous. *Clin Kidney J*. 2015;8(2):168-179.
36. Ramezani A, Massy ZA, Meijers B, Evenepoel P, Vanholder R, Raj DS. Role of the Gut Microbiome in Uremia: A Potential Therapeutic Target. *Am J Kidney Dis*. 2016;67(3):483-498.
37. Rysz J, Franczyk B, Ławiński J, Olszewski R, Ciałkowska-Rysz A, Gluba-Brzózka A. The Impact of CKD on Uremic Toxins and Gut Microbiota. *Toxins*. 2021;13(4):252.
38. Guldris SC, Parra EG, Amenós AC. Gut microbiota in chronic kidney disease. *Nefrologia*. 2017;37(1):9-19.
39. Ivarenga L, Reis DCMV, Kemp JA, Teixeira KTR, Fouque D, Mafra D. Using the concept of food as medicine to mitigate inflammation in patients undergoing peritoneal dialysis. *Ther Apher Dial*. 2024;28:341-353.
40. Wang P, Zhang X, Zheng X, Gao J, Shang M, Xu J, et al. Folic Acid Protects against Hyperuricemia in C57BL/6J Mice via Ameliorating Gut–Kidney Axis Dysfunction. *J Agric Food Chem* 2022;70:15787–803.
41. Peters BA, Qi Q, Usyk M, Daviglius ML, Cai J, Franceschini N, et al. Association of the gut microbiome with kidney function and damage in the Hispanic Community Health Study/Study of Latinos (HCHS/SOL). *Gut Microbes* 2023;15:2186685.
42. Watanabe T, Moriya K, Tsubokawa N, Baba H. Effect of remifentanyl on postoperative nausea and vomiting: a randomized pilot study. *J Anesth*. 2018;32(5):781-785.
43. Milovanova LYu, Volkov AV, Milovanova SY, Taranova MV, Nezhdanov KS. Soy Protein as a Part of a Low-protein Diet is a New Direction in Cardio- and Nephroprotection in Patients With 3b-4 Stages of Chronic kidney Disease. *J Ren Nutr*. 2023;33(4):435-442.
44. Srivastava T, Ju W, Milne GL, Rezaiekhaliq MH, Staggs VS, Alon US, et al. Urinary prostaglandin E2 is a biomarker of early adaptive hyperfiltration in solitary functioning kidney. *Prostagl Other Lipid Mediat*. 2020;146:106403.
45. Adeva-Andany MM, Fernández-Fernández C, Funcasta-Calderón R, Ameneiros-Rodríguez E, Adeva-Contreras L, Castro-Quintela E. Insulin Resistance is Associated with Clinical Manifestations of Diabetic Kidney Disease (Glomerular Hyperfiltration, Albuminuria, and Kidney Function Decline). *Curr Diab Rev*. 2022;18:e171121197998.
46. Sasaki G, Sasaki T, Okabayashi Y, Haruhara K, Koike K, et al. Dietary Protein Restriction Improves Single-Nephron Glomerular Filtration Rate. *Nutrients*. 2020;12(9):2549.
47. Gounden V. Novel glomerular filtration markers. *Adv Clin Chem*. 2019;119.

