

**DAFTAR PUSTAKA**

1. Ade RJ, Hamid F, Budu B, Natzir R, Hala Y, Alam G, Agus R, Bahar B, Syukri A, Primaguna MR, Dwiyantri R, Febrianti A, Sabir M, Azhar A, Hatta M. A potential mechanism of Miana (*Coleus scutellarioides*) and quercetin via NF- $\kappa$ B in *Salmonella typhi* infection. *Heliyon*. 2023 Nov 14;9(11):e22327. DOI: <https://doi.org/10.1016/j.heliyon.2023.e22327>. PMID: 38058621; PMCID: PMC10696054.
2. Alves CC, Donadi EA, Giuliatti S. Structural Characterization of the Interaction of Hypoxia Inducible Factor-1 with Its Hypoxia Responsive Element at the -964G > A Variation Site of the HLA-G Promoter Region. *International Journal of Molecular Sciences*. 2021; 22(23):13046. <https://doi.org/10.3390/ijms222313046>
3. Barreto E, Liu Y, Xiang D, Zhang H, Yao H, Wang Y. Hypoxia-Inducible Factor-1: A Potential Target to Treat Acute Lung Injury. *Oxidative Medicine and Cellular Longevity*. 2020, Article ID 8871476, 13. <https://doi.org/10.1155/2020/8871476>
4. Borsi E, Terragna C, Brioli A, Tacchetti P, Martello M, Cavo M. Therapeutic targeting of hypoxia and hypoxia-inducible factor 1 alpha in multiple myeloma. *Transl Res*. 2015 Jun;165(6):641-50. doi: 10.1016/j.trsl.2014.12.001. Epub 2014 Dec 11. PMID: 25553605.
5. Botusan IR, Sunkari VG, Savu O, Catrina AI, Grünler J, Lindberg S, Pereira T, Ylä-Herttua S, Poellinger L, Brismar K, Catrina SB. Stabilization of HIF-1alpha is critical to improve wound healing in diabetic mice. *Proc Natl Acad Sci U S A*. 2008 Dec 9;105(49):19426-31. doi: 10.1073/pnas.0805230105. Epub 2008 Dec 4. PMID: 19057015; PMCID: PMC2614777.
6. Cano Sanchez M, Lancel S, Boulanger E, Neviere R. Targeting Oxidative Stress and Mitochondrial Dysfunction in the Treatment of Impaired Wound Healing: A Systematic Review. *Antioxidants*. 2018; 7(8):98. <https://doi.org/10.3390/antiox7080098>
7. Casse JM, Martrille L, Vignaud JM, et al. Skin wounds vitality markers in forensic pathology: an updated review. *Med Sci Law*. 2016; 56:128–137.
8. Cecchi R. Estimating wound age: looking into the future. *Int J Legal Med*. 2010;124:523–536.

9. Chujia Lin, Guoshu Yin, Miaoqiong Ou, Shaoyu Zheng. The effects of HIF-1 $\alpha$  and VEGF on wound healing in diabetic mice. *Biomedical Research* 2017; 28 (18): 8121-8124.
10. Cyran AM, Zhitkovich A. HIF1, HSF1, and NRF2: Oxidant-Responsive Trio Raising Cellular Defenses and Engaging Immune System. *Chem Res Toxicol.* 2022 Oct 17;35(10):1690-1700. doi: 10.1021/acs.chemrestox.2c00131. Epub 2022 Aug 10. PMID: 35948068; PMCID: PMC9580020.
11. Dahlan,S. Ilmu Kedokteran Forensik Pedoman Bagi Dokter dan Penegak Hukum. Semarang: Badan Penerbit Universitas Diponegoro; 2000: 47-62
12. Devraj G, Beerlage C, Brüne B, Kempf VA. Hypoxia and HIF-1 activation in bacterial infections. *Microbes Infect.* 2017 Mar;19(3):144-156. doi: 10.1016/j.micinf.2016.11.003. Epub 2016 Nov 27. PMID: 27903434.
13. Emilie V, Chiara C, Emilio BM. (2015). DAMPs from Cell Death to New Life. *Frontiers in Immunology.* 6: 422.DOI=10.3389/fimmu.2015.00422
14. Fachri, M., Hatta, M., Massi, M.N. et al. The strong correlation between ADAM33 expression and airway inflammation in chronic obstructive pulmonary disease and candidate for biomarker and treatment of COPD. *Scientific Reports.* 2021; 11:23162. <https://doi.org/10.1038/s41598-021-02615-2>.
15. Fikri LH, As'ad S, Hatta M, Syamsuddin S, Prasetya P, Maddeppungeng M, Aminuddin A, Zainuddin AA, Hasanah I. Hypnotherapy influence on maternal psychological, breastfeeding, mRNA-OXTR expression, and OXTR protein, *American Journal of Clinical Hypnosis.* 2024;1-15. DOI: <https://doi.org/10.1080/00029157.2023.2293288>
16. Fronczek J, Lulf R, Korkmaz HI. Analysis of inflammatory cells and mediators in skin wound biopsies to determine wound age in living subjects in forensic medicine. *Forensic Sci Int.* 2015;247:7–13.
17. Fronczek J, Lulf R, Korkmaz HI, et al. . Analysis of morphological characteristics and expression levels of extracellular matrix proteins in skin wounds to determine wound age in living subjects in forensic medicine. *Forensic Sci Int.* 2015;246:86–91.
18. Gauchotte G, Wissler MP, Casse JM, et al. . FVIIIra, CD15, and tryptase performance in the diagnosis of skin stab wound vitality in forensic pathology. *Int J Legal Med.* 2013;127:957–965.

19. Goova MT, Li J, Kislinger T, Qu W, Lu Y, Bucciarelli LG, Nowygrod S, Wolf BM, Caliste X, Yan SF, Stern DM, Schmidt AM. Blockade of receptor for advanced glycation end products restores effective wound healing in diabetic mice. *Am J Pathol* 2001;159:513-25.
20. Greijer AE, van der Wall E. The role of hypoxia inducible factor 1 (HIF-1) in hypoxia induced apoptosis. *Journal of Clinical Pathology* 2004; 57:1009-1014.
21. Grellner W, Madea B. Demands on scientific studies: vitality of wounds and wound age estimation. *Forensic Sci Int.* 2007;165:150–154.
22. Gunawan I, Hatta M, Benyamin AF, Islam AA. The Hypoxic Response Expression as a Survival Biomarkers in Treatment-Naive Advanced Breast Cancer. *Asian Pac J Cancer Prev.* 2020 Mar 1;21(3):629-637. doi: <https://doi.or.10.31557/APJCP.2020.21.3.629>. PMID: 32212787; PMCID: PMC7437329.
23. Hatta, M., Surachmanto, E.E., Islam, A.A., Wahid, S. Expression of mRNA IL-17F and sIL-17F in atopic asthma patients. *BMC Research Notes*, 2017;10 (1), art. no. 202, DOI: <https://doi.org.10.1186/s13104-017-2517-9>
24. Hoediyanto H. *Buku Ajar Ilmu Kedokteran Forensik dan Medikolegal*. Surabaya: Badan Penerbit FK UNAIR; 2012
25. Hong WX, Hu MS, Esquivel M, Liang GY, Rennert RC, McArdle A, Paik KJ, Duscher D, Gurtner GC, Lorenz HP, Longaker MT. The Role of Hypoxia-Inducible Factor in Wound Healing. *Adv Wound Care (New Rochelle)*. 2014 May 1;3(5):390-399. doi: 10.1089/wound.2013.0520. PMID: 24804159; PMCID: PMC4005494.
26. Hutami IR, Izawa T, Khurel-Ochir T, Sakamaki T, Iwasa A, Tanaka E. Macrophage Motility in Wound Healing Is Regulated by HIF-1 $\alpha$  via S1P Signaling. *Int J Mol Sci.* 2021 Aug 20;22(16):8992. doi: 10.3390/ijms22168992. PMID: 34445695; PMCID: PMC8396560.
27. Idries A. *Pedoman Ilmu Kedokteran Forensik*. Jakarta: Penerbit Binarupa Aksara; 1997:67-72
28. Indiaminovich I.S., NorbuvayevicA.K., Farhatovich A, *Forensic Medical Aspects of Bicycle Injury*, *American Journal of Medicine and Medical Sciences*, Vol. 11 No. 5, 2021, pp. 415-418. doi: 10.5923/j.ajmms.20211105.12.

29. Infantino V, Santarsiero A, Convertini P, Todisco S, Iacobazzi V. Cancer Cell Metabolism in Hypoxia: Role of HIF-1 as Key Regulator and Therapeutic Target. *International Journal of Molecular Sciences*. 2021; 22(11):5703. <https://doi.org/10.3390/ijms22115703>
30. Kaczorowski DJ, Tsung A, Billiar TR. Innate immune mechanisms in ischemia/ reperfusion. *Front Biosci (Elite Ed)* 2009;1:91e8.
31. Kewley RJ, Whitelaw ML, Chapman-Smith A. The mammalian basic helix-loop-helix/PAS family of transcriptional regulators. *Int J Biochem Cell Biol*. 2004 Feb;36(2):189-204. doi: 10.1016/s1357-2725(03)00211-5. PMID: 14643885.
32. Ishida Y, Kimura A, Nosaka M, et al. . Detection of endothelial progenitor cells in human skin wounds and its application for wound age determination. *Int J Legal Med*. 2015;129:1049–1054.
33. Kaczorowski DJ, Tsung A, Billiar TR. (2009). Innate immune mechanisms in ischemia/ reperfusion. *Front Biosci (Elite Ed)* 2009;1:91e8.
34. Li G, Ko CN, Li D, Yang C, Wang W, Yang GJ, Di Primo C, Wong VKW, Xiang Y, Lin L, Ma DL, Leung CH. A small molecule HIF-1 $\alpha$  stabilizer that accelerates diabetic wound healing. *Nat Commun*. 2021 Jun 7;12(1):3363. doi: 10.1038/s41467-021-23448-7. PMID: 34099651; PMCID: PMC8184911.
35. Li N, Du Q, Bai R, Sun J. Vitality and wound-age estimation in forensic pathology: review and future prospects. *Forensic Sci Res*. 2018;29;5(1):15-24. doi: 10.1080/20961790.2018.1445441.
36. Mao S, Fu F, Dong X. Supplementary pathway for vitality of wounds and wound age estimation in bruises using the electric impedance spectroscopy technique. *J Forensic Sci*. 2011; 56:925–929
37. Martinotti S, Ranzato E. Propolis: a new frontier for wound healing? *Burns Trauma*. 2015 Jul 22;3:9. doi: 10.1186/s41038-015-0010-z. PMID: 27574655; PMCID: PMC4964312.
38. Mitroshina EV, Savyuk MO, Ponimaskin E, Vedunova MV. Hypoxia-Inducible Factor (HIF) in Ischemic Stroke and Neurodegenerative Disease. *Front. Cell Dev. Biol*. 2021; 9:703084. doi:10.3389/fcell.2021.703084
39. Oehmichen M. Vitality and time course of wounds. *Forensic Sci Int*. 2004;144:221–231.

40. Ozgok Kangal MK, Regan JP. Wound Healing. 2023 May 1. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2024 Jan–. PMID: 30571027.
41. Pan Y, Ke H, Yan Z, Geng Y, Asner N, Palani S, Zheng G. The western-type diet induces anti-HMGB1 autoimmunity in Apoe<sup>-/-</sup> mice. *HHS Public Access*, 2017;1(1), 31–38. <https://doi.org/10.1016/j.atherosclerosis.2016.05.027>.
42. Park JW, Chun YS, Kim MS. Hypoxia-inducible factor 1-related diseases and prospective therapeutic tools. *J Pharmacol Sci*. 2004 Mar;94(3):221-32. doi: 10.1254/jphs.94.221. PMID: 15037806
43. Reed K R, Song F, Young MA, Hassan N, Daniel J, Gemici NB, Jenkins JR. Secreted HMGB1 from Wnt activated intestinal cells is required to maintain a crypt progenitor phenotype. 2016; 77(32), 51665– 51673. [www.impactjournals.com/oncotarget/](http://www.impactjournals.com/oncotarget/)
44. Rosamarlina R, Hatta M, Djaharuddin I, Patellongi I, Susanto A. D, Islam A. A, Massi M. N, Bukhari A, Santoso A, Tabri N. A, Murtiani F, Junita A. R, Saleh A. S, Dwiyantri R, Pakadang S. R. The Changes of HIF-1 $\alpha$  and ICAM-1 Expression after Miana (*Coleus Scutellarioides* [L]) Treatment in Balb/C Mice with *Mycobacterium Tuberculosis* Infection. *Biomed Pharmacol J*. 2022;15(1), 73-81. DOI: <https://dx.doi.org/10.13005/bpj/2344>
45. Rusyati LMM, Hatta M, Widiara IGR, Adiguna MS, Wardana M, Dwiyantri R, Noviyanti RA, Sabir M, Yasir Y, Paramit, S, Junita AR, Primaguna MR. Higher *treg* FOXP3 and TGF- $\beta$  mRNA expression in type 2 reaction ENL (*Erythema nodosum leprosum*) patients in *mycobacterium leprae* infection. *Open Microbiology Journal*. 2020; 14 (1), pp. 304-309. DOI: <https://doi.org.10.2174/1874434602014010304>
46. Ruthenborg RJ, Ban JJ, Wazir A, Takeda N, Kim JW. Regulation of wound healing and fibrosis by hypoxia and hypoxia-inducible factor-1. *Mol Cells*. 2014 Sep;37(9):637-43. doi: 10.14348/molcells.2014.0150. Epub 2014 Jun 24. PMID: 24957212; PMCID: PMC4179131.
47. Samec M, Liskova A, Koklesova L, Mersakova S, Strnadel J, Kajo K, Pec M, Zhai K, Smejkal K, Mirzaei S, Hushmandi K, Ashrafzadeh M, Saso L, Brockmueller A, Shakibaei M, Büsselberg D, Kubatka P. Flavonoids Targeting HIF-1: Implications on Cancer Metabolism. *Cancers*. 2021; 13(1):130. <https://doi.org/10.3390/cancers13010130>

48. Santos SAD, Andrade DR Júnior. HIF-1 $\alpha$  and infectious diseases: a new frontier for the development of new therapies. *Rev Inst Med Trop Sao Paulo*. 2017 Dec 21;59:e92. doi: 10.1590/S1678-9946201759092. PMID: 29267600; PMCID: PMC5738998.
49. Shao G, Gao CY, Lu GW. Alterations of Hypoxia-Inducible Factor-1 Alpha in the Hippocampus of Mice Acutely and Repeatedly Exposed to Hypoxia. *Neurosignals* 2005;14:255-261. doi: 10.1159/000088641
50. Stavrou D. Neovascularisation in wound healing. *J Wound Care* 2008;17:298-300, 2.
51. Sumantri S, Hatta M, Natzir R, Rasyid H, Rengganis I, Massi MN, Islam AA, Lawrence G, Patellongi I, Benyamin AF Metformin improves FOXP3 mRNA expression through suppression of interferon gamma levels in pristane-induced murine models of lupus. (2021) *F1000Research*, 9, art. no. 342. DOI: <https://doi.org/10.12688/f1000research.23471.2>
52. Sun JH, Zhu XY, Dong TN. An “up, no change, or down” system: Time-dependent expression of mRNAs in contused skeletal muscle of rats used for wound age estimation. *Forensic Sci Int*. 2017; 272:104–110.
53. Surachmanto EE, Hatta M, Islam AA, Wahid S. Association between asthma control and Interleukin-17F expression levels in adult patients with atopic asthma. *Saudi Medical Journal*, 2018; 39 (7): 662-667. DOI: <https://doi.org/10.15537/smj.2018.7.22055>
54. Suresh MV, Balijepalli S, Zhang B, Singh VV, Swamy S, Panicker S, Dolgachev VA, Subramanian C, Ramakrishnan SK, Thomas B, Rao TC, Delano MJ, Machado-Aranda D, Shah YM, Raghavendran K. Hypoxia-Inducible Factor (HIF-1 $\alpha$ ) Promotes Inflammation and Injury Following Aspiration-Induced Lung Injury in Mice. *Shock*. 2019 Dec;52(6):612-621. doi: 10.1097/SHK.0000000000001312. PMID: 30601332; PMCID: PMC6591116.
55. Suzuki T, Sakata H, Kato C. Astrocyte activation and wound healing in intact-skull mouse after focal brain injury. *Eur J Neurosci*. 2012;36:3653–3664.
56. Syawal P, Budu B, Hatta M, Massi MN, Ichsan AM, Minhajat R. Comparison between the triamcinolone and bevacizumab subconjunctivals and changes in Interleukin-1 mRNA expression in pterygium. *J Taibah Univ Med Sci*. 2021 Aug 16;17(1):67-71. doi: <https://doi.org/10.1016/j.jtumed.2021.07.009>. PMID: 35140567; PMCID: PMC8801476.

57. Syukri A, Budu, Hatta M, Amir M, Rohman MS, Mappangara I, Kaelan C, Wahyuni S, Bukhari A, Junita AR, Primaguna MR, Dwiyantri R, Febrianti A. Doxorubicin induced immune abnormalities and inflammatory responses via HMGB1, HIF1- $\alpha$  and VEGF pathway in progressive of cardiovascular damage. *Ann Med Surg (Lond)*. 2022 Mar 21;76:103501. doi: <https://doi.org/10.1016/j.amsu.2022.103501>. PMID: 35340325; PMCID: PMC8943401.
58. Takamiya M, Fujita S, Saigusa K, et al. . Simultaneous detections of 27 cytokines during cerebral wound healing by multiplexed bead-based immunoassay for wound age estimation. *J Neurotrauma*. 2007;24:1833–1844.
59. Tang D, Kang R, Zeh 3rd HJ, Lotze MT. (2011). High-mobility group box 1, oxidative stress, and disease. *Antioxid Redox Signal*. 14:1315e35
60. Taufik S A, Dirja BT, Utomo DN, Usman MA, Sakti M, Saleh MR, Hatta M, Budu. Double membrane platelet-rich fibrin (PRF) - Synovium succeeds in regenerating cartilage defect at the knee: An experimental study on rabbit. *Heliyon*. 2023 Jan 22;9(2):e13139. doi: <https://doi.org/10.1016/j.heliyon.2023.e13139>. PMID: 36747521; PMCID: PMC9898638.
61. Undang-Undang Republik Indonesia Nomor 36 Tahun 2009 Tentang Kesehatan [internet].C2009 [dilihat 15 Agustus 2020]. Tersedia dari : [http://www.depdagri.go.id/media/documents/2009/10/13/UU\\_No.36](http://www.depdagri.go.id/media/documents/2009/10/13/UU_No.36)
62. VanPatten S, Al-Abed Y. High Mobility Group Box-1 (HMGB1): Current Wisdom and Advancement as a Potential Drug Target: Miniperspective. *J. Med. Chem*. 2018; 61, 5093–5107. <https://doi.org/10.1021/acs.jmedchem.7b01136>
63. Wang LL, Zhao R, Liu CS. A fundamental study on the dynamics of multiple biomarkers in mouse excisional wounds for wound age estimation. *J Forensic Leg Med*. 2016;39:138–146.
64. Weidemann A, Johnson R. Biology of HIF-1 $\alpha$ . *Cell Death Differ*.2008; 15, 621–627 . <https://doi.org/10.1038/cdd.2008.12>
65. Wilkinson HN, Hardman MJ. Wound healing: cellular mechanisms and pathological outcomes. *Open Biol*. 2020 Sep;10(9):200223. doi: 10.1098/rsob.200223. Epub 2020 Sep 30. PMID: 32993416; PMCID: PMC7536089.

66. Wu AH, He L, Long W, Zhou Q, Zhu S, Wang P, Fan S, Wang H. Novel Mechanisms of Herbal Therapies for Inhibiting HMGB1 Secretion or Action. Evidence-based complementary and alternative medicine : eCAM, 2015, 456305. <https://doi.org/10.1155/2015/456305>
67. Xu J, Liu X, Zhao F. HIF1 $\alpha$  overexpression enhances diabetic wound closure in high glucose and low oxygen conditions by promoting adipose-derived stem cell paracrine function and survival. Stem Cell Res Ther. 2020; 11, 148. <https://doi.org/10.1186/s13287-020-01654-2>
68. Yagi Y, Murase T, Kagawa S. Immunohistochemical detection of CD14 and combined assessment with CD32B and CD68 for wound age estimation. Forensic Sci Int. 2016;262: 113–120.
69. Yang C, Zhong ZF, Wang SP, Vong CT, Yu B, Wang YT. HIF-1: structure, biology and natural modulators. Chin J Nat Med. 2021 Jul;19(7):521-527. doi: 10.1016/S1875-5364(21)60051-1. PMID: 34247775.
70. Yanto TA, Hatta M, Bukhari A, Natzir R. Molecular and Immunological Mechanisms of Miana Leaf (*Coleus Scutellariodes* [L] Benth) in Infectious Diseases. Biomed Pharmacol J. 2020; 13(4):1-8. DOI : <https://dx.doi.org/10.13005/bpj/2036>
71. Zetterström CK, Strand ML, Söder O. The high mobility group box chromosomal protein 1 is expressed in the human and rat testis where it may function as an antibacterial factor. Human Reproduction. 2006;21(11); 2801–2809
72. Zhang J, Niu F, Dong H. Characterization of protein alterations in damaged axons in the brainstem following traumatic brain injury using fourier transform infrared microspectroscopy: a preliminary study. J Forensic Sci. 2015; 60:759–763.