

DAFTAR PUSTAKA

- Abebe TB, Gebreyohannes EA, Tefera YG, Bhagavathula AS, Erku DA, Belachew SA, et al. The prognosis of heart failure patients: Does sodium level play a significant role? PLOS ONE. PLoS One. 2018; 13(11): 1-14. doi: 10.1371/journal.pone.0207242
- Arévalo-Lorido JC, Gómez JC, Formiga F, Conde-Martel A, Carrera-Izquierdo M, Muela-Molinero A, et al. High serum osmolarity at admission determines a worse outcome in patients with heart failure Is a new target emerging. *Int J Cardiol.* 2016; 221: 238-42. doi: 10.1016/j.ijcard.2016.07.084
- Chamberlain RS, Sond J, Mahendraraj K, Lau CS, Siracuse BL. Determining 30-day readmission risk for heart failure patients: the Readmission After Heart Failure scale.. *International Journal of General Medicine.* 2018; 9(11): 127-141. doi: 10.2147/IJGM.S150676
- Donzé J, Aujesky D, Williams D, Schnipper JL. Potentially avoidable 30-day hospital readmissions in medical patients: derivation and validation of a prediction model. *JAMA Intern Med.* 2013; 173(8): 632-8. doi: 10.1001/jamainternmed.2013.3023
- Feng J, Zhang Y, Zhang J. Epidemiology and Burden of Heart Failure in Asia. *JACC: Asia.* 2024; 4(4): 249-264. doi: 10.1016/j.jacasi.2024.01.013
- Gheorghiade M, Vaduganathan M, Fonarow GC, Bonow RO. Rehospitalization for heart failure: problems and perspectives. *J Am Coll Cardiol.* 2013; 61(4): 391-403. doi: 10.1016/j.jacc.2012.09.038
- Groenewegen A, Rutten FH, Mosterd A, Hoes AW. Epidemiology of heart failure. *European Journal of Heart Failure,* 2020; 22: 1342-1356. doi: 10.1002/ejhf.1858
- Khan NA, Ma I, Thompson CR, Humphries K, Salem DN, Sarnak MJ, et al. Kidney Function and Mortality among Patients with Left Ventricular Systolic Dysfunction. *J Am Soc Nephrol.* 2006; 17(1): 244-53. doi: 10.1681/ASN.2005030270
- Kilgore M, Patel HK, Kielhorn A, Maya JF, Sharma P. Economic burden of hospitalizations of Medicare beneficiaries with heart failure. *Risk Manag Healthc Policy.* 2017; 10(10): 63-70. doi: 10.2147/RMHP.S130341
- Komukai K, Ogawa T, Yagi H, Date T, Sakamoto H, Kanzaki Y, et al. Decreased renal function as an independent predictor of re-hospitalization for congestive heart failure. *Circ J.* 2008; 72(7): 1152-7. doi: 10.1253/circj.72.1152
- Kwok CS, Abramov D, Purwani P, Ghosh RK, Kittleson M, Ahmad FZ, et al. Cost of inpatient heart failure care and 30-day readmissions in the United States. *International Journal of Cardiology.* 2021; 15(329): 115-122. doi: 10.1016/j.ijcard.2020.12.020

- LeJemtel TH, Serrano C. Vasopressin dysregulation: Hyponatremia, fluid retention and congestive heart failure. *Int J Cardiol.* 2007; 120(1): 1-9. doi: 10.1016/j.ijcard.2006.11.113
- Lu DY, Cheng HM, Cheng YL, Hsu PF, Huang WM, Guo CY, et al. Hyponatremia and worsening sodium levels are associated with Long - Term outcome in patients hospitalized for acute heart failure. *Journal of the American Heart Association.* 2016; 5(3): 1-8. doi: 10.1161/JAHA.115.002668
- O'Connor CM, Miller AB, Blair JE, Konstam MA, Wedge P, Bohit MC, et al. Causes of death and rehospitalization in patients hospitalized with worsening heart failure and reduced left ventricular ejection fraction: Results from efficacy of vasopressin antagonism in heart failure outcome study with tolvaptan (EVEREST) program. *American Heart Journal.* 2010; 159(5): 841-849. doi: 10.1016/j.ahj.2010.02.023
- Radovanovic D, Seifert B, Urban P, Eberli FR, Rickli H, Bertel O, et al. Validity of Charlson Comorbidity Index in patients hospitalised with acute coronary syndrome. Insights from the nationwide AMIS Plus registry 2002–2012. *Heart,* 2014; 100(4): 288-94. doi: 10.1136/heartjnl-2013-304588
- Rangaswami J, Bhalla V, Blair JEA, Chang TI, Costa S, Lentine KL, et al. Cardiorenal Syndrome: Classification, Pathophysiology, Diagnosis, and Treatment Strategies: A Scientific Statement From the American Heart Association. *Circulation,* 2019; 139(16): e840-e878. doi: 10.1161/CIR.0000000000000664
- Robinson R, Hudali T. The HOSPITAL score and LACE index as predictors of 30 day readmission in a retrospective study at a university-affiliated community hospital. *PeerJ.* 2017; 29(5): 1-13. doi: 10.7717/peerj.3137
- Ronco C, Haapio M, House AA, Anavekar N, Bellomo R. Cardiorenal Syndrome. *J Am Coll Cardiol.* 2008; 52(19): 1527-39. doi: 10.1016/j.jacc.2008.07.051
- Şorodoc V, Asaftei A, Puha G, Ceasovschi A, Lionte C, Sîrbu O, et al. Management of Hyponatremia in heart Failure: Practical considerations. *J Pers Med.* 2023; 13(1): 1-17. doi: 10.3390/jpm13010140.
- Stolfo D, Uijl A, Vedin O, Strömberg A, Faxén UL, Rosano GMC, et al. Sex-Based Differences in Heart Failure Across the Ejection Fraction Spectrum: Phenotyping, and Prognostic and Therapeutic Implications. *JACC: Heart Fail.* 2019; 7(6): 505-515. doi: 10.1016/j.jchf.2019.03.011
- Thongprayoon C, Cheungpasitporn W, Petnak T, Ghamrawi R, Thirunavukkarasu S, Chewcharat A, et al. The prognostic importance of serum sodium levels at hospital discharge and one-year mortality among hospitalized patients. *International Journal of Clinical Practice.* 2020; 74(10): 1-19. doi: 10.1111/ijcp.13581.

- Udani SM, Koyner JL. The effects of heart failure on renal function. *Cardiol Clin.* 2010; 28(3): 453-65. doi: 10.1016/j.ccl.2010.04.004.
- van Walraven C, Dhalla IA, Bell C, Etchells E, Stiell IG, Zarnke K, et al. Derivation and validation of an index to predict early death or unplanned readmission after discharge from hospital to the community. *Canadian Medical Association Journal.* 2010; 182(6): 551-557. doi: 10.1503/cmaj.091117
- Vindhya MR, Khayyat S, Shaaban A, Duran BA, Kallail KJ. Decreased Renal Function is Associated with Heart Failure Readmissions. *Cureus.* 2018;10(8): 1-9. doi: 10.7759/cureus.3122.
- Zamora E, Lupón J, Urrutia A, González B, Mas D, Díez C, et al. Prognostic significance of creatinine clearance rate in patients with heart failure and normal serum creatinine. *Revista española de cardiología.* 2007; 60(12): 1315-1318. doi: 10.1157/13113938.