

DAFTAR PUSTAKA

- Arioli, T., Mattner, S. W., & Winberg, P. C. (2015). Applications of seaweed extracts in Australian agriculture: past, present and future. *Journal of Applied Phycology*, 27(5), 2007–2015. <https://doi.org/10.1007/s10811-015-0574-9>
- Astorga-España, M. S., Rodríguez-Galdón, B., Rodríguez-Rodríguez, E. M., & Díaz-Romero, C. (2016). Amino acid content in seaweeds from the Magellan Straits (Chile). *Journal of Food Composition and Analysis*, 53, 77–84. <https://doi.org/10.1016/J.JFCA.2016.09.004>
- Azizah, R. T. N. (2006). *Experiments on Various Cultivation Methods of Latoh (Caulerpa racemosa as an Effort ti Support Production Continuity* (Vol. 11, Issue 2).
- Bachtiar, subchan Y., Tjahjaningsih, W., & Sianita, N. (2012). Effect of algae brown (sargassum sp.) Extract against bacterial growth of escherichia coli. *Journal of Marine and Coastal Science*, 1(1), 53–60.
- Barka, E. A., Vatsa, P., Sanchez, L., Gaveau-Vaillant, N., Jacquard, C., Klenk, H.-P., Clément, C., Ouhdouch, Y., & van Wezel, G. P. (2016). Taxonomy, Physiology, and Natural Products of Actinobacteria. *Microbiology and Molecular Biology Reviews*, 80(1), 1–43. <https://doi.org/10.1128/mmb.00019-15>
- Baweja, P., Kumar, S., Sahoo, D., & Levine, I. (2016). Biology of Seaweeds. *Seaweed in Health and Disease Prevention*, 41–106. <https://doi.org/10.1016/B978-0-12-802772-1.00003-8>
- Boga, C., Del Vecchio, E., Forlani, L., & Franceschetti, M. (2014). Beneficall and Effective Microorganisms for a sustainable Agriculture and Environment. *Environmental Chemistry Letters*, 12(3), 429–434. <https://doi.org/10.1007/s10311-014-0465-3>
- Burtin, P. (2003). Nutritional Value Of Seaweeds. *Electron. J. Environ. Agric. Food Chem*



intra, S. C., & Patel, N. V. (2008). Biochemical composition of algae collected from Sunderban. In *Indian Journal of Marine* . 37, Issue 3).

Chattopadhyay, K., Adhikari, U., Lerouge, P., & Ray, B. (2007). Polysaccharides from *Caulerpa racemosa*: Purification and structural features. *Carbohydrate Polymers*, 68(3), 407–415. <https://doi.org/10.1016/J.CARBPOL.2006.12.010>

Chin, Y. Y., Chang, K. A., Ng, W. M., Eng, Z. P., Chew, L. Y., Neo, Y. P., Yan, S. W., Wong, C. L., Kong, K. W., & Ismail, A. (2023). A comparative evaluation of nutritional composition and antioxidant properties of six Malaysian edible seaweeds. *Food Chemistry Advances*, 3, 100426. <https://doi.org/10.1016/J.FOCHA.2023.100426>

Davis, T. A., Volesky, B., & Mucci, A. (2003). A review of the biochemistry of heavy metal biosorption by brown algae. In *Water Research* (Vol. 37, Issue 18, pp. 4311–4330). Elsevier Ltd. [https://doi.org/10.1016/S0043-1354\(03\)00293-8](https://doi.org/10.1016/S0043-1354(03)00293-8)

Dj Ratoe Oedjoe, M. (2023). Nutritional and Carrageenan Content of Seaweed (*Kappaphycus alvarezii*). *JVIP*, 4(1), 1–1.

Fatmawati, R. , E. (2019). *Seaweed Cultivation Technique (Caulerpa racemosa Using the Spreading Method at the Brackish Water Aquaculture Center in Jepara, Central Java*.

Hani Norziah, M., & Yen Ching, C. (2002). *Nutritional composition of edible seaweed Gracilaria changgi*. www.elsevier.com/locate/foodchem

Hiddink, G. A., Termorshuizen, A. J., & van Bruggen, A. H. C. (2010). *Mixed Cropping and Suppression of Soilborne Diseases* (pp. 119–146). https://doi.org/10.1007/978-90-481-8741-6_5

Holdt, S. L., & Kraan, S. (2011). Bioactive compounds in seaweed: Functional food applications and legislation. In *Journal of Applied Phycology* (Vol. 23, Issue 3, pp. 543–597). <https://doi.org/10.1007/s10811-010-9632-5>

Ikhsan F, I. H. W. R. (2022). Growth Rate of *Kappaphycus alvarezii* Seaweed Green Variety. *Intek Akuakultur*, 6, 82–91.

Irwan Nurdiansyah, S. (2020). Seaweed Growth Rate *Kappaphycus alvarezii* with Spreading Method in the Waters of Natuna Sea Village Sepempang. *Laut Khatulistiwa* (Vol. 3, Issue 1). tan.ac.id/index.php/lk



Ismianti, J., Diniarti, N., & Ghazali, M. (2018). *The Effect of Depth on the Growth of Sea Grapes (Caulerpa racemosa) Using the Longline Method in Tanjung Bele Village, Moyo Hilir District, Sumbawa Regency.*

Khairy, H. M., & El-Shafay, S. M. (2013). Seasonal variations in the biochemical composition of some common seaweed species from the coast of Abu Qir Bay, Alexandria, Egypt. *Oceanologia*, 55(2), 435–452. <https://doi.org/10.5697/OC.55-2.435>

Khan, W., Rayirath, U. P., Subramanian, S., Jithesh, M. N., Rayorath, P., Hodges, D. M., Critchley, A. T., Craigie, J. S., Norrie, J., & Prithiviraj, B. (2009). Seaweed extracts as biostimulants of plant growth and development. In *Journal of Plant Growth Regulation* (Vol. 28, Issue 4, pp. 386–399). <https://doi.org/10.1007/s00344-009-9103-x>

Kitadai, Y., & Kadowaki, S. (2007). Growth, nitrogen and phosphorous uptake rates and O₂ production rate of seaweeds cultured on coastal fish farms. *Graduate School of Agricultural Science, Kogoshima University, Shimoarata, Kagoshima, Japan*, 19, 149–154.

Kumar, M., Gupta, V., Kumari, P., Reddy, C. R. K., & Jha, B. (2011). Assessment of nutrient composition and antioxidant potential of Caulerpaceae seaweeds. *Journal of Food Composition and Analysis*, 24(2), 270–278. <https://doi.org/10.1016/J.JFCA.2010.07.007>

Li, Z., Dong, Y., Zhang, Y., Zheng, M., Jiang, Z., Zhu, Y., Deng, S., Li, Q., & Ni, H. (2023). Lactobacillus-fermentation enhances nutritional value and improves the inhibition on pancreatic lipase and oral pathogens of edible red seaweed *Bangia fuscopurpurea*. *LWT*, 179, 114643. <https://doi.org/10.1016/J.LWT.2023.114643>

Liu, Z., Chen, J., Zhang, J., Wang, K., & Zhang, S. (2024). The evaluation of C, N, P release and contribution to the aquatic environment during *Sargassum* litters biomass decay. *Regional Studies in Marine Science*, 80, 103892. <https://doi.org/10.1016/j.rsma.2024.103892>

Luo, J., Ye, C., Wen, Y., Bao, Y., & Quek, S. Y. (2025). Investigation of solid-state of red seaweed (*Pyropia* spp.) with lactic acid Bacteria: Effects of fermentation time on chemical composition, antioxidant activity, and in vitro digestibility. *Food Chemistry*, 469, 142667. <https://doi.org/10.1016/J.FOODCHEM.2024.142667>

Mahmoud, S., Mustapha, N. M., Muhammad, K., & Ming, C. H. (2023). Antioxidant activities and phenolics content of eight species of seaweeds from the coast of Sabah, Malaysia. *Journal of Food Composition and Analysis*, 118, 105785. <https://doi.org/10.1016/j.jfca.2023.105785>



seaweeds from north Borneo. *Journal of Applied Phycology*, 20(4), 367–373. <https://doi.org/10.1007/s10811-007-9264-6>

Nasmia, Rosyida, E., Masyahoro, A., Putera, F. H. A., & Natsir, S. (2021). The utilization of seaweed-based liquid organic fertilizer to stimulate *Gracilaria verrucosa* growth and quality. *International Journal of Environmental Science and Technology*, 18(6), 1637–1644. <https://doi.org/10.1007/s13762-020-02921-8>

Oliyai, N., Zekri, S., Iraj, A., Oliyai, A., Tanideh, R., Mussin, N. M., Tamadon, A., & Tanideh, N. (2025). Health benefits of algae and marine-derived bioactive metabolites for modulating ulcerative colitis symptoms. *Journal of Functional Foods*, 125, 106690. <https://doi.org/10.1016/J.JFF.2025.106690>

Olle, M., & Williams, I. H. (2013). Effective microorganisms and their influence on vegetable production - A review. In *Journal of Horticultural Science and Biotechnology* (Vol. 88, Issue 4, pp. 380–386). Headley Brothers Ltd. <https://doi.org/10.1080/14620316.2013.11512979>

Ortiz, J., Romero, N., Robert, P., Araya, J., Lopez-Hernández, J., Bozzo, C., Navarrete, E., Osorio, A., & Rios, A. (2006). Dietary fiber, amino acid, fatty acid and tocopherol contents of the edible seaweeds *Ulva lactuca* and *Durvillaea antarctica*. *Food Chemistry*, 99(1), 98–104. <https://doi.org/10.1016/J.FOODCHEM.2005.07.027>

Pal, A., Kamthania, M. C., & Kumar, A. (2014). Bioactive Compounds and Properties of Seaweeds—A Review. *OALib*, 01(04), 1–17. <https://doi.org/10.4236/oalib.1100752>

Piazzi L, Ceccherelli G, & Cinelli F. (2001). Effects Of The Introduced Green Alga *Caulerpa racemosa* In The Mediterranean. *Marine Ecology Progress Series*, 210: 149-159.

Pramita, S., Khali, M., & Muliani, dan. (2022). Cultivation of seaweed *Caulerpa racemosa* on a laboratory scale using liquid organic fertilizer. In *Acta Aquatica: Aquatic Sciences Journal* (Vol. 9).



wan, M., Singh, P., Das, P., Srivastava, U., & Bast, F. (2025a). I factors and heavy metals in edible seaweeds: Challenges, tions, and strategies for safer consumption. *Journal of Food and Analysis*, 140, 107283. [/10.1016/J.JFCA.2025.107283](https://doi.org/10.1016/J.JFCA.2025.107283)

Prashant, N., Sangwan, M., Singh, P., Das, P., Srivastava, U., & Bast, F. (2025b). Anti-nutritional factors and heavy metals in edible seaweeds: Challenges, health implications, and strategies for safer consumption. *Journal of Food Composition and Analysis*, 140, 107283. <https://doi.org/10.1016/J.JFCA.2025.107283>

Purwaningsih, S., Ramadhan, W., Nabila, W. T., Deskawati, E., & Baabud, H. M. (2024). Dataset on the chemical composition and bioactive compound of estuarine seaweed *Gracilaria* from four different cultivation area in Java and Lombok Island, Indonesia. *Data in Brief*, 56, 110825. <https://doi.org/10.1016/J.DIB.2024.110825>

Ratana-Arporn, P., & Chirapart, A. (2006). Nutritional Evaluation of Tropical Green Seaweeds *Caulerpa lentillifera* and *Ulva reticulata*. In *Nat. Sci.* (Vol. 40).

Rodrigues, D., Freitas, A. C., Pereira, L., Rocha-Santos, T. A. P., Vasconcelos, M. W., Roriz, M., Rodríguez-Alcalá, L. M., Gomes, A. M. P., & Duarte, A. C. (2015). Chemical composition of red, brown and green macroalgae from Buarcos bay in Central West Coast of Portugal. *Food Chemistry*, 183, 197–207. <https://doi.org/10.1016/J.FOODCHEM.2015.03.057>

Sa'adah, N., & Widyaningsih, S. (2018). The Effect of CO₂ on Water pH on the Growth of *Caulerpa racemosa* var. *uvifera*. *Journal of Tropical Marine*, 21(1), 17. <https://doi.org/10.14710/jkt.v21i1.2460>

Sahabuddin, Kaheriyah, A., & Chadijah, A. (2017). *The Effect of Increasing Carbon Dioxide (CO₂) Concentration on Population Growth and Performance of Adopted Phytoplankton (Emiliana Huxleyi sp.) on a Laboratory Scale* (Vol. 6, Issue 1).

Sánchez-Machado, D. I., López-Cervantes, J., López-Hernández, J., & Paseiro-Losada, P. (2004). Fatty acids, total lipid, protein and ash contents of processed edible seaweeds. *Food Chemistry*, 85(3), 439–444. <https://doi.org/10.1016/J.FOODCHEM.2003.08.001>

Sitorus, E. R., Santosa, G. W., & Pramesti, R. (2020). Effect of Low Light Intensity *racemosa* (Forsskål) 1873 (Ulvophyceae:Caulerpaeae). *of Marine Research*, 9(1), 13–17. [/10.14710/jmr.v9i1.25376](https://doi.org/10.14710/jmr.v9i1.25376)

nkatesalu, V., Anantharaj, M., & Chandrasekaran, M. (2006). Seaweed extracts on the growth and biochemical constituents of



Vigna sinensis. *Bioresource Technology*, 97(14), 1745–1751.
<https://doi.org/10.1016/J.BIORTECH.2005.06.016>

Sobuj, M. K. A., Rahman, S., & Ali, M. Z. (2024). A review on commercially important seaweed resources from the Bangladesh coast. In *Food Chemistry Advances* (Vol. 4). Elsevier Ltd.
<https://doi.org/10.1016/j.focha.2024.100655>

Suthar, P., Gajaria, T. K., & Reddy, C. R. K. (2019). Production of quality seaweed biomass through nutrient optimization for the sustainable land-based cultivation. *Algal Research*, 42, 101583.
<https://doi.org/10.1016/J.ALGAL.2019.101583>

Tapotubun, A. M. (2018). Chemical Composition of Seaweed (*Caulerpa lentillifera*) from Kei Waters, Maluku with Different Drying Methods. *Jurnal Pengolahan Hasil Perikanan Indonesia*, 21(1), 13.
<https://doi.org/10.17844/jphpi.v21i1.21257>

Tapotubun, A. M., Matrutty, T. E. A. A., Riry, J., Tapotubun, E. J., Fransina, E. G., Mailoa, M. N., Riry, W. A., Setha, B., & Rieuwpassa, F. (2020). Seaweed *Caulerpa* sp position as functional food. *IOP Conference Series: Earth and Environmental Science*, 517(1). <https://doi.org/10.1088/1755-1315/517/1/012021>

Taslim, N. A., Hardinsyah, H., Radu, S., Mayulu, N., Tsopmo, A., Kurniawan, R., Tallei, T. E., Herlina, T., Maksum, I. P., & Nurkolis, F. (2024a). Functional food candidate from Indonesian green algae *Caulerpa racemosa* (Försskal) J. Agardh by two extraction methods: Metabolite profile, antioxidant activity, and cytotoxic properties. *Journal of Agriculture and Food Research*, 18, 101513. <https://doi.org/10.1016/J.JAFR.2024.101513>

Taslim, N. A., Hardinsyah, H., Radu, S., Mayulu, N., Tsopmo, A., Kurniawan, R., Tallei, T. E., Herlina, T., Maksum, I. P., & Nurkolis, F. (2024b). Functional food candidate from Indonesian green algae *Caulerpa racemosa* (Försskal) J. Agardh by two extraction methods: Metabolite profile, antioxidant activity, and cytotoxic properties. *Journal of Agriculture and Food Research*, 18, 101513. <https://doi.org/10.1016/J.JAFR.2024.101513>



Manivannan, K., Devi, G. K., Anantharaman, P., & rian, T. (2009). Proximate Composition of Different Group of m Vedalai Coastal Waters (Gulf of Mannar): Southeast Coast le-East Journal of Scientific Research, 4(2), 72–77.

Windarto, S., Prastiwahyudi, A. H., Susilowati, T., Haditomo, A. H. C., & Harwanto, D. (2021). *Effect of Different Substrates on Growth and Protein Content of Caulerpa Racemosa* (Vol. 48, Issue 7).

Yusuf Bachtiar, S., Tjahjaningsih, W., & Sianita, N. (2012). EFFECT OF ALGAE BROWN (*Sargassum* sp.) EXTRACT AGAINST BACTERIAL GROWTH OF *Escherichia coli*. In *Journal of Marine and Coastal Science* (Vol. 1, Issue 1).

Zakharikhina, L. V., Gorbunova, T. L., Ryndin, A. V., Lesnikova, P. S., & Rogozhina, E. V. (2024). Chemical composition and growth responses of brown, green, and red macroalgae to pollution in the Russian black sea. *Marine Pollution Bulletin*, 207, 116831. <https://doi.org/10.1016/J.MARPOLBUL.2024.116831>

