

DAFTAR PUSTAKA

- Alentorn-Geli, E., Alvarez-Diaz, P., Ramon, S., Marin, M., Steinbacher, G., Boffa, J. J., Cuscó, X., Ballester, J., & Cugat, R. (2015). Assessment of neuromuscular risk factors for anterior cruciate ligament injury through tensiomyography in male soccer players. *Knee Surgery, Sports Traumatology, Arthroscopy: Official Journal of the ESSKA*, 23(9), 2508–2513. <https://doi.org/10.1007/s00167-014-3018-1>
- Alsayed, H. N., Alkhateeb, M. A., Aldossary, A. A., Houbani, K. M., Aljamaan, Y. M., & Alrashidi, Y. A. (2022). Risk of anterior cruciate ligament injury in population with elevated body mass index. *Medicinski Glasnik*, 20(1), 83–87. <https://doi.org/10.17392/1517-22>
- Ashwini T, Aditi Jain, & Ashok Adekal Kumar. (2018, July 1). *MRI Correlation of ACL Injuries with Femoral Intercondylar Notch, Posterior Tibial Slopes and Medial Tibial Plateau Depth in The Indian Population*. International Journal of Anatomy, Radiology and Surgery.
- Balcarek, P., Jung, K., Frosch, K.-H., & Stürmer, K. M. (2011). Value of the Tibial Tuberosity–Trochlear Groove Distance in Patellar Instability in the Young Athlete. *The American Journal of Sports Medicine*, 39(8), 1756–1762. <https://doi.org/10.1177/0363546511404883>
- Basukala, B., Joshi, A., & Pradhan, I. (2020). The Effect of the Intercondylar Notch Shape and Notch Width Index on Anterior Cruciate Ligament Injuries. *Journal of Nepal Health Research Council*, 17(4), 532–536. <https://doi.org/10.33314/jnhrc.v17i4.1858>
- Bayer, S., Meredith, S. J., Wilson, K., De Sa, D., Pauyo, T., Byrne, K., McDonough, C. M., & Musahl, V. (2020). Knee Morphological Risk Factors for Anterior Cruciate Ligament Injury: A Systematic Review. *Journal of Bone and Joint Surgery - American Volume*, 102(8), 703–718. <https://doi.org/10.2106/JBJS.19.00535>
- Beynon, B. D., Hall, J. S., Sturnick, D. R., DeSarno, M. J., Gardner-Morse, M., Tourville, T. W., Smith, H. C., Slaughterbeck, J. R., Shultz, S. J., Johnson, R. J., & Vacek, P. M. (2014). Increased Slope of the Lateral Tibial Plateau Subchondral Bone Is Associated With Greater Risk of Noncontact ACL Injury in Females but Not in Males: A Prospective Cohort Study With a Nested, Matched Case-Control Analysis. *The American Journal of Sports Medicine*, 42(5), 1039–1048. <https://doi.org/10.1177/0363546514523721>
- Cantrell, A. J., Imonugo, O., & Varacallo, M. A. (2025). Anatomy, Bony Pelvis and Lower Limb: Leg Bones. In *StatPearls*. StatPearls Publishing. <http://www.ncbi.nlm.nih.gov/books/NBK537024/>
- Chen, K. J., Lee, E. J., Kliethermes, S. A., & Scerpella, T. A. (2023). Association of Tibial Tubercle–Trochlear Groove Distance With Risk of ACL Graft Failure. *Orthopaedic Journal of Sports Medicine*, 11(6), 23259671231180860. <https://doi.org/10.1177/23259671231180860>

- Choi, N.-H., Lee, D.-M., & Victoroff, B. N. (2023). Bone Morphological Characteristics as Risk Factors for Anterior Cruciate Ligament Injury: Comparison Between Contact and Noncontact Injury. *Orthopaedic Journal of Sports Medicine*, 11(6), 23259671231179757. <https://doi.org/10.1177/23259671231179757>
- Christensen, J. J., Krych, A. J., Engasser, W. M., Vanhees, M. K., Collins, M. S., & Dahm, D. L. (2015). Lateral Tibial Posterior Slope Is Increased in Patients With Early Graft Failure After Anterior Cruciate Ligament Reconstruction. *The American Journal of Sports Medicine*, 43(10), 2510–2514. <https://doi.org/10.1177/0363546515597664>
- Dare, D. M., Fabricant, P. D., McCarthy, M. M., Rebolledo, B. J., Green, D. W., Cordasco, F. A., & Jones, K. J. (2015). Increased Lateral Tibial Slope Is a Risk Factor for Pediatric Anterior Cruciate Ligament Injury: An MRI-Based Case-Control Study of 152 Patients. *The American Journal of Sports Medicine*, 43(7), 1632–1639. <https://doi.org/10.1177/0363546515579182>
- Dean, R. S., DePhillipo, N. N., & LaPrade, R. F. (2022). Posterior Tibial Slope in Patients With Torn ACL Reconstruction Grafts Compared With Primary Tear or Native ACL: A Systematic Review and Meta-analysis. *Orthopaedic Journal of Sports Medicine*, 10(4), 23259671221079380. <https://doi.org/10.1177/23259671221079380>
- Dejour, H., Walch, G., Nove-Josserand, L., & Guier, Ch. (1994). Factors of patellar instability: An anatomic radiographic study. *Knee Surgery, Sports Traumatology, Arthroscopy*, 2(1), 19–26. <https://doi.org/10.1007/BF01552649>
- Etzel, C. M., Meghani, O., Owens, B. D., Kocher, M. S., & Field, A. E. (2024). Predictors of Anterior Cruciate Ligament Tears in Adolescents and Young Adults. *Orthopaedic Journal of Sports Medicine*, 12(9), 23259671241272699. <https://doi.org/10.1177/23259671241272699>
- Evans, J., Mabrouk, A., & Nielson, J. I. (2025). Anterior Cruciate Ligament Knee Injury. In *StatPearls*. StatPearls Publishing. <http://www.ncbi.nlm.nih.gov/books/NBK499848/>
- Fahim, S. M., Dhawan, T., Jagadeesh, N., & Ashwathnarayan, Y. P. (2021). The relationship of anterior cruciate ligament injuries with MRI based calculation of femoral notch width, notch width index, notch shape—A randomized control study. *Journal of Clinical Orthopaedics and Trauma*, 17, 5–10. <https://doi.org/10.1016/j.jcot.2021.01.006>
- Fang, Z., & Liu, W. (2024). Obesity-associated outcomes after ACL reconstruction: A propensity-score-matched analysis of the US Nationwide Inpatient Sample 2005–2018. *Journal of Orthopaedics and Traumatology*, 25(1), 36. <https://doi.org/10.1186/s10195-024-00779-x>
- Fares, A., Horteur, C., Abou Al Ezz, M., Hardy, A., Rubens-Duval, B., Karam, K., Gaulin, B., & Pailhe, R. (2023). Posterior tibial slope (PTS) ≥ 10 degrees is a risk factor for further anterior cruciate ligament (ACL) injury; BMI is not.

- European Journal of Orthopaedic Surgery & Traumatology*, 33(5), 2091–2099. <https://doi.org/10.1007/s00590-022-03406-9>
- Faude, O., Junge, A., Kindermann, W., & Dvorak, J. (2006). Risk Factors for Injuries in Elite Female Soccer Players. *British Journal of Sports Medicine*, 40(9), 785–790. <https://doi.org/10.1136/bjsm.2006.027540>
- Feucht, M. J., Mauro, C. S., Brucker, P. U., Imhoff, A. B., & Hinterwimmer, S. (2013). The role of the tibial slope in sustaining and treating anterior cruciate ligament injuries. *Knee Surgery, Sports Traumatology, Arthroscopy*, 21(1), 134–145. <https://doi.org/10.1007/s00167-012-1941-6>
- Gemayel, A. C., & Varacallo, M. A. (2025). Total Knee Replacement Techniques. In *StatPearls*. StatPearls Publishing. <http://www.ncbi.nlm.nih.gov/books/NBK538208/>
- Görmeli, C. A., Görmeli, G., öztürk, B. Y., Yildirim, O., Gözükar, H., & özdemir, Z. (2015). The effect of the intercondylar notch width index on anterior cruciate ligament injuries: A study on groups with unilateral and bilateral ACL injury. *Acta Orthopædica Belgica*, 81.
- Gupton, M., Imonogo, O., Black, A. C., Launico, M. V., & Terreberry, R. R. (2025). Anatomy, Bony Pelvis and Lower Limb, Knee. In *StatPearls*. StatPearls Publishing. <http://www.ncbi.nlm.nih.gov/books/NBK500017/>
- Hasegawa, A., Otsuki, S., Pauli, C., Miyaki, S., Patil, S., Steklov, N., Kinoshita, M., Koziol, J., D’Lima, D. D., & Lotz, M. K. (2012). Anterior cruciate ligament changes in the human knee joint in aging and osteoarthritis. *Arthritis and Rheumatism*, 64(3), 696–704. <https://doi.org/10.1002/art.33417>
- Hashemi, J., Breighner, R., Chandrashekar, N., Hardy, D. M., Chaudhari, A. M., Shultz, S. J., Slauterbeck, J. R., & Beynnon, B. D. (2011). Hip extension, knee flexion paradox: A new mechanism for non-contact ACL injury. *Journal of Biomechanics*, 44(4), 577–585. <https://doi.org/10.1016/j.jbiomech.2010.11.013>
- Hashemi, J., Chandrashekar, N., Gill, B., Beynnon, B. D., Slauterbeck, J. R., Schutt, R. C., Mansouri, H., & Dabezies, E. (2008). The geometry of the tibial plateau and its influence on the biomechanics of the tibiofemoral joint. *The Journal of Bone and Joint Surgery. American Volume*, 90(12), 2724–2734. <https://doi.org/10.2106/JBJS.G.01358>
- Hashemi, J., Chandrashekar, N., Mansouri, H., Gill, B., Slauterbeck, J. R., Schutt, R. C., Dabezies, E., & Beynnon, B. D. (2010). Shallow Medial Tibial Plateau and Steep Medial and Lateral Tibial Slopes: New Risk Factors for Anterior Cruciate Ligament Injuries. *The American Journal of Sports Medicine*, 38(1), 54–62. <https://doi.org/10.1177/0363546509349055>
- Iriuchishima, T., Ryu, K., & Fu, F. H. (2019). Evaluation of age-related differences in anterior cruciate ligament size. *Knee Surgery, Sports Traumatology, Arthroscopy*, 27(1), 223–229. <https://doi.org/10.1007/s00167-018-5336-1>
- Jagadeesh, N., Kapadi, S., Deva, V., Kariya, A., Jagadeesh, N., Kapadi, S., Deva, V., & Kariya, A. (2021). Risk Factors of ACL Injury. In *Arthroscopy*. IntechOpen. <https://doi.org/10.5772/intechopen.99952>

- Kacprzak, B., Stańczak, M., Surmacz, J., & Hagner-Derengowska, M. (2024). Biophysics of ACL Injuries. *Orthopedic Reviews*, *16*, 126041. <https://doi.org/10.52965/001c.126041>
- Lemeshow, S., Hosmer, D. W., Klar, J., & Lwanga, S. K. (1990). *Adequacy of Sample Size in Health Studies*. Wiley.
- Li, H., Zeng, C., Wang, Y., Wei, J., Yang, T., Cui, Y., Xie, D., Liu, H., & Lei, G. (2018). Association Between Magnetic Resonance Imaging–Measured Intercondylar Notch Dimensions and Anterior Cruciate Ligament Injury: A Meta-analysis. *Arthroscopy: The Journal of Arthroscopic & Related Surgery*, *34*(3), 889–900. <https://doi.org/10.1016/j.arthro.2017.08.299>
- Lippert, L. S. (2011). *Clinical Kinesiology and Anatomy* (5th ed.). F. A. Davis Company.
- Liu, Z., Jiang, J., Yi, Q., Teng, Y., Liu, X., He, J., Zhang, K., Wang, L., Teng, F., Geng, B., Xia, Y., & Wu, M. (2022). An increased posterior tibial slope is associated with a higher risk of graft failure following ACL reconstruction: A systematic review. *Knee Surgery, Sports Traumatology, Arthroscopy*, *30*(7), 2377–2387. <https://doi.org/10.1007/s00167-022-06888-6>
- Loupatty, F., Sukmaningtyas, H., Novriansyah, R., Satoto, B., Priharsanti, C. H. N., & Baskoro, N. (2023). Correlation of tibial tuberosity–trochlear groove distance, anterolateral ligament injury, and the type of trochlear dysplasia with the grade of anterior cruciate ligaments tear. *Egyptian Journal of Radiology and Nuclear Medicine*, *54*(1), 206. <https://doi.org/10.1186/s43055-023-01158-y>
- Maheshwari, S. G., Kuber, R., Lamghare, P., Thomas, J., Arkar, R., Avhad, M., Tharmalingam, T., Bhamare, D. S., Katuri, S., Nagraj, H., Redla, S., Elsayad, M., & Vrizedou, S. (2023). Anterior cruciate ligament (ACL) injury: Correlation with MRI morphometry. *Egyptian Journal of Radiology and Nuclear Medicine*, *54*(1), 130. <https://doi.org/10.1186/s43055-023-01073-2>
- Marouane, H., Shirazi-Adl, A., & Hashemi, J. (2015). Quantification of the role of tibial posterior slope in knee joint mechanics and ACL force in simulated gait. *Journal of Biomechanics*, *48*(10), 1899–1905. <https://doi.org/10.1016/j.jbiomech.2015.04.017>
- McCarthy, M. M., & Hannafin, J. A. (2014). The mature athlete: Aging tendon and ligament. *Sports Health*, *6*(1), 41–48. <https://doi.org/10.1177/1941738113485691>
- Montalvo, A. M., Schneider, D. K., Webster, K. E., Yut, L., Galloway, M. T., Heidt, R. S., Kaeding, C. C., Kremcheck, T. E., Magnussen, R. A., Parikh, S. N., Stanfield, D. T., Wall, E. J., & Myer, G. D. (2019). Anterior Cruciate Ligament Injury Risk in Sport: A Systematic Review and Meta-Analysis of Injury Incidence by Sex and Sport Classification. *Journal of Athletic Training*, *54*(5), 472–482. <https://doi.org/10.4085/1062-6050-407-16>
- Moore, K. L., Dalley, A. F., & Agur, A. M. R. (2013). *Clinically Oriented Anatomy*. Lippincott Williams & Wilkins.

- Murray, M. M., Magarian, E. M., Harrison, S. L., Mastrangelo, A. N., Zurakowski, D., & Fleming, B. C. (2010). The effect of skeletal maturity on functional healing of the anterior cruciate ligament. *The Journal of Bone and Joint Surgery. American Volume*, 92(11), 2039–2049. <https://doi.org/10.2106/JBJS.I.01368>
- Napier, R. J., Garcia, E., Devitt, B. M., Feller, J. A., & Webster, K. E. (2019). Increased Radiographic Posterior Tibial Slope Is Associated with Subsequent Injury Following Revision Anterior Cruciate Ligament Reconstruction. *Orthopaedic Journal of Sports Medicine*, 7(11), 2325967119879373. <https://doi.org/10.1177/2325967119879373>
- Neumann, D. A. (2017). *Kinesiology of The Musculoskeletal System, Foundations for Rehabilitation*. Mosby Elsevier. <http://evolve.elsevier.com/Neumann/>
- Olsen, O. E., Myklebust, G., Engebretsen, L., Holme, I., & Bahr, R. (2003). Relationship between floor type and risk of ACL injury in team handball. *Scandinavian Journal of Medicine & Science in Sports*, 13(5), 299–304. <https://doi.org/10.1034/j.1600-0838.2003.00329.x>
- Orchard, J., Seward, H., McGivern, J., & Hood, S. (2001). Intrinsic and extrinsic risk factors for anterior cruciate ligament injury in Australian footballers. *The American Journal of Sports Medicine*, 29(2), 196–200. <https://doi.org/10.1177/03635465010290021301>
- Piotrowski, S. (2014). *The Motion of the Anterior Cruciate Ligament During Internal and External Rotation of the Knee: A Cadaveric Study*. University of Manitoba.
- Pitman, J., & Ghasemi, A. (2025). *Pitman—2025—Study Finds Gender Gap in Knee Injuries*. Radiological Society of North America. <https://www.rsna.org/media/press/2025/2618>
- Raj, M. A., Mabrouk, A., & Varacallo, M. A. (2025). Posterior Cruciate Ligament Knee Injuries. In *StatPearls*. StatPearls Publishing. <http://www.ncbi.nlm.nih.gov/books/NBK430726/>
- Schoettle, P. B., Zanetti, M., Seifert, B., Pfirrmann, C. W. A., Fucentese, S. F., & Romero, J. (2006). The tibial tuberosity–trochlear groove distance; a comparative study between CT and MRI scanning. *The Knee*, 13(1), 26–31. <https://doi.org/10.1016/j.knee.2005.06.003>
- Seitlinger, G., Scheurecker, G., Högler, R., Labey, L., Innocenti, B., & Hofmann, S. (2014). The position of the tibia tubercle in 0°–90° flexion: Comparing patients with patella dislocation to healthy volunteers. *Knee Surgery, Sports Traumatology, Arthroscopy*, 22(10), 2396–2400. <https://doi.org/10.1007/s00167-014-3173-4>
- Shelbourne, K. D., Davis, T. J., & Klootwyk, T. E. (1998). The Relationship Between Intercondylar Notch Width of the Femur and the Incidence of Anterior Cruciate Ligament Tears. *The American Journal of Sports Medicine*, 26(3), 402–408. <https://doi.org/10.1177/03635465980260031001>
- Smith, H. C., Vacek, P., Johnson, R. J., Slauterbeck, J. R., Hashemi, J., Shultz, S., & Beynon, B. D. (2012a). Risk factors for anterior cruciate ligament injury:

- A review of the literature—Part 1: Neuromuscular and anatomic risk. *Sports Health*, 4(1), 69–78. <https://doi.org/10.1177/1941738111428281>
- Smith, H. C., Vacek, P., Johnson, R. J., Slauterbeck, J. R., Hashemi, J., Shultz, S., & Beynon, B. D. (2012b). Risk Factors for Anterior Cruciate Ligament Injury: A Review of the Literature—Part 2: Hormonal, Genetic, Cognitive Function, Previous Injury, and Extrinsic Risk Factors. *Sports Health: A Multidisciplinary Approach*, 4(2), 155–161. <https://doi.org/10.1177/1941738111428282>
- Song, E. K., Seon, J. K., Kim, M. C., Seol, Y.-J., & Lee, S. H. (2016). Radiologic Measurement of Tibial Tuberosity-Trochlear Groove (TT-TG) Distance by Lower Extremity Rotational Profile Computed Tomography in Koreans. *Clinics in Orthopedic Surgery*, 8(1), 45–48. <https://doi.org/10.4055/cios.2016.8.1.45>
- Sonnery-Cottet, B., & Colombet, P. (2016). Partial tears of the anterior cruciate ligament. *Orthopaedics & Traumatology: Surgery & Research*, 102(1), S59–S67. <https://doi.org/10.1016/j.otsr.2015.06.032>
- Souryal, T. O., & Freeman, T. R. (1993). Intercondylar notch size and anterior cruciate ligament injuries in athletes: A prospective study. *The American Journal of Sports Medicine*, 21(4), 535–539. <https://doi.org/10.1177/036354659302100410>
- Stijak, L., Herzog, R. F., & Schai, P. (2008). Is there an influence of the tibial slope of the lateral condyle on the ACL lesion? A case-control study. *Knee Surgery, Sports Traumatology, Arthroscopy: Official Journal of the ESSKA*, 16(2), 112–117. <https://doi.org/10.1007/s00167-007-0438-1>
- Tensho, K., Kumaki, D., Yoshida, K., Shimodaira, H., Horiuchi, H., & Takahashi, J. (2023). Does posterior tibial slope laterality exist? A matched cohort study between ACL-injured and non-injured knees. *Journal of Experimental Orthopaedics*, 10(1), 132. <https://doi.org/10.1186/s40634-023-00702-z>
- Thompson, J. C. (2010). Netter's Concise Orthopaedic Anatomy. In *Saunders Elsevier* (2nd ed.). Saunders Elsevier.
- Vairo, G. L., Moya-Angeler, J., Siorta, M. A., Anderson, A. H., & Sherbondy, P. S. (2019). Tibial Tubercle-Trochlear Groove Distance Is a Reliable and Accurate Indicator of Patellofemoral Instability. *Clinical Orthopaedics & Related Research*, 477(6), 1450–1458. <https://doi.org/10.1097/CORR.0000000000000711>
- Van Eck, C. F., Martins, C. A. Q., Lorenz, S. G. F., Fu, F. H., & Smolinski, P. (2010). Assessment of correlation between knee notch width index and the three-dimensional notch volume. *Knee Surgery, Sports Traumatology, Arthroscopy*, 18(9), 1239–1244. <https://doi.org/10.1007/s00167-010-1131-3>
- Vaswani, R., Meredith, S. J., Lian, J., Li, R., Nickoli, M., Fu, F. H., & Musahl, V. (2020). Intercondylar Notch Size Can Be Predicted on Preoperative Magnetic Resonance Imaging. *Arthroscopy, Sports Medicine, and Rehabilitation*, 2(1), e17–e22. <https://doi.org/10.1016/j.asmr.2019.10.004>

- Volpi, P., Bisciotti, G. N., Chamari, K., Cena, E., Carimati, G., & Bragazzi, N. L. (2018). *Risk factors of anterior cruciate ligament injury in football players: A systematic review of the literature*. *Muscles, Ligaments and Tendons Journal*.
- Webb, J. M., Salmon, L. J., Leclerc, E., Pinczewski, L. A., & Roe, J. P. (2013). Posterior Tibial Slope and Further Anterior Cruciate Ligament Injuries in the Anterior Cruciate Ligament–Reconstructed Patient. *The American Journal of Sports Medicine*, 41(12), 2800–2804. <https://doi.org/10.1177/0363546513503288>
- Wetters, N., Weber, A. E., Wuerz, T. H., Schub, D. L., & Mandelbaum, B. R. (2016). Mechanism of Injury and Risk Factors for Anterior Cruciate Ligament Injury. *Operative Techniques in Sports Medicine*, 24(1), 2–6. <https://doi.org/10.1053/j.otsm.2015.09.001>
- Wilson, R., & Barhorst, A. A. (2018). Intercondylar Notch Impingement of the Anterior Cruciate Ligament: A Cadaveric In Vitro Study Using Robots. *Journal of Healthcare Engineering*, 2018, 1–27. <https://doi.org/10.1155/2018/8698167>
- Windt, J., & Gabbett, T. J. (2017). How do training and competition workloads relate to injury? The workload—injury aetiology model. *British Journal of Sports Medicine*, 51(5), 428–435. <https://doi.org/10.1136/bjsports-2016-096040>
- Woo, S. L., Hollis, J. M., Adams, D. J., Lyon, R. M., & Takai, S. (1991). Tensile properties of the human femur-anterior cruciate ligament-tibia complex. The effects of specimen age and orientation. *The American Journal of Sports Medicine*, 19(3), 217–225. <https://doi.org/10.1177/036354659101900303>
- Yasen, S. K. (2023). Common knee injuries, diagnosis and management. *Surgery (Oxford)*, 41(4), 215–222. <https://doi.org/10.1016/j.mpsur.2023.02.003>
- Yellin, J. L., Parisien, R. L., Talathi, N. S., Farooqi, A. S., Kocher, M. S., & Ganley, T. J. (2021). Narrow Notch Width is a Risk Factor for Anterior Cruciate Ligament Injury in the Pediatric Population: A Multicenter Study. *Arthroscopy, Sports Medicine, and Rehabilitation*, 3(3), e823–e828. <https://doi.org/10.1016/j.asmr.2021.01.024>
- Zeng, C., Gao, S., Wei, J., Yang, T., Cheng, L., Luo, W., Tu, M., Xie, Q., Hu, Z., Liu, P., Li, H., Yang, T., Zhou, B., & Lei, G. (2013). The influence of the intercondylar notch dimensions on injury of the anterior cruciate ligament: A meta-analysis. *Knee Surgery, Sports Traumatology, Arthroscopy*, 21(4), 804–815. <https://doi.org/10.1007/s00167-012-2166-4>