

REFERENSI

- A. Skolarios, H. Jung, A. Neisius, A. Petřík, B. Somani, T. Tailly, G. Gambaro, 2024. Urolithiasis EAU Guidelines on Urolithiasis. Arnhem, The Netherlands.
- Abufaraj, M., Xu, T., Cao, C., Waldhoer, T., Seitz, C., D'andrea, D., Siyam, A., Tarawneh, R., Fajkovic, H., Schernhammer, E., Yang, L., Shariat, S.F., 2021. Prevalence and Trends in Kidney Stone Among Adults in the USA: Analyses of National Health and Nutrition Examination Survey 2007-2013;2018 Data. *Eur Urol Focus* 7, 1468–1475.
<https://doi.org/10.1016/j.euf.2020.08.011>
- Beigi, F.M.A., Hafshejani, Z.K., Aghahoseini, M., Shirani, M., 2019. Impact of body mass index on success, complications and failure of extracorporeal shock wave lithotripsy. *J Renal Inj Prev* 8, 221–224.
<https://doi.org/10.15171/jrip.2019.41>
- Bishop, K., Momah, T., Ricks, J., 2020. Nephrolithiasis. Primary Care: Clinics in Office Practice 47, 661–671.
<https://doi.org/https://doi.org/10.1016/j.pop.2020.08.005>
- Chaussy, C., Eisenberger, F., Forssmann, B., 2007. Extracorporeal shockwave lithotripsy (ESWL®): A chronology. *J Endourol* 21, 1249–1253. <https://doi.org/10.1089/end.2007.9880>
- Chen, P.C., Liu, Y.T., Hsieh, J.H., Wang, C.C., 2017. A practical formula to predict the stone-free rate of patients undergoing extracorporeal shock

wave lithotripsy. Urol Sci 28, 215–218.

<https://doi.org/10.1016/j.urols.2016.05.004>

Crum, L.A., 1988. Cavitation microjets as a contributory mechanism for renal calculi disintegration in ESWL. J Urol 140, 1587–1590.

[https://doi.org/10.1016/s0022-5347\(17\)42132-x](https://doi.org/10.1016/s0022-5347(17)42132-x)

Damanik, Y.M.B., Rahman, E.Y., Istiana, 2021. LITERATURE REVIEW:

ANALISIS EFIKASI ESWL SEBAGAI TATALAKSANA

NEFROLITIASIS. Homeostasis 4, 795–804.

Dhayal, I.R., Gupta, R., Srivastava, A., Rai, P., 2023. A Comparative Study

on Outcomes of Retrograde Intrarenal Surgery (RIRS), Mini

Percutaneous Nephrolithotomy (PCNL), and Extracorporeal Shock

Wave Lithotripsy (ESWL) for Lower Pole Renal Calculi of < 1.5 cm.

Nephrourol Mon 15. <https://doi.org/10.5812/numonthly-128168>

Ganpule, A.P., Vijayakumar, M., Malpani, A., Desai, M.R., 2016.

Percutaneous nephrolithotomy (PCNL) a critical review. International

Journal of Surgery. <https://doi.org/10.1016/j.ijssu.2016.11.028>

Kallidonis, P., Tsaturyan, A., Lattarulo, M., Liatsikos, E., 2020. Minimally

invasive percutaneous nephrolithotomy (Pcnl): Techniques and

outcomes. Turk J Urol 46, 58–63.

<https://doi.org/10.5152/tud.2020.20161>

Khamchun, S., Yoodee, S., Thongboonkerd, V., 2021. Dual modulatory

effects of diosmin on calcium oxalate kidney stone formation processes:

Crystallization, growth, aggregation, crystal-cell adhesion, internalization into renal tubular cells, and invasion through extracellular matrix. *Biomedicine and Pharmacotherapy* 141. <https://doi.org/10.1016/j.biopha.2021.111903>

Köseoğlu, H., Eroğlu, T., Miriyev, T., 2023. Status of SWL Indicated Patients with Urolithiasis during the Era of SARS-Cov-2 Pandemia. *SAS Journal of Medicine* 9, 1174–1177. <https://doi.org/10.36347/sasjm.2023.v09i11.008>

Leslie, S.W., Sajjad, H., Murphy, P.B., 2025. Renal Calculi, Nephrolithiasis. *StatPearls [Internet]* 2025.

Lin, B.B., Lin, M.E., Huang, R.H., Hong, Y.K., Lin, B.L., He, X.J., 2020. Dietary and lifestyle factors for primary prevention of nephrolithiasis: A systematic review and meta-analysis. *BMC Nephrol* 21. <https://doi.org/10.1186/s12882-020-01925-3>

Liu, S., Nie, P., Wang, H., Guo, J., Shang, Q., Xu, W., Feng, W., 2020. Application of Digital Tomosynthesis in the Diagnosis of Urolithiasis: Comparison with MDCT. *J Endourol* 34, 145–150. <https://doi.org/10.1089/end.2019.0327>

Liu, Y., Chen, Y., Liao, B., Luo, D., Wang, K., Li, H., Zeng, G., 2018. Epidemiology of urolithiasis in Asia. *Asian J Urol*. <https://doi.org/10.1016/j.ajur.2018.08.007>

- Manzoor, H., Leslie, S.W., Saikali, S.W., 2024. Extracorporeal Shockwave Lithotripsy. StatPearls.
- Mishra, P., Pandey, C., Singh, U., Keshri, A., Sabaretnam, M., 2019. Selection of appropriate statistical methods for data analysis. *Ann Card Anaesth* 22, 297–301. https://doi.org/10.4103/aca.ACA_248_18
- Oliveira, B., Teixeira, B., Magalhães, M., Vinagre, N., Fraga, A., Cavadas, V., 2024. Extracorporeal shock wave lithotripsy: retrospective study on possible predictors of treatment success and revisiting the role of non-contrast-enhanced computer tomography in kidney and ureteral stone disease. *Urolithiasis* 52. <https://doi.org/10.1007/s00240-024-01570-7>
- Pricop, C., Radavoi, G.D., Puia, D., Vechiu, C., Jinga, V., 2019. Obesity: A delicate issue choosing the eswl treatment for patients with kidney and ureteral stones? *Acta Endocrinol (Copenh)* 15, 133–138. <https://doi.org/10.4183/aeb.2019.133>
- Reynolds, L.F., Krocak, T., Pace, K.T., 2018. Indications and contraindications for shock wave lithotripsy and how to improve outcomes. *Asian J Urol*. <https://doi.org/10.1016/j.ajur.2018.08.006>
- Setthawong, V., Srisubat, A., Potisat, S., Lojanapiwat, B., Pattanittum, P., 2023. Extracorporeal shock wave lithotripsy (ESWL) versus percutaneous nephrolithotomy (PCNL) or retrograde intrarenal surgery (RIRS) for kidney stones. *Cochrane Database of Systematic Reviews* 2023. <https://doi.org/10.1002/14651858.CD007044.pub4>

Shah, R., Kandel, T.D., Adhikari, D., 2023. Retrograde Intrarenal Surgery versus Mini-Percutaneous Nephrolithotomy for Lower Calyceal Stones of ≤ 2 cm. Journal of College of Medical Sciences-Nepal 19, 399–405. <https://doi.org/10.3126/jcmsn.v19i4.56590>

Shinde, S., Al Balushi, Y., Hossny, M., Jose, S., Al Busaidy, S., 2018. Factors affecting the outcome of extracorporeal shockwave lithotripsy in urinary stone treatment. Oman Med J 33, 209–217. <https://doi.org/10.5001/omj.2018.39>

Sorokin, I., Pearle, M.S., 2018. Medical therapy for nephrolithiasis: State of the art. Asian J Urol. <https://doi.org/10.1016/j.ajur.2018.08.005>

Stamatelou, K., Goldfarb, D.S., 2023. Epidemiology of Kidney Stones. Healthcare (Switzerland). <https://doi.org/10.3390/healthcare11030424>

Whitfield, H.N., 2007. Re: Diabetes Mellitus and Hypertension Associated With Shock Wave Lithotripsy of Renal and Proximal Ureteral Stones at 19 Years of Followup. Eur Urol 51, 281–282. <https://doi.org/https://doi.org/10.1016/j.eururo.2006.09.033>

Yu, C., Longfei, L., Long, W., Feng, Z., Jiping, N., Mao, L., Lin, Q., Hequn, C., 2014. A systematic review and meta-analysis of new onset hypertension after extracorporeal shock wave lithotripsy. Int Urol Nephrol 46, 719–725. <https://doi.org/10.1007/s11255-013-0588-7>

Yusuf, F., Elmekresh, A., Kumar, S., Bagheri, F., 2022. Percutaneous
Nephrolithotripsy in Morbidly Obese Patient: A Case Report. Case Rep
Urol 2022, 1–5. <https://doi.org/10.1155/2022/5899896>