

DAFTAR PUSTAKA

1. Kamath S, Hammad Altaq H, Abdo T. Management of Sepsis and Septic Shock: What Have We Learned in the Last Two Decades? *Microorganisms*. 2023;11(9).
2. Kelm DJ, Perrin JT, Cartin-Ceba R, Gajic O, Schenck L, Kennedy CC. Fluid overload in patients with severe sepsis and septic shock treated with early goal-directed therapy is associated with increased acute need for fluid-related medical interventions and hospital death. *Shock*. 2015;43(1):68–73.
3. Mizera J, Bystroń A, Matuszewska K, Pilch J, Giordano U. Knowledge, management, and complications of sepsis and septic shock: A significant therapeutic challenge in the intensive care unit. *J Acute Dis*. 2024;13(2):53–9.
4. Xiao W, Liu W, Zhang J, Huang L, Liu Y, Hu J, et al. Early persistent exposure to high CVP is associated with increased mortality and AKI in septic shock: A retrospective study. *Am J Emerg Med* [Internet]. 2023;74:146–51. Tersedia pada: <https://doi.org/10.1016/j.ajem.2023.09.043>
5. Malbrain MLNG, Wong A, Nasa P, Ghosh S. Rational Use of Intravenous Fluids in Critically Ill Patients. Springer. Swizerland: Springer; 2024. 47–74 hal.
6. Messmer AS, Dill T, Müller M, Pfortmueller CA. Active fluid de-resuscitation in critically ill patients with septic shock: A systematic review and meta-analysis. *Eur J Intern Med* [Internet]. 2023;109(May 2022):89–96. Tersedia pada: <https://doi.org/10.1016/j.ejim.2023.01.009>
7. Linardi V, Handrawan SE, Liana P, Lestari MI. Positive cumulative fluid balance is a risk factor of mortality in critically septic patients. *Crit Care Shock*. 2021;2021-Decem(6):271–80.
8. Malbrain MLNG, Marik PE, Witters I, Cordemans C, Kirkpatrick AW, Roberts DJ, et al. Fluid overload, de-resuscitation, and outcomes in critically ill or injured patients: A systematic review with suggestions for clinical practice. *Anaesthesiol Intensive Ther*. 2014;46(5):361–80.
9. Silversides JA, Fitzgerald E, Manickavasagam US, Lapinsky SE, Nisenbaum R, Hemmings N, et al. Deresuscitation of patients with iatrogenic fluid overload is associated with reduced mortality in critical illness. *Crit Care Med*. 2018;46(10):1600–7.

10. Silversides JA, McAuley DF, Blackwood B, Fan E, Ferguson AJ, Marshall JC. Fluid management and deresuscitation practices: A survey of critical care physicians. *J Intensive Care Soc.* 2020;21(2):111–8.
11. Munroe E, Weinstein J, Bozyk P, Cahill M, Digiovine B, Esteitie R, et al. Fluid Overload and Deresuscitation Practices In Michigan ICUS. *Chest* [Internet]. 2024;166(4):A6379–80. Tersedia pada: <https://doi.org/10.1016/j.chest.2024.07.021>
12. La Via L, Vasile F, Perna F, Zawadka M. Prediction of fluid responsiveness in critical care: Current evidence and future perspective. *Trends Anaesth Crit Care* [Internet]. 2024;54(November 2023):101316. Tersedia pada: <https://doi.org/10.1016/j.tacc.2023.101316>
13. Gawalkar AA, Batta A. Ultrasound based estimate of central venous pressure: Are we any closer? *World J Cardiol.* 2024;16(6):310–3.
14. Kim K, Choi HS, Chung SP, Kwon WY. Septic Shock. *Essentials Shock Manag.* 2018;
15. Slobod D, Yao H, Mardini J, Natkaniec J, Correa JA, Jayaraman D, et al. Bioimpedance-measured volume overload predicts longer duration of mechanical ventilation in intensive care unit patients. *Can J Anesth.* 2019;66(12):1458–63.
16. Jeong H, Park I, Lee JH, Kim D, Baek S, Kim S, et al. Feasibility study using longitudinal bioelectrical impedance analysis to evaluate body water status during fluid resuscitation in a swine sepsis model. *Intensive Care Med Exp* [Internet]. 2022;10(1). Tersedia pada: <https://doi.org/10.1186/s40635-022-00480-5>
17. Guarino M, Perna B, Cesaro AE, Maritati M, Spampinato MD, Contini C, et al. 2023 Update on Sepsis and Septic Shock in Adult Patients: Management in the Emergency Department. *J Clin Med.* 2023;12(9).
18. Ferreira FL, Bota DP, Bross A, Mélot C, Vincent J-L. Serial evaluation of the SOFA score to predict outcome in critically ill patients. *Jama.* 2001;286(14):1754–8.
19. Haydar S, Spanier M, Weems P, Wood S, Strout T. Comparison of QSOFA score and SIRS criteria as screening mechanisms for emergency department sepsis. *Am J Emerg Med.* 2017;35(11):1730–3.
20. Latten GHP, Polak J, Merry AHH, Muris JWM, Ter Maaten JC, Olgers TJ, et al. Frequency of alterations in qSOFA, SIRS, MEWS and NEWS scores during the emergency department stay in infectious patients: a prospective study. *Int J Emerg Med.* 2021;14(1):1–7.

21. Majmundar A, Lakhani S, Shah S, Lakhani JD. A Comparative Study between SIRS, QSOFA, NEWS AND MEWS Score as Predictors for Inhospital Mortality of Patients Getting Admitted To ICU of a Tertiary Care Teaching Hospital. *Asian J Res Infect Dis*. 2023;12(1):8–14.
22. Basodan N, Al Mehmadi AE, Al Mehmadi AE, Aldawood SM, Hawsawi A, Fatini F, et al. Septic Shock: Management and Outcomes. *Cureus*. 2022;14(12).
23. Mehta A, Khalid A, Swaroop M. Sepsis and Septic Shock. *Intech Open, Slov Univ Ljublijana*. 2018;
24. Malbrain MLNG, Van Regenmortel N, Saugel B, De Tavernier B, Van Gaal PJ, Joannes-Boyau O, et al. Principles of fluid management and stewardship in septic shock: it is time to consider the four D's and the four phases of fluid therapy. *Ann Intensive Care* [Internet]. 2018;8(1). Tersedia pada: <https://doi.org/10.1186/s13613-018-0402-x>
25. Malbrain MLNG, Martin G, Ostermann M. Everything you need to know about deresuscitation. *Intensive Care Med* [Internet]. 2022;48(12):1781–6. Tersedia pada: <https://doi.org/10.1007/s00134-022-06761-7>
26. Pfortmueller CA, Dabrowski W, Wise R, van Regenmortel N, Malbrain MLNG. Fluid accumulation syndrome in sepsis and septic shock: pathophysiology, relevance and treatment—a comprehensive review. *Ann Intensive Care* [Internet]. 2024;14(1). Tersedia pada: <https://doi.org/10.1186/s13613-024-01336-9>
27. Silversides JA, Major E, Ferguson AJ, Mann EE, McAuley DF, Marshall JC, et al. Conservative fluid management or deresuscitation for patients with sepsis or acute respiratory distress syndrome following the resuscitation phase of critical illness: a systematic review and meta-analysis. Vol. 43, *Intensive Care Medicine*. 2017. 155–170 hal.
28. Firestone R, Newton B, Duby J, Cabri A, Sebat C. Effect Of Fluid Deresuscitation On Duration Of Mechanical Ventilation In Septic Shock Patients. *Crit Care Med*. 2017;46(1):1.
29. Lantang EY, George Y, Sugiarto A, Diana A. Central venous pressure as end-point fluid removal in drowning patients: A case report. *Bali J Anesthesiol*. 2022;6(4):247–50.
30. De Backer D, Vincent JL. Should we measure the central venous pressure to guide fluid management? Ten answers to 10 questions. *Crit Care*. 2018;22(1):1–6.
31. Lubis B, Amelia P. High Central Venous Pressure Associated with Mortality in ICU. *Maced J Med Sci*. 2022;20(10):1052–5.

32. Song X, Li J, Li S, Tang Z, Hu X, Zhu Y, et al. Exploring the optimal range of central venous pressure in sepsis and septic shock patients: A retrospective study in 208 hospitals. *Am J Med Sci*. 2024;368(4).
33. H. George Y, Cindryani M. Lower central venous pressure target contributes to higher survival among COVID-19 patients. *Neurol Spinale Med Chir*. 2022;5(3):146.
34. Pomero F, Re R, Meschi M, Montemurro D, Spadafora L, Borretta V, et al. Approach to leg Edema. *Ital J Med*. 2017;11(3):267–77.
35. Marik PE, Byrne L, van Haren F. Fluid resuscitation in sepsis: The great 30 mL per kg hoax. *J Thorac Dis*. 2020;2(1):S37–47.
36. Moschopoulos CD, Dimopoulou D, Dimopoulou A, Dimopoulou K, Protopapas K, Zavras N, et al. New Insights into the Fluid Management in Patients with Septic Shock. *Medicina (B Aires)* [Internet]. 2018;365(1):2. Tersedia pada: <https://www.scopus.com/inward/record.uri?eid=2-s2.0-0032126564&doi=10.1016%2FS0306-3623%2897%2900424-2&partnerID=40&md5=e70f13173e905d39509a5672ebc09801>
37. Dargent A, Dumargne H, Labruyère M, Brezillon S, Brassart-Pasco S, Blot M, et al. Role of the interstitium during septic shock: a key to the understanding of fluid dynamics? *J Intensive Care* [Internet]. 2023;11(1):1–13. Tersedia pada: <https://doi.org/10.1186/s40560-023-00694-z>
38. Yanagisawa N, Koshiyama M, Watanabe Y, Sato S, Sakamoto SI. A quantitative method to measure skin thickness in leg edema in pregnant women using B-scan portable ultrasonography: A comparison between obese and non-obese women. *Med Sci Monit*. 2019;25:1–9.
39. Mourik N Van, Metske HA, Hofstra JJ, Binnekade JM, Geerts BF, Schultz MJ, et al. Cumulative fluid balance predicts mortality and increases time on mechanical ventilation in ARDS patients: An observational cohort study. *PLoS One*. 2019;14(10):1–13.
40. Dash A, Jena SK, Chhotray P, Panda S, Mishra SB, Nayak B. Association between cumulative fluid balance and clinical outcome in patients with septic shock: A retrospective observational study. *Int J Acad Med*. 2023;9(3):121–5.
41. Cunha ARL, Lobo SMA. What happens to the fluid balance during and after recovering from septic shock? *Rev Bras Ter Intensiva*. 2015;27(1):10–7.

42. Tigabu BM, Davari M, Kebriaeezadeh A, Mojtahedzadeh M. Fluid volume, fluid balance and patient outcome in severe sepsis and septic shock: A systematic review. *J Crit Care* [Internet]. 2018;48:153–9. Tersedia pada: <https://doi.org/10.1016/j.jcrc.2018.08.018>
43. Chung YJ, Kim EY. Usefulness of bioelectrical impedance analysis and ECW ratio as a guidance for fluid management in critically ill patients after operation. *Sci Rep* [Internet]. 2021;11(1):1–10. Tersedia pada: <https://doi.org/10.1038/s41598-021-91819-7>
44. Malbrain MLNG, Huygh J, Dabrowski W, De Waele JJ, Staelens A, Wauters J. The use of bio-electrical impedance analysis (BIA) to guide fluid management, resuscitation and deresuscitation in critically ill patients: A bench-to-bedside review. *Anaesthesiol Intensive Ther.* 2014;46(5):381–91.
45. Rochweg B, Cheung JH, Ribic CM, Lalji F, Clarke FJ, Gantareddy S, et al. Assessment of Postresuscitation Volume Status by Bioimpedance Analysis in Patients with Sepsis in the Intensive Care Unit: A Pilot Observational Study. *Can Respir J.* 2016;2016.
46. Cleymaet R, D'Hondt M, Scheinok T, Malbrain L, De Laet I, Schoonheydt K, et al. Comparison of Bioelectrical Impedance Analysis (BIA)-Derived Parameters in Healthy Volunteers and Critically Ill Patients. *Life.* 2024;14(1):1–19.
47. Cimmino F, Petrella L, Cavaliere G, Ambrosio K, Trinchese G, Monda V, et al. A Bioelectrical Impedance Analysis in Adult Subjects : The Relationship between Phase Angle and Body Cell Mass. *J Funct Morphol Kinesiol.* 2023;8.
48. Marra M, Sammarco R, Lorenzo A De, Iellamo F, Siervo M, Pietrobelli A, et al. Assessment of Body Composition in Health and Disease Using Bioelectrical Impedance Analysis (BIA) and Dual Energy X-Ray Absorptiometry (DXA): A Critical Overview. *Contrast Media Mol Imaging.* 2019;2019.
49. Jha V, Jairam A, Sharma MC, Sakhuja V, Piccoli A, Parthasarathy S. Body composition analysis with bioelectric impedance in adult Indians with ESRD : Comparison with healthy population. *Kidney Int* [Internet]. 2006;69(9):1649–53. Tersedia pada: <http://dx.doi.org/10.1038/sj.ki.5000293>
50. Nisini N, Corda A, Biretoni F, Miglio A, Teresa M. Bioelectrical Impedance Analysis (BIA) detects body resistance increase in dogs undergoing blood donation. *Vet Res Commun.* 2024;48:3889–97.
51. Moonen HPFX, Van Zanten ARH. Bioelectric impedance analysis for body composition

- measurement and other potential clinical applications in critical illness. *Curr Opin Crit Care*. 2021;27(4):344–53.
52. Silva AM, Campa F, Stagi S, Gobbo LA, Buffa R, Toselli S, et al. The bioelectrical impedance analysis (BIA) international database : aims , scope , and call for data. *Eur J Clin Nutr*. 2023;(7).
 53. Shin J, Park I, Lee JH, Han JS, Kim B, Jang DH, et al. Comparison of body water status and its distribution in patients with non-septic infection, patients with sepsis, and healthy controls. *Clin Exp Emerg Med*. 2021;8(3):173–81.
 54. Khalil SF, Mohktar MS, Ibrahim F. The theory and fundamentals of bioimpedance analysis in clinical status monitoring and diagnosis of diseases. *Sensors (Switzerland)*. 2014;14(6):10895–928.
 55. Jalalzadeh M, Hajiesmaeili M. Bio-electrical impedance analysis in patients with critical illnesses. *Nephrourol Mon*. 2017;9(1):1–5.
 56. Dumitriu AM, Cobilinschi C, Dumitriu B, Sebastian V, Marina I, Ungureanu R, et al. Advancing Nutritional Care Through Bioelectrical Impedance Analysis in Critical Patients. *Nutrients*. 2025;17:380.
 57. Suh SW, Park HJ, Choi YS. Preoperative volume assessment using bioelectrical impedance analysis for minimizing blood loss during hepatic resection. *Hpb [Internet]*. 2022;24(4):568–74. Tersedia pada: <https://doi.org/10.1016/j.hpb.2021.09.009>
 58. Ishiyama D, Toi K, Aoyagi Y, Suzuki K, Takayama T, Yazu H, et al. The extracellular-to-total body water ratio reflects improvement in the activities of daily living in patients who experienced acute stroke. *J Stroke Cerebrovasc Dis [Internet]*. 2024;33(8):107810. Tersedia pada: <https://doi.org/10.1016/j.jstrokecerebrovasdis.2024.107810>
 59. Ruste M, Chabanol C, Fellahi J-L, Jacquet-Lagrèze M. Bioelectrical impedance analysis during deresuscitation: correlation and agreement with cumulative fluid balance in ICU patients. *J Clin Monit Comput*. 2023;37(2):679–87.
 60. Kyosebekirov E, Kazakov D, Nikolova-Kamburova S, Stoilov V, Mitkovski E, Pavlov G, et al. Bioimpedance Analysis for Fluid Status Assessment in Critically Ill Septic Patients. *Folia Med (Plovdiv)*. 2024;66(3):323–31.
 61. Sugiyono. *Metode Penelitian Kuantitatif Kualitatif & RND*. Bandung: Alfabeta; 2016.
 62. Schober P, Boer C, Schwarte LA. Correlation Coefficients: Appropriate Use and

- Interpretation. *Anesth Analg*. 2018;126(5):1763–8.
63. Naorungroj T, Prajantasen U, Sanla-eat T, Viarasilpa T, Tongyoo S. Restrictive fluid management with early de-escalation versus usual care in critically ill patients (reduce trial): a feasibility randomized controlled trial. *Crit Care*. 2025;29:405.
 64. Bissell BD, Wieruszewski PM, Newsome AS, Verkerk BS, Witenko CJ, Ammar MA, et al. Fluid management and dereuscitation practices – An opportunity for collaboration. *J Intensive Care Soc*. 2020;24(3):1–3.
 65. Serra-Prat M, Lorenzo I, Palomera E, Ramírez S, Yébenes JC. Total body water and intracellular water relationships with muscle strength, frailty and functional performance in an elderly population. a cross-sectional study. *J Nutr Heal Aging*. 2019;23(1):96–101.
 66. Piccoli A, Pittoni G, Facco E, Favaro E, Pillon L. Relationship between central venous pressure and bioimpedance vector analysis in critically ill patients. *Crit Care Med*. 2000;28(1):132–7.
 67. Sakaguchi T, Hirata A, Kashiwase K, Higuchi Y, Ohtani T, Sakata Y, et al. Relationship of Central Venous Pressure to Body Fluid Volume Status and Its Prognostic Implication in Patients With Acute Decompensated Heart Failure. *J Card Fail* [Internet]. 2020;26(1):15–23. Tersedia pada: <https://doi.org/10.1016/j.cardfail.2018.06.001>
 68. Karpaviciute J, Skarupskiene I, Balciuviene V, Vaiciunien R, Žiginskiene E, Bumblyte IA. Assessment of Fluid Status by Bioimpedance Analysis and Central Venous Pressure Measurement and Their Association with the Outcomes of Severe Acute Kidney Injury. *Medicina (B Aires)*. 2021;57:518.
 69. Dahlan F, Nurdin H, Palinrungi AS, Salam SH, Muchtar F, Tan CW. The relationship of bioimpedance analysis to central venous pressure, degree of edema, and cumulative fluid balance in septic shock patients in the intensive care unit. *Crit Care Shock*. 2025;28(6).
 70. George YWH, Cindryani M. Lower Central Venous Pressure Target Contributes to Higher Survival among COVID-19 Patients. *Neurol Spinale Med Chir*. 2022;5:146–9.
 71. Shah P, Louis MA. Physiology, Central Venous Pressur. *StatPearls* [Internet] Treasure Isl StatPearls Publ. 2025;
 72. Farcy DA, Jain A, Dalley M, Scalea TM. Pitfalls in Using Central Venous Pressure as a Marker of Fluid Responsiveness. *Emerg Med*. 2016;
 73. Mohammedin AS, AlSaid AH, Almalki AM, Alsaiani AR, Alghamdi FN, Jalalah AA, et al.

- Assessment of Hydration Status and Blood Pressure in a Tertiary Care Hospital at Al-Khobar. *Cureus*. 2022;14(8):10–5.
74. Akimoto T, Tasaki K, Ishihara M. Association of Body Water Balance , Nutritional Risk , and Sarcopenia with Outcome in Patients with Acute Ischemic. *Nutrients*. 2024;16:2165.
 75. Amano K, Liu D, Bruera E, Hui D. Collapse of Fluid Balance and Association with Survival in Patients with Advanced Cancer Admitted to a Palliative Care Unit : Preliminary Findings. *J Palliat Med*. 2020;23(4):459–65.
 76. Ge YZ, Ruan GT, Zhang Q, Dong WJ, Zhang X, Song MM. Extracellular water to total body water ratio predicts survival in cancer patients with sarcopenia : a multi - center cohort study. *Nutr Metab (Lond)* [Internet]. 2022;1–9. Tersedia pada: <https://doi.org/10.1186/s12986-022-00667-3>
 77. Levenbrown Y, Costarino AT. CHAPTER 17 Edema [Internet]. Third Edit. *Nephrology and Fluid/Electrolyte Physiology*. Elsevier Inc.; 2019. 271–290 hal. Tersedia pada: <https://doi.org/10.1016/B978-0-323-53367-6.00017-0>
 78. Nieto ORP, Wong A, Fermín JL, Iván E, López Z, Antonio J, et al. Aiming for zero fluid accumulation : First , do no harm. *Anaesthesiol Intensive Ther*. 2021;53(2).
 79. Cleymaet R, D'Hondt M, Scheinok T, Malbrain L, De Laet I, Schoonheydt K, et al. Comparison of Bioelectrical Impedance Analysis (BIA)-Derived Parameters in Healthy Volunteers and Critically Ill Patients. *Life*. 2024;14(1).
 80. Wang Y, Liu J, Hao H, Lu Q, Zhang L, Wei G. Non-linear association between extracellular water / total body water ratio and all-cause mortality : a population-based cohort study. *Sci Rep*. 2025;15:1–12.
 81. Joskova V, Patkova A, Havel E, Najpaverova S, Uramova D. The Bioimpedance Spectroscopy as Useful Tool for Measuring the Fluid Excess and Fluid Management in Severe Polytrauma Patients. *Physiol Res*. 2019;68:255–64.
 82. Balik M, Sedivy J, Waldauf P, Kolar M, Smejkalova V, Pachl J. Can bioimpedance determine the volume of distribution of antibiotics in sepsis? *Anaesth Intensive Care*. 2005;33(3):345–50.
 83. Ruste M, Chabanol C, Fellahi J-L, Jacquet-Lagrèze M. Bioelectrical impedance analysis during deresuscitation: correlation and agreement with cumulative fluid balance in ICU patients. *J Clin Monit Comput*. 2023;37:679–687.

84. Malbrain MLNG, Langer T, Annane D, Gattinoni L, Elbers P, Hahn RG, et al. Intravenous fluid therapy in the perioperative and critical care setting : Executive summary of the International Fluid Academy (IFA). *Ann Intensive Care* [Internet]. 2020;10. Tersedia pada: <https://doi.org/10.1186/s13613-020-00679-3>