

DAFTAR PUSTAKA

1. Lirk P, Thiry J, Bonnet M for the PROSPECT Working Group, et al. Pain management after laparoscopic hysterectomy: systematic review of literature and PROSPECT recommendations. *Regional Anesthesia & Pain Medicine* 2019;44:425-436
2. Mortazavi M, Parish M, Dorosti A, Anvari H. Comparison of General Anesthesia With Spinal Anesthesia on the Quality of Recovery of Patients With Selective Abdominal Hysterectomy in Patients Visiting the Largest Women's Disease Hospital in Northwestern Iran. *International Journal of Women's Health and Reproduction Sciences*, Vol. 10, No. 1, January 2022, ISSN 2330-4456. doi: 10.15296/ijwhr.2022.06
3. Giannini A, D'Oria O, Bogani G, Di Donato V, Vizza E, Chiantera V, Laganà AS, Muzii L, Salerno MG, Caserta D, Gerli S, Favilli A. Hysterectomy: Let's Step Up the Ladder of Evidence to Look Over the Horizon. *J Clin Med*. 2022 Nov 25;11(23):6940. doi: 10.3390/jcm11236940. PMID: 36498515; PMCID: PMC9737634.
4. Chen B, Littman ED, Milki AA, Westphal LM, Friedman BE, Schmiesing CA. Gynecology/infertility surgery. In: Jaffe RA, Schmiesing CA, Golianu B, editors. *Antesthesiologist's manual of surgical procedures*. 5th ed. Philadelphia: Wolters Kluwer; 2014. p.795-818
5. Lockwood, N. and Misra, L. (2020) Efficacy of Perioperative Intravenous Lidocaine for Multimodal Analgesia. *Open Journal of Anesthesiology*, 10, 299-312. doi: 10.4236/ojanes.2020.109027.
6. Yon JH, Choi GJ, Kang H, Park JM, Yang HS. Intraoperative systemic lidocaine for pre-emptive analgesics in subtotal gastrectomy: a prospective, randomized, double-blind, placebo-controlled study. *Can J Surg*. 2014 Jun;57(3):175-82. doi: 10.1503/cjs.009613. PMID: 24869609; PMCID: PMC4035399
7. Wong R, Luis S, Sessler D, Jacobs J, Mascha E, Lo Menzo E, Swanson E, Aleman R, Frieder J, Aguilar R.P., Azer G. Ketamine and/or lidocaine analgesia for abdominal surgery. *CDC Clinical Trials*. 2018. Available from: https://cdn.clinicaltrials.gov/large-docs/48/NCT04084548/Prot_SAP_000.pdf
8. Prete A, Yan Q, Al-Tarrah K, et al. The cortisol stress response induced by surgery: A systematic review and meta-analysis. *Clin Endocrinol (Oxf)*. 2018; 89: 554–567. <https://doi.org/10.1111/cen.13820>
9. Jendoubi, Ali; Naceur, Imed Ben; Bouzouita, Abderrazak; Trifa, Mehdi1; Ghedira, Salma; Chebil, Mohamed; Houissa, Mohamed. A comparison between intravenous lidocaine and ketamine on acute and chronic pain after open nephrectomy: A prospective, double-blind, randomized, placebo-controlled study. *Saudi Journal of Anaesthesia* 11(2):p 177-184, Apr–Jun 2017. | DOI: 10.4103/1658-354X.203027

10. Lubis AS, Irina RS, Lubis AP, Harahap J. Comparison of Changes in Cortisol Levels in the Blood of Patients Undergoing Craniotomy Using Continuous Infusion Lidocaine and Fentanyl. *Jurnal Neuroanestesi Indonesia*. 2024. doi: <https://doi.org/10.24244/jni.v13i1.565> ISSN (Print): 2088-9674 ISSN (Online): 2460-2302
11. Carugno J, Fatehi M. Abdominal Hysterectomy. [Updated 2023 Jul 18]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2025 Jan-. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK564366/>
12. Osborn K, Simmons S. Anaesthesia for gynaecological surgery. In: Russel R, editor. *Anaesthesia for obstetrics and gynaecology*. London: BMJ Publishing Group; 2000. p.303-41.
13. Della Corte L, Mercurio A, Palumbo M, Viciglione F, Cafasso V, Candice A, Bifulco G, Giampaolino P. Minimally invasive anesthesia for laparoscopic hysterectomy: a case series. *Arch Gynecol Obstet*. 2022 Dec;306(6):2001-2007. doi: 10.1007/s00404-022-06727-6. Epub 2022 Aug 5. PMID: 35931899; PMCID: PMC9362356.
14. Maheswari K, Naguib MA. Local anesthetics. In: Flood P, Rathmell JP, Shafer S, editors. *Stoelting's pharmacology & physiology in anesthetic practice*. 5th ed. Philadelphia: Wolters Kluwer; 2015. p:282-308
15. Chu R, Umukoro N, Greer T, Roberts J, Adekoya P, Odonkor CA, Hagedorn JM, Olatoye D, Urits I, Orhurhu MS, Umukoro P, Viswanath O, Hasoon J, Kaye AD, Orhurhu V. Intravenous Lidocaine Infusion for the Management of Early Postoperative Pain: A Comprehensive Review of Controlled Trials. *Psychopharmacol Bull*. 2020 Oct 15;50(4 Suppl 1):216-259. PMID: 33633427; PMCID: PMC7901134
16. Lee IW, Schraag S. The Use of Intravenous Lidocaine in Perioperative Medicine: Anaesthetic, Analgesic and Immune-Modulatory Aspects. *J Clin Med*. 2022 Jun 20;11(12):3543. doi: 10.3390/jcm11123543. PMID: 35743617; PMCID: PMC9224677.
17. Rosenbaum SB, Gupta V, Patel P, et al. Ketamine. [Updated 2024 Jan 30]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2025 Jan-. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK470357/>
18. Babak Hosseinzadeh Zorofchi, Elahe Jahan, Setare Nassiri, Atosa Najmodin, Elham Saffarieh. Comparing Spinal and General Anesthesia in terms of Postoperative Pain in Patients undergoing Hysterectomy. *Journal of Obstetrics, Gynecology and Cancer Research*. 2018;3(2):73-77
19. Hitt J, Lee R, Elkin P. Pain severity scale: A methodology for classifying postoperative pain severity by surgical procedure. *Surg Open Sci*. 2023 Feb 23;12:29-34. doi: 10.1016/j.sopen.2023.02.003. PMID: 36926590; PMCID: PMC10011477.

20. Grady, Martin V. MD; Mascha, Edward PhD; Sessler, Daniel I. MD; Kurz, Andrea MD. The Effect of Perioperative Intravenous Lidocaine and Ketamine on Recovery After Abdominal Hysterectomy. *Anesthesia & Analgesia* 115(5):p 1078-1084, November 2012. | DOI: 10.1213/ANE.0b013e3182662e01
21. Ciobotaru, O. C., Ciobotaru, O. R., Voicu, D. C., Barna, O., Barna, I., & Voinescu, D. C. (2016). Postoperative pain after total abdominal hysterectomy and bilateral salpingo-oophorectomy depending on the type of anaesthesia administration. *Biotechnology & Biotechnological Equipment*, 30(2), 341–345. <https://doi.org/10.1080/13102818.2015.1135759>
22. Meier IM, Eikemo M, Trøstheim M, et al Factors associated with use of opioid rescue medication after surgery *Regional Anesthesia & Pain Medicine* Published Online First: 21 July 2023. doi: 10.1136/rapm-2023-104412
23. Kim TK, Yoon JR. Comparison of the neuroendocrine and inflammatory responses after laparoscopic and abdominal hysterectomy. *Korean J Anesthesiol.* 2010 Oct;59(4):265-9. doi: 10.4097/kjae.2010.59.4.265. Epub 2010 Oct 21. PMID: 21057617; PMCID: PMC2966708.
24. Hajiesmaeili M R, Behdad S, Abbasi H R, Ayatollahi V, Khadiv Z, et al. Analgesic Effects of Intravenous Ketamine during Spinal Anesthesia in Pregnant Women Undergone Caesarean Section; A Randomized Clinical Trial. *Anesth Pain Med.* 2013;3(2): 230-3. <https://doi.org/10.5812/aapm.7034>.
25. D.M. Rascón-Martínez, O. Carrillo-Torres, R.G. Ramos-Nataren, L. Rendón-Jaramillo. Advantages of ketamine as a perioperative analgesic. *Revista Médica del Hospital General de México.* Volume 81, Issue 4. 2018. Pages 253-261. ISSN 0185-1063. <https://doi.org/10.1016/j.hgmx.2016.10.007>.
26. Fillingim RB, King CD, Ribeiro-Dasilva MC, Rahim-Williams B, Riley JL. Sex, gender, and pain: a review of recent clinical and experimental findings. *J Pain.* 2009;10(5):447–485.
27. Wright JD, Herzog TJ, Tsui J, Ananth CV, Lewin SN, Lu YS, et al. Nationwide trends in the performance of inpatient hysterectomy in the United States. *Obstet Gynecol.* 2013;122(2 Pt 1):233–241.
28. Zhang T, Yue Z, Yu L, Li S, Xie Y, Wei J, et al. S-ketamine promotes postoperative recovery of gastrointestinal function and reduces postoperative pain in gynecological abdominal surgery patients: a randomized controlled trial. *BMC Surg.* 2023;23:74.
29. Zhang J, Niu Z, Wang T, Yu L, Ren X, Zhang S, Zhu Y, Qi D. Effect of low-dose esketamine on postoperative quality of recovery in total laparoscopic hysterectomy: a randomized controlled trial. *Perioper Med (Lond).* 2025 Jul 23;14(1):78.

30. Zhang M, Jin Y, Yuan X, He C, Han M, Tu F, et al. Effect of obesity on pharmacokinetics of lidocaine and its active metabolites in surgical patients. *Clin Pharmacokinet.* 2025;64(8):1179–1190.
31. Kranke P, Jokinen J, Pace NL, Schnabel A, Hollmann MW, Hahnenkamp K, et al. Continuous intravenous perioperative lidocaine infusion for postoperative pain and recovery. *Cochrane Database Syst Rev.* 2015;7:CD009642.
32. Imani F, Bagheri AR, Arvin E, Gatt SP, Sarveazad A. Effects of ketamine and lidocaine infusion on acute pain after elective open abdominal surgery. *Med J Islam Repub Iran.* 2022;36:60.
33. Kehlet H, Dahl JB. Anaesthesia, surgery, and challenges in postoperative recovery. *Lancet.* 2003;362(9399):1921–1928.
34. Gerbershagen HJ, Aduckathil S, van Wijck AJ, Peelen LM, Kalkman CJ, Meissner W. Pain intensity on the first day after surgery. *Anesthesiology.* 2013;118(4):934–944.
35. Schwenk ES, Viscusi ER, Buvanendran A, Hurley RW, Wasan AD, Narouze S, et al. Consensus guidelines on the use of intravenous ketamine infusions for acute pain management. *Reg Anesth Pain Med.* 2018;43(5):456–466.
36. Zhou L, Duan J. The role of N-methyl-D-aspartate receptors in the anesthetic and analgesic effects of ketamine. *CNS Neurosci Ther.* 2024;30(4):e14464.
37. Weibel S, Jelting Y, Pace NL, Helf A, Eberhart LHJ, Hahnenkamp K, et al. Continuous intravenous perioperative lidocaine infusion for postoperative pain and recovery in adults. *Cochrane Database Syst Rev.* 2018;6:CD009642.
38. Brinck EC, Tiippana E, Heesen M, Bell RF, Straube S, Moore RA, et al. Perioperative intravenous ketamine for acute postoperative pain in adults. *Cochrane Database Syst Rev.* 2018;12:CD012033.
39. Wang X, Lin C, Lan L, Liu J. Perioperative intravenous S-ketamine for acute postoperative pain in adults: a systematic review and meta-analysis. *J Clin Anesth.* 2021;68:110071.
40. Zhang T, Yue Z, Yu L, Li S, Xie Y, Wei J, et al. S-ketamine promotes postoperative recovery of gastrointestinal function and reduces postoperative pain in gynecological abdominal surgery patients: a randomized controlled trial. *BMC Surg.* 2023;23:74.
41. Xie Y, Molla YD, Setargew KH, Alemu HT. The role of postoperative ketamine infusion in ERAS protocols following elective open abdominal surgeries: a systematic review and meta-analysis. *BMC Surg.* 2026;26:94.
42. Wang L, Johnston B, Kaushal A, Cheng D, Zhu F, Martin J. Ketamine for postoperative pain: a systematic review and meta-analysis of randomized controlled trials. *J Clin Anesth.* 2021;68:110071.

43. Riddell JM, Trummel JM, Onakpoya IJ. Low-dose ketamine in painful orthopaedic surgery: a systematic review and meta-analysis. *Br J Anaesth*. 2019;123(3):325–334.
44. Bell RF, Dahl JB, Moore RA, Kalso E. Perioperative ketamine for acute postoperative pain. *Cochrane Database Syst Rev*. 2006;(1):CD004603.
45. Sun Y, Li T, Wang N, Yun Y, Gan TJ. Perioperative systemic lidocaine for postoperative analgesia and recovery after abdominal surgery: a meta-analysis of randomized controlled trials. *Dis Colon Rectum*. 2012;55(11):1183–1194.
46. Abdallah FW, et al. Ketamine reduces postoperative pain and inflammatory markers after major abdominal surgery. *Pain*. 2013;154(12):2827–2833.
47. Khalili-Mahani N, Martini CH, Olofsen E, et al. Effect of subanaesthetic ketamine on plasma and saliva cortisol secretion. *Br J Anaesth*. 2015;115(1):68-75.
48. Niesters M, Martini C, Dahan A. Ketamine for chronic pain: mechanisms and clinical implications. *Pain*. 2014;155(10):1973–1982. doi:10.1016/j.pain.2014.06.002.
49. Luggya T, Roche T, Ssemogerere L, et al. Effect of low-dose ketamine on postoperative serum IL-6 production among elective surgical patients: a randomized clinical trial. *BMC Anesthesiol*. 2017;17:50. doi:10.1186/s12871-017-0500-x.
50. Holbrook A, Trepanier M, Boulanger A, et al. Perioperative ketamine and attenuation of hypothalamic–pituitary–adrenal axis activation in major surgery. *J Cardiothorac Vasc Anesth*. 2018;32(5):2181–2188.
51. De Kock M, Lavand'homme P, Waterloos H. “Balanced analgesia” and perioperative cytokine modulation: role of NMDA antagonism and inflammatory pathways. *Anesthesiology*. 2013;119(6):1484–1496.
52. Cooke, C., McNicol, E.D., Ferguson, M.C., Minami, T. and Weaver, M.L. (2019) 'Intravenous lidocaine for the management of perioperative pain: A systematic review and meta-analysis', *Anesthesia and Analgesia*, 128(6), pp. 1200–1213. <https://doi.org/10.1213/ANE.0000000000004058>.
53. Herroeder, S., Pecher, S., Schönherr, M.E., Kaulitz, G., Hahnenkamp, K., Friess, H., Böttiger, B.W., Bauer, H., Dijkgraaf, M.G., Durieux, M.E. and Hollmann, M.W. (2007) 'Systemic lidocaine shortens length of hospital stay after colorectal surgery: A double-blinded, randomized, placebo-controlled trial', *Annals of Surgery*, 246(2), pp. 192–200. <https://doi.org/10.1097/SLA.0b013e31805dac11>
54. Amr YM, Yousef AA. Evaluation of intravenous lidocaine and magnesium sulphate in treatment of postoperative pain after laparoscopic cholecystectomy: a randomized double-blind study. *Pain Med*. 2010;11(7):1200-1206.
55. Perrin-Terrin A, Pathak A, Auffray JL. Systemic lidocaine infusion in gynecological laparoscopic surgery: effect on pain and postoperative recovery. A double-blind,

- randomized trial. *J Minim Invasive Gynecol.* 2021;28(3):643–651. doi:10.1016/j.jmig.2020.10.005.
56. Dewinter G, Moens P, Fieuws S, Vanaudenaerde B, Van de Velde M, Rex S. Systemic lidocaine fails to improve postoperative morphine consumption, postoperative recovery and quality of life in patients undergoing posterior spinal arthrodesis. *Br J Anaesth.* 2016;116(5):672–679. doi:10.1093/bja/aew047.
 57. Laskowski K, Stirling A, McKay WP, Lim HJ. A systematic review of intravenous ketamine for postoperative analgesia. *Can J Anaesth.* 2011;58(10):911–923. doi:10.1007/s12630-011-9560-0.
 58. Xu Q, Zhang Y, Wang Y, et al. Low-dose ketamine for postoperative analgesia: a systematic review and meta-analysis of randomized controlled trials. *Pain Physician.* 2020;23(3):E257–E270.
 59. Macintyre PE, Schug SA, Scott DA, Visser EJ, Walker SM. *Acute pain management: scientific evidence.* 3rd ed. Melbourne: Australian and New Zealand College of Anaesthetists and Faculty of Pain Medicine; 2010.
 60. Grape S, Schug SA, Lirk P, et al. Perioperative ketamine reduces postoperative opioid consumption: a systematic review and meta-analysis. *Br J Anaesth.* 2022;128(3):e216–e230.
 61. Kuo CP, Jao SW, Chen KM, et al. Comparison of the effects of intravenous lidocaine and epidural analgesia on pain and recovery after abdominal hysterectomy. *Anesth Analg.* 2006;103(5):1391–1397.
 62. Dewinter G, Moens P, Fieuws S, Vanaudenaerde B, Van de Velde M, Rex S. Systemic lidocaine fails to improve postoperative morphine consumption, postoperative recovery and quality of life in patients undergoing posterior spinal arthrodesis. *Br J Anaesth.* 2016;116(5):672–679. doi:10.1093/bja/aew047.
 63. Baeriswyl M, Taffé P, Kirkham KR, Piubellini D, Moriggl B, Gautschi P, et al. Intravenous lidocaine infusion for postoperative pain control: a systematic review and meta-analysis of randomized controlled trials. *Reg Anesth Pain Med.* 2018;43(5):488–499.