

DAFTAR PUSTAKA

- American Diabetes Association. (2015) *Diabetes Care*. 38 : 8-16. Jakarta : EGC
- American Diabetes Association (2015) '*Diagnosis and Classification of Diabetes Melitus.*'. *Diabetes Care*. 34-40. Jakarta : EGC
- Azami. et al. (2018) 'Long Working Hours and Skipping Breakfast Concomitant with Late Evening Meals are Associated with Suboptimal Glycemic Control Among Japanese Patient with Type 2 Diabetes", doi: <https://doi.org/10.1111/jdi.12852>
- Buku Ajar Ilmu Penyakit Dalam. *Jilid III. Edisi V*. Jakarta : Balai Penerbit FKUI. 2014
- Fauci, A.S, et al (2008) '*Harrison Principles of Internal Medicine*'. USA : The McGraw-Hill Companies, Inc. pp.
- Gouda. et al (2018) 'Associations between eating habits and glycemic control and obesity in
- Internasional Diabetes Federation (2014) '*Diabetes Atlas*'. Available at: <https://www.idf.org/atlasmap/atlasmap>. (Accessed: 28 July 2019)
- Iwasaki, T. et al (2019) 'Association between eating behavior and poor glycemic control in Japanese adults', 2019. doi: <https://doi.org/10.1038/s41598-019-39001-y>
- Kementerian Kesehatan Republik Indonesia (2014) *Infodatin Diabetes Melitus*. Available at: <http://www.depkes.go.id/article/view/18110200003/infodatin-diabetes-melitus-2014.html> (Accessed: 28 July 2019).
- Kementerian Kesehatan Republik Indonesia (2019) *Potret sehat Indonesia dari Riskesdas 2018*. Available at: <http://www.depkes.go.id/article/view/18110200003/potret-sehat-indonesia-dari-riskesdas-2018.html> (Accessed: 28 July 2019).
- Megan. et al. (2020) 'Night eating in patients with type 2 diabetes. Associations with glycemic control, eating patterns, sleep, and mood, doi: 10.1016/j.appet.2014.04.009
- Mekary. et al. (2012) 'Eating patterns and type 2 diabetes risk in men: breakfast omission, eating frequency, and snacking', American Society for Nutrition, USA.
- Nakajima. et al. (2015) 'Association of hyperglycemia in a general Japanese population with late-night-dinner eating alone, but not breakfast skipping alone', doi: 10.1186/s40200-015-0147-0
- Nishiyama. et al. (2009) 'The Combined Unhealthy Behaviors of Breakfast Skipping and Smoking Are Associated with the Prevalence of Diabetes Melitus', Tochigi, Japan.
- PERKENI. (2015) *Konsensus Pengelolaan dan Pencegahan Diabetes Mellitus Tipe 2*

- Soegondo. et al. (2010) 'Penatalaksanaan Diabetes Melitus Terpadu, Sebagai Panduan Penatalaksanaan Bagi Dokter Maupun Edukator. Jakarta;FKUI.
- Takeshita. et al. (2020) 'Skipping breakfast, late-night eating and current smoking are associated with medication adherence in Japanese patients with diabetes, doi: 10.1016/j.pcd.2020.05.002
- Tjokroprawiro, et al. (2007) 'Ilmu Penyakit Dalam, Surabaya.Airlangga University.
- Uemura, M. et al. (2015) 'Breakfast Skipping is Positively Associated With Incidence of Type 2 Diabetes Mellitus: Evidence From the Aichi Workers', doi: <http://dx.doi.org/10.2188/jea.JE20140109>
- World Health Organization (2016) '*Global Report on Diabetes*'. Available at: <https://www.who.int/diabetes/global-report/en/>. (Accessed: 28 July 2019)
- WHO (2018) 'Obesity and overweight'. Available at: <https://www.who.int/news-room/fact-sheets/detail/obesity-and-overweight>
- World Health Organization (2018) 'Diabetes Melitus'. Available at: <https://www.who.int/news-room/fact-sheets/detail/DiabetesMelitus> (Accessed: 28 July 2019).
- Yaguchi. et al. (2020) 'Skipping breakfast, late-night eating and current smoking are associated with medication adherence in Japanese patients with diabetes, doi: 10.1016/j.pcd.2020.05.002

