

DAFTAR PUSTAKA

- Ahmed, S. B. and Ramesh, S. (2016) 'Sex hormones in women with kidney disease', *Nephrology Dialysis Transplantation*, 31(11), pp. 1787–1795. doi: 10.1093/ndt/gfw084.
- American Diabetes Association (2021) '11. chronic kidney disease and risk management: standards of medical care in diabetes—2022', *Diabetes Care*, 45(Supplement_1), S175–184. doi:10.2337/dc22-s011.
- Argyropoulos, C.P. et al. (2017) 'Rediscovering Beta-2 microglobulin as a biomarker across the spectrum of kidney diseases', *Frontiers in Medicine*, 4. doi:10.3389/fmed.2017.00073.
- Badro, D.A. (2023) 'Chronic kidney disease management in developing countries', *Handbook of Medical and Health Sciences in Developing Countries*, pp. 1–146. doi:10.1007/978-3-030-74786-2_27-1.
- Behairy, O.G. et al. (2017) 'Role of serum cystatin-C and Beta-2 microglobulin as early markers of renal dysfunction in children with beta thalassemia major', *International Journal of Nephrology and Renovascular Disease*, Volume 10, pp. 261–268. doi:10.2147/ijnrd.s142824.
- Bikbov, B. et al. (2020) 'Global, regional, and national burden of chronic kidney disease, 1990–2017: A systematic analysis for the global burden of disease study 2017', *The Lancet*, 395(10225), pp. 709–733. doi:10.1016/s0140-6736(20)30045-3.
- Bikbov, B., Perico, N. and Remuzzi, G. (2018) 'Disparities in chronic kidney disease prevalence among males and females in 195 countries: Analysis of the global burden of disease 2016 study', *Nephron*, 139(4), pp. 313–318. doi:10.1159/000489897.
- de Boer, I. H. et al. (2014) 'Renal outcomes in patients with type 1 diabetes and macroalbuminuria.', *Journal of the American Society of Nephrology: JASN*, 25(10), pp. 2342–2350. doi: 10.1681/ASN.2013091004.
- I.C. et al. (2019) 'Evaluation of beta 2-microglobulin, cystatin C, and calin-2 as renal biomarkers for patients with Fabry disease', *Nephron*, 139(4), pp. 217–227. doi:10.1159/000500570.



- Braga, P.C. et al. (2022) 'Mitochondrial pathophysiology on chronic kidney disease', *International Journal of Molecular Sciences*, 23(3), p. 1776. doi:10.3390/ijms23031776.
- Brenner, B. M. and Bazilinski, N. G. (2020) 'Brenner and Rector's the kidney', *JAMA- Journal of the American Medical Association-International Edition*, 284(1), p. 99.
- Chawla, L. S. et al. (2014) 'Acute kidney injury and chronic kidney disease as interconnected syndromes.', *The New England journal of medicine*, 371(1), pp. 58–66. doi: 10.1056/NEJMra1214243.
- Chen, T.K., Knicely, D.H. and Grams, M.E. (2019) 'Chronic kidney disease diagnosis and management', *JAMA*, 322(13), p. 1294. doi:10.1001/jama.2019.14745.
- Chida, S. et al. (2016) 'Levels of albuminuria and risk of developing macroalbuminuria in type 2 diabetes: historical cohort study', *Scientific Reports*, 6(1), p. 26380. doi: 10.1038/srep26380.
- Chu, C. D. et al. (2023) 'Estimated Prevalence and Testing for Albuminuria in US Adults at Risk for Chronic Kidney Disease', *JAMA Network Open*, 6(7), pp. e2326230–e2326230. doi: 10.1001/jamanetworkopen.2023.26230.
- Coresh, J., Grams, M.E. and Chen, T.K. (2021) 'Using GFR, albuminuria, and their changes in clinical trials and clinical care', *American Journal of Kidney Diseases*, 78(3), pp. 333–334. doi:10.1053/j.ajkd.2021.04.003.
- Decroli, E. (2019) in *Diabetes Melitus Tipe 2*. 1st edn. Padang: Pusat Penerbitan Bagian Ilmu Penyakit Dalam Fakultas Kedokteran Universitas Andalas, pp. 1–27.
- Egubve, A.O. et al. (2019) 'The role of urine albumin creatinine ratio and serum β 2 microglobulin as biomarkers of chronic kidney disease', *Universa Medicina*, 38(3), pp. 172–178. doi:10.18051/univmed.2019.v38.172-178.
- Estakhri, M.A. et al. (2018) 'An investigation of the relationship between beta-2 microglobulin (B2M) and inflammatory factors (serum levels of CRP and albumin) and high density lipoproteins (HDL) in hemodialysis patients', *Modern dical Laboratory Journal*, 1(1), pp. 17–22. doi:10.30699/mmlj17-01-04.
- M. et al. (2022) 'A Narrative Review of Chronic Kidney Disease in Clinical ctice: Current Challenges and Future Perspectives', *Advances in Therapy*, 1), pp. 33–43. doi: 10.1007/s12325-021-01927-z.



- Foster, M. C. *et al.* (2016) 'Serum β_2 -Trace Protein and β_2 -Microglobulin as Predictors of ESRD, Mortality, and Cardiovascular Disease in Adults With CKD in the Chronic Renal Insufficiency Cohort (CRIC) Study', *American Journal of Kidney Diseases*, 68(1), pp. 68–76. doi: 10.1053/j.ajkd.2016.01.015.
- Fox, C., Neuhaus, K. and Vassalotti, J. (2013) 'The importance urine albumin-creatinine ratio in the diagnosis and prognosis of chronic kidney disease', *OA Nephrology*, 1(3). doi:10.13172/2053-0293-1-3-1027.
- Grangé, G. *et al.* (2015) 'Fetal Beta2-Microglobulin as a Biomarker of Kidney Disease', in Patel, V. B. (ed.) *Biomarkers in Kidney Disease*. Dordrecht: Springer Netherlands, pp. 1–23. doi: 10.1007/978-94-007-7743-9_21-1.
- Hall, J.E. (2011) 'The Body Fluids and Kidneys', in Guyton and Hall Textbook of Medical Physiology . 12th edn. Philadelphia: Elsevier, pp. 304–307.
- Heerspink, H. J. L., Rabelink, T., & de Zeeuw, D. (2020) 'Pathophysiology of Proteinuria: Albuminuria as a Target for Treatment', *Chronic Renal Disease*, 211–224. doi:10.1016/b978-0-12-815876-0.00015-2
- Hill, N.R. *et al.* (2016) 'Global prevalence of chronic kidney disease – a systematic review and meta-analysis', *PLOS ONE*, 11(7). doi:10.1371/journal.pone.0158765.
- Hull, K.L. *et al.* (2021) 'Indications and considerations for kidney biopsy: An overview of clinical considerations for the non-specialist', *Clinical Medicine*, 22(1), pp. 34–40. doi:10.7861/clinmed.2021-0472.
- Hussein, Z., Abdulsattar, S. and Salman, I. (2022) 'The usefulness of serum beta-2 microglobulin as a biomarker for evaluating renal function decline in type II diabetes mellitus', *Baghdad Journal of Biochemistry and Applied Biological Sciences*, 3(01), pp. 51–59. doi:10.47419/bjbabs.v3i01.40.
- Jotwani, V. *et al.* (2020) 'Tubular Biomarkers and Chronic Kidney Disease Progression in SPRINT Participants.', *American journal of nephrology*, 51(10), pp. 797–805. doi: 10.1159/000509978.
- Kalantar-Zadeh, K. *et al.* (2021) 'Chronic kidney disease', *The Lancet*, 398(10302), 786–802. doi:10.1016/s0140-6736(21)00519-5.
- rian Kesehatan RI. (2017) 'Situasi Penyakit Ginjal Kronis', *InfoDATIN Pusat a dan Informasi Kementerian Kesehatan RI*. 2017, pp. 1–10.



- Keown, P.A. (2013) 'Predicting long-term outcome in renal transplantation', *Kidney International*, 84(4), pp. 650–652. doi:10.1038/ki.2013.275.
- Kovesdy, C.P. (2022) 'Epidemiology of chronic kidney disease: An update 2022', *Kidney International Supplements*, 12(1), pp. 7–11. doi:10.1016/j.kisu.2021.11.003.
- Koye, D.N. et al. (2018) 'The global epidemiology of diabetes and kidney disease', *Advances in Chronic Kidney Disease*, 25(2), pp. 121–132. doi:10.1053/j.ackd.2017.10.011.
- Kumar, M. et al. (2023) 'The Bidirectional Link Between Diabetes and Kidney Disease: Mechanisms and Management.', *Cureus*, 15(9), p. e45615. doi: 10.7759/cureus.45615.
- Li, L., Dong, M. and Wang, X.-G. (2016) 'The implication and significance of beta 2 microglobulin', *Chinese Medical Journal*, 129(4), pp. 448–455. doi:10.4103/0366-6999.176084.
- Lombardi, G. et al. (2024) 'Sex differences in chronic kidney disease–related complications and mortality across levels of glomerular filtration rate', *Nephrology Dialysis Transplantation*, 39(12), pp. 2005–2015. doi: 10.1093/ndt/gfae087.
- López-Novoa, J.M. et al. (2011) 'Etiopathology of chronic tubular, glomerular and renovascular nephropathies: Clinical implications', *Journal of Translational Medicine*, 9(1). doi:10.1186/1479-5876-9-13.
- Lousa, I. et al. (2020) 'New potential biomarkers for chronic kidney disease management—a review of the literature', *International Journal of Molecular Sciences*, 22(1), p. 43. doi:10.3390/ijms22010043.
- Lubis, R. and Thristy, I. (2023) 'Perbandingan Kadar Asam Urat Dan Laju Filtrasi Glomerulus (Lfg) Pada Pasien Gagal Ginjal Kronik Sebelum Dan Sesudah Hemodialisa', *Jurnal Ilmiah Kohesi*, 7(1), pp. 1–8. Available at: <https://kohesi.sciencemakarioz.org/index.php/JIK/article/download/382/387>.
- Lydia, A. (2023) 'Merawat Kesehatan Ginjal Generasi Muda Indonesia: Peran Reaksi Dini Glomerulonefritis'.



Mon, R.A. and Pincus, M.R. (2017) *Henry's clinical diagnosis and management by laboratory methods* (23rd ed.). Philadelphia, PA: Elsevier.

- Mumtaz, A. *et al.* (2010) 'Beta-2 microglobulin levels in hemodialysis patients.', *Saudi journal of kidney diseases and transplantation : an official publication of the Saudi Center for Organ Transplantation, Saudi Arabia*, 21(4), pp. 701–706.
- Pardede, S.O. (2016) 'Podosit Dan Slit diafragma serta perannya', *Sari Pediatri*, 6(3), p. 119. doi:10.14238/sp6.3.2004.119-24.
- Perazella, M.A. (2015) 'The urine sediment as a biomarker of kidney disease', *American Journal of Kidney Diseases*, 66(5), pp. 748–755. doi:10.1053/j.ajkd.2015.02.342.
- Pugh, D., Gallacher, P. J. and Dhaun, N. (2019) 'Management of Hypertension in Chronic Kidney Disease.', *Drugs*, 79(4), pp. 365–379. doi: 10.1007/s40265-019-1064-1.
- Rajput, S. *et al.* (2020) 'Diagnostic & Clinical Utilities of Serum β eta2 Microglobulin and Urinary Diagnostic & Clinical Utilities of Serum β eta 2 Microglobulin and Urinary Albumin Creatinine Ratio for Identification of Pre-Chronic Kidney Disease', (November), pp. 1–12.
- Ren, F. *et al.* (2021) 'Urine albumin-to-creatinine ratio within the normal range and risk of hypertension in the general population: A meta-analysis', *The Journal of Clinical Hypertension*, 23(7), pp. 1284–1290. doi:10.1111/jch.14263.
- Ren, J. and Dai, C. (2019) 'Pathophysiology of chronic kidney disease', *Chronic Kidney Disease*, pp. 13–32. doi:10.1007/978-981-32-9131-7_2.
- Rifai, N. (2022) 'Kidney Disease', in *Tietz Textbook of Laboratory Medicine*. 7th edn. Philadelphia: Elsevier, pp. 603-671.
- Romagnani, P. *et al.* (2017) 'Chronic kidney disease', *Nature Reviews Disease Primers*, 3(1). doi:10.1038/nrdp.2017.88.
- Sedighi, O., Abediankenari, S. and Omranifar, B. (2015) 'Association between plasma Beta-2 microglobulin level and cardiac performance in patients with chronic kidney disease.', *Nephro-urology monthly*, 7(1), p. e23563. doi: 10.5812/numonthly.23563.
- Stanga, Z. *et al.* (2013) 'Factors other than the glomerular filtration rate that ermine the serum beta-2-microglobulin level', *PLoS ONE*, 8(8). doi:10.1371/journal.pone.0072073.



- Stevens, P. E. *et al.* (2024) 'KDIGO 2024 Clinical Practice Guideline for the Evaluation and Management of Chronic Kidney Disease', *Kidney International*. United States, pp. S117–S314. doi: 10.1016/j.kint.2023.10.018.
- TopciuShufta, V. *et al.* (2016) 'Association of Beta-2 microglobulin with inflammation and dislipidemia in high-flux membrane hemodialysis patients', *Medical Archives*, 70(5), p. 348. doi:10.5455/medarh.2016.70.348-350.
- Vaidya, S.R. (2022) Chronic kidney disease, StatPearls [Internet]. Available at: <https://www.ncbi.nlm.nih.gov/books/NBK535404/> (Accessed: 18 February 2024).
- Webster, A.C. *et al.* (2017) 'Chronic kidney disease', *The Lancet*, 389(10075), pp. 1238–1252. doi:10.1016/s0140-6736(16)32064-5.
- Whittier, W.L. and Lewis, E. (2018) 'Development and progression of chronic kidney disease', National Kidney Foundation's Primer on Kidney Diseases [Preprint]. doi:10.1016/b978-0-323-47794-9.00051-2.
- Williams, M. E. (2013) 'Diabetic kidney disease in elderly individuals', *Medical Clinics*, 97(1), pp. 75–89.
- Wilson, S. *et al.* (2021) 'Chronic kidney disease: Definition, updated epidemiology, staging, and mechanisms of increased cardiovascular risk', *The Journal of Clinical Hypertension*, 23(4), pp. 831–834. doi:10.1111/jch.14186.
- Winchester, J.F., Salsberg, J.A. and Levin, N.W. (2003) 'Beta-2 microglobulin in ESRD: An in-depth review', *Advances in Renal Replacement Therapy*, 10(4), pp. 279–309. doi:10.1053/j.arrt.2003.11.003.
- Zainuddin, A. *et al.* (2022) Geospatial Analysis of Diabetes Type 2 and Hypertension in South Sulawesi, Indonesia. doi: 10.21203/rs.3.rs-2031474/v1.

