

DAFTAR PUSTAKA

1. De Vico P, Biasucci DG, Aversano L, Polidoro R, Zingaro A, Millarelli FR, et al. Feasibility and safety of deep sedation with propofol and remifentanyl in spontaneous breathing during endoscopic retrograde cholangiopancreatography: an observational prospective study. *BMC Anesthesiol.* 2023;23(1):1–8.
2. Salama HZ, Alnajjar YA, Owais TA, Jobran AWM, Safi R, Bahar M, et al. Endoscopic retrograde cholangiopancreatography utilisation and outcomes in the first advanced endoscopy centre in Palestine at Al-Ahli Hospital: A retrospective cohort study. *BMJ Open.* 2023;13(12):1–9.
3. Sarraj R, Theiler L, Vakilzadeh N, Krupka N, Wiest R. Propofol sedation in routine endoscopy: A case series comparing target controlled infusion vs manually controlled bolus concept . *World J Gastrointest Endosc.* 2024;16(1):11–7.
4. Simanjuntak VE, Oktaliansah E, Redjeki IS. Perbandingan Waktu Induksi, Perubahan Tekanan Darah, dan Pulih Sadar antara Total Intravenous Anesthesia Profopofol Target Controlled Infusion dan Manual Controlled Infusion. *J Anestesi Perioper.* 2013;1(3):158–66.
5. Chiang MH, Wu SC, You CH, Wu KL, Chiu YC, Ma CW, et al. Target-controlled infusion vs manually controlled infusion of propofol with alfentanil for bidirectional endoscopy: A randomized controlled trial. *Endoscopy.* 2013;45(11):907–14.
6. Motiaa Y, Bensghir M, Jaafari A, Meziane M, Ahtil R, Drissi Kamili N. Anesthesia for endoscopic retrograde cholangiopancreatography: Target-controlled infusion versus standard volatile anesthesia. *Ann Gastroenterol.* 2016;29(4):530–5.
7. Sahu A, Tripathy DK, Gupta B, Talawar P, Gupta R. Recovery time in target controlled infusion versus manual infusion of propofol in total intravenous anaesthesia for endoscopic retrograde cholangiopancreatography procedure using laryngeal mask airway-gastro: A

- randomised comparative study. *Indian J Anaesth.* 2019;49(4):257–62.
8. Godoroja-Diarto D, Constantin A, Moldovan C, Rusu E, Sorbello M. Efficacy and Safety of Deep Sedation and Anaesthesia for Complex Endoscopic Procedures—A Narrative Review. *Diagnostics.* 2022;12(7):1–12.
 9. American Society of Anesthesiologists. Statement on Continuum of Depth of Sedation: Definition of General Anesthesia and Levels of Sedation/Analgesia. *Comm Qual Manag Dep Adm.* 2019;
 10. Sahinovic MM, Struys MMRF, Absalom AR. Clinical Pharmacokinetics and Pharmacodynamics of Propofol. *Clin Pharmacokinet [Internet].* 2018;57(12):1539–58. Tersedia pada: <https://doi.org/10.1007/s40262-018-0672-3>
 11. Fan W, Zhu X, Wu L, Wu Z, Li D, Huang F, et al. Propofol: An anesthetic possessing neuroprotective effects. *Eur Rev Med Pharmacol Sci.* 2015;19(8):1520–9.
 12. Bas SS. Remifentanil-Propofol versus Fentanyl-Propofol for Circumcision Operations in Pediatric Ambulatory Surgical Patients. *Osmangazi J Med.* 2020;42(1):1–6.
 13. Jo YY, Kwak HJ. Sedation strategies for procedures outside the operating room. *Yonsei Med J.* 2019;60(6):491–9.
 14. Egbuta C, Mason KP. Current state of analgesia and sedation in the pediatric intensive care unit. *J Clin Med.* 2021;10(9).
 15. Su H, Eleveld DJ, Struys MMRF, Colin PJ. Mechanism-based pharmacodynamic model for propofol haemodynamic effects in healthy volunteers☆. *Br J Anaesth [Internet].* 2022;128(5):806–16. Tersedia pada: <https://doi.org/10.1016/j.bja.2022.01.022>
 16. Kupeli E, Feller-Kopman D, Mehta AC. Diagnostic Bronchoscopy. *Med Manag Thorac Surg Patient.* 2009;65–72.
 17. Hassan AH, Amer IA. Fentanyl versus Dexmedetomidine as an Adjuvant to Propofol for Fiberoptic Intubation in Patients with Temporomandibular Joint Ankylosis. *Egypt J Ear, Nose, Throat Allied Sci.* 2022;23(23):1–7.

18. Al-Rifai Z, Mulvey D. Principles of total intravenous anaesthesia: basic pharmacokinetics and model descriptions. *BJA Educ* [Internet]. 2016;16(3):92–7. Tersedia pada: <http://dx.doi.org/10.1093/bjaceaccp/mkv021>
19. Jasper A, Efe BP, Nosa EP. Total Intravenous Anaesthesia with Propofol-a comparison of Target-controlled infusion (TCI) with manual controlled infusion (MCI). A randomized study. *Authorea* [Internet]. 2022;(8):1–14. Tersedia pada: <https://www.authorea.com/doi/full/10.22541/au.165936244.49446058>
20. Anderson BJ, Bagshaw O. Practicalities of Total Intravenous Anesthesia and Target-controlled Infusion in Children. *Anesthesiology*. 2019;131(1):164–85.
21. Bajwa SJS, Vinayagam S, Shinde S, Dalal S, Vennel J, Nanda S. Recent advancements in total intravenous anaesthesia and anaesthetic pharmacology. *Indian J Anaesth*. 2022;49(4):257–62.
22. Anderson BJ, Houghton J. Total Intravenous Anesthesia and Target-Controlled Infusion [Internet]. Sixth Edit. *A Practice of Anesthesia for Infants and Children*. Elsevier Inc.; 2018. 177-198.e3 hal. Tersedia pada: <https://doi.org/10.1016/B978-0-323-42974-0.00008-2>
23. Gale T, Leslie K, Kluger M. Propofol anaesthesia via target controlled infusion or manually controlled infusion: Effects on the bispectral index as a measure of anaesthetic depth. *Anaesth Intensive Care*. 2001;29(6):579–84.
24. Absalom AR, Mani V, De Smet T, Struys MMRF. Pharmacokinetic models for propofol- Defining and illuminating the devil in the detail. *Br J Anaesth*. 2009;103(1):26–37.
25. Arslan G, Sezen Ö. Comparison of target controlled infusion and manual infusion of propofol for sedation in spinal anesthesia. *Ain-Shams J Anesthesiol*. 2020;12(1).
26. Laso LF, López-Picado A, de La Fuente EO, Murua AM, Sánchez-Castro C, Ruilope LP, et al. Manual vs. target-controlled infusion induction with

- propofol: An observational study. *Colomb J Anesthesiol* [Internet]. 2016;44(4):272–7. Tersedia pada: <http://dx.doi.org/10.1016/j.rcae.2016.06.005>
27. Russell D, Wilkes MP, Hunter SC, Glen JB, Hutton P, Kenny GNC. Manual compared with target-controlled infusion of propofol. *Br J Anaesth*. 1995;75(5):562–6.
 28. Yildirim SA, Dogan L, Sarikaya ZT, Ulugol H, Gucyetmez B, Toraman F. Hypotension after Anesthesia Induction: Target-Controlled Infusion Versus Manual Anesthesia Induction of Propofol. *J Clin Med*. 2023;12(16).
 29. Hsieh ML, Lu YT, Lin CC, Lee CP. Comparison of the target-controlled infusion and the manual infusion of propofol anesthesia during electroconvulsive therapy: an open-label randomized controlled trial. *BMC Psychiatry*. 2021;21(1):1–11.
 30. Yeganeh N, Roshani B, Latifi H, Almasi A. Comparison of target-controlled infusion of sufentanil and remifentanil in blunting hemodynamic response to tracheal intubation. *J Inj Violence Res*. 2013;5(2):101–7.
 31. Mu J, Jiang T, Xu XB, Yuen VM, Irwin MG. Comparison of target-controlled infusion and manual infusion for propofol anaesthesia in children. *Br J Anaesth*. 2018;120(5):1049–55.
 32. Kateliya R, Madhukant, Dubey M, Chandra S, Sahay N. Comparison of recovery profiles in target-controlled infusions (TCI) versus manually controlled infusions for total intravenous anesthesia (TIVA) in laparoscopic surgeries. A randomized controlled trial. *J Anaesthesiol Clin Pharmacol*. 2023;34(3):46–50.
 33. Chen G, Buell O, Gruenewald M, Fudickar A, Scholz J, Bein B. A comparison between target-controlled and manually controlled propofol infusions in patients undergoing routine surgical procedures. *Eur J Anaesthesiol*. 2009;26(11):928–35.
 34. Leslie K, Clavisi O, Hargrove J. Target-controlled infusion versus manually-controlled infusion of propofol for general anaesthesia or sedation in adults. *Cochrane Database Syst Rev*. 2016;2016(7).

35. Passot S, Servin F, Allary R, Pascal J, Prades JM, Auboyer C, et al. Target-controlled versus manually-controlled infusion of propofol for direct laryngoscopy and bronchoscopy. *Anesth Analg*. 2002;94(5):1212–6.
36. Sharma S, Mudgal S, Thakur K, Gaur R. How to calculate sample size for observational and experiential nursing research studies? *Natl J Physiol Pharm Pharmacol*. 2019;10(0):1.
37. Fabus MS, Sleight JW, Warnaby CE. Effect of Propofol on Heart Rate and Its Coupling to Cortical Slow Waves in Humans. *Anesthesiology*. 2024;140(1):62–72.
38. Tantri AR, Sukmono RB, Atmadja LS. Comparison of recovery time with target controlled infusion of propofol with sevoflurane anesthesia using bispectral index monitoring in vitrectomy surgery. *Anaesthesia, Pain Intensive Care*. 2021;25(6):707–12.
39. Yeganeh N, Roshani B, Yari M, Almasi A. Target-controlled infusion anesthesia with propofol and remifentanyl compared with manually controlled infusion anesthesia in mastoidectomy surgeries. *Middle East J Anesthesiol*. 2010;20(6):785–93.
40. Wong SSC, Wang F, Chan TCW, Cheung CW. The analgesic effect of total intravenous anaesthesia with propofol versus inhalational anaesthesia for acute postoperative pain after hepatectomy: a randomized controlled trial. *BMC Anesthesiol* [Internet]. 2023;23(1):1–11. Tersedia pada: <https://doi.org/10.1186/s12871-023-02063-7>
41. Irwin MG, Chung CKE, Ip KY, Wiles MD. Influence of propofol-based total intravenous anaesthesia on peri-operative outcome measures: a narrative review. *Anaesthesia*. 2020;75(S1):e90–100.
42. Joe HB, Lee SY, Kim JS, Chang HS, Jeong Y, Jeong H, et al. Effect of total intravenous anaesthesia and prophylactic ramosetron on postoperative nausea and vomiting after thyroidectomy: A prospective, randomized controlled study. *J Int Med Res*. 2016;44(1):81–8.
43. Marsaban A, Sumaratih L, Pryambodho. Comparison between Intermittent Propofol Bolus Techniques and Target-Controlled Infusion in Patients

Underwent Gastrointestinal Endoscopy. *Indones J Gastroenterol Hepatol Dig Endosc.* 2013;14(3):132–8.

44. Lee JG, Yoo KS, Byun YJ. Continuous infusion versus intermittent bolus injection of propofol during endoscopic retrograde cholangiopancreatography. *Korean J Intern Med.* 2020;35(6):1338–45.



REKOMENDASI PERSETUJUAN ETIK

Nomor : 520/UN4.6.4.5.31/ PP36/ 2024

Tanggal: 4 Juli 2024

Dengan ini Menyatakan bahwa Protokol dan Dokumen yang Berhubungan Dengan Protokol berikut ini telah mendapatkan Persetujuan Etik :

No Protokol	UH24060384	No Sponsor	
Peneliti Utama	dr. Ira Ulil Inayah Wahid, S.Ked	Sponsor	
Judul Peneliti	PERBANDINGAN WAKTU INDUKSI, PERUBAHAN HEMODINAMIK, WAKTU PULIH SADAR, DAN KEBUTUHAN OBAT ANTARA TOTAL INTRAVENOUS ANESTHESIA PROPOFOL TARGET CONTROLLED INFUSION DAN MANUAL CONTROLLED INFUSION PADA PASIEN YANG MENJALANI TINDAKAN ENDOSCOPIC RETROGRADE CHOLA		
No Versi Protokol	2	Tanggal Versi	27 Juni 2024
No Versi PSP	2	Tanggal Versi	27 Juni 2024
Tempat Penelitian	RSUP. Dr. Wahidin Sudirohusodo Makassar		
Jenis Review	<input type="checkbox"/> Exempted <input type="checkbox"/> Expedited <input checked="" type="checkbox"/> Fullboard Tanggal 25 Juni 2024	Masa Berlaku 4 Juli 2024 sampai 4 Juli 2025	Frekuensi review lanjutan
Ketua KEP Universitas Hasanuddin	Prof. dr. Muh Nasrum Massi, PhD, SpMK, Subsp. Bakt(K)	Tanda tangan	
Sekretaris KEP Universitas Hasanuddin	dr. Firdaus Hamid, PhD, SpMK(K)	Tanda tangan	

Kewajiban Peneliti Utama:

- Menyerahkan Amandemen Protokol untuk persetujuan sebelum di implementasikan
- Menyerahkan Laporan SAE ke Komisi Etik dalam 24 Jam dan dilengkapi dalam 7 hari dan Lapo SUSAR dalam 72 Jam setelah Peneliti Utama menerima laporan
- Menyerahkan Laporan Kemajuan (progress report) setiap 6 bulan untuk penelitian resiko tinggi dan setiap setahun untuk penelitian resiko rendah
- Menyerahkan laporan akhir setelah Penelitian berakhir
- Melaporkan penyimpangan dari prokol yang disetujui (protocol deviation / violation)
- Mematuhi semua peraturan yang ditentukan

CURICULUM VITE

1. DATA PRIBADI

Nama Lengkap : dr. Ira Ulil Inayah Wahid
Tempat tanggal lahir : Bulukumba, 20 Februari 1989
Jenis Kelamin : Perempuan
Kewarganegaraan : Warga Negara Indonesia
Agama : Islam
Alamat : Royal Sentraland, Cluster East Sunderland E5-3,
Moncongloe, Maros
Email : ranayahwahid@gmail.com

2. RIWAYAT PENDIDIKAN

No	Jenjang Pendidikan	Institusi	Tahun	Tempat
1	SD	SD Negeri 7 Matajang	1995-2001	Bulukumba
2	SMP	SMP Negeri 1 Bulukumba	2001-2004	Bulukumba
3	SMA	SMA Negeri 17 Makassar	2004-2007	Makassar
4	S1 & Pofesi	FK Universitas Hasanuddin Makassar	2007-2013	Makassar
5	PPDS	Departemen Ilmu Anestesi, Terapi Intensif dan Manajemen Nyeri FK Universitas Hasanuddin	2020-2025	Makassar