

DAFTAR PUSTAKA

- Afifah, A.R., Liana, P., Fertilita, S., Salim, N.A., Hilda, F., Amin, C.A. & Umar, T.P. (2023), “C-Reactive Protein as The Predictor of Mortality for COVID-19 Patients in Indonesia”, Vol. 29 No. 2, pp. 180–184.
- Ali, N. (2020). Elevated level of C-reactive protein may be an early marker to predict risk for severity of COVID-19. In *Journal of Medical Virology* (Vol. 92, Issue 11, pp. 2409–2411). John Wiley and Sons Inc. <https://doi.org/10.1002/jmv.26097>
- Alwani, M., Yassin, A., Al-Zoubi, R.M., Aboumarzouk, O.M., Nettleship, J., Kelly, D., Al-Qudimat, A.R., et al. (2021), “Sex-based differences in severity and mortality in COVID-19.”, *Reviews in Medical Virology, England*, Vol. 31 No. 6, p. e2223.
- Amano, K., Maeda, I., Morita, T., Baba, M., Miura, T., Hama, T., Mori, I., et al. (2017), “C-reactive protein, symptoms and activity of daily living in patients with advanced cancer receiving palliative care”, *Journal of Cachexia, Sarcopenia and Muscle*, Vol. 8 No. 3, pp. 457–465.
- Ariani, P., Kurnia, D., Meinapuri, M., Effendi, R., Afriant, R. & Gustia, R. (2023), “Correlation between C-Reactive Protein level and outcome in coronavirus disease 2019 patients at covid intensive care unit RSUP. Dr. M. Djamil Padang”, *Jambi Medical Journal*, No. December 2021, pp. 268–278.
- Asghar, M.S., Haider Kazmi, S.J., Ahmed Khan, N., Akram, M., Ahmed Khan, S., Rasheed, U., Hassan, M., et al. (2020), “Clinical Profiles, Characteristics, and Outcomes of the First 100 Admitted COVID-19 Patients in Pakistan: A Single-Center Retrospective Study in a Tertiary Care Hospital of Karachi.”, *Cureus, United States*, Vol. 12 No. 6, p. e8712.
- Barek, M.A., Aziz, M.A. & Islam, M.S. (2020), “Impact of age, sex, comorbidities and clinical symptoms on the severity of COVID-19 cases: A meta-analysis with 55 studies and 10014 cases.”, *Heliyon, England*, Vol. 6 No. 12, p. e05684.
- Belice T, Demir I, & Yuksel A. (2020). Role of neutrophil-lymphocyte-ratio in the mortality of males diagnosed with COVID-19. In *Iranian Journal of Microbiology* (Vol. 296, Issue 2, pp. E15–E25). Radiological Society of North America Inc. <https://doi.org/10.1148/radiol.2020200490>
- Bienvenu, L.A., Noonan, J., Wang, X. & Peter, K. (2020), “Higher mortality of COVID-19 in males: Sex differences in immune response and cardiovascular comorbidities”, *Cardiovascular Research*, Vol. 116 No. 14, pp. 2197–2206.
- Buonacera, A., Stancanelli, B., Colaci, M., & Malatino, L. (2022). Neutrophil to Lymphocyte Ratio: An Emerging Marker of the Relationships between the Immune System and Diseases. In *International Journal of Molecular Sciences* (Vol. 23, Issue 7). MDPI. <https://doi.org/10.3390/ijms23073636>
- Burhan, E., Susanto, A. D., Nasution, S. A., Eka, G., Pitoyo, ceva W., Susilo, A., Firdaus, I., Santoso, A., Juzar, D. A., & Arif, S. K. (2022). Pedoman tatalaksana COVID-19 edisi 4.
- Cascella M; Rajnik M; Aleem. (2023). Features, Evaluation, and Treatment of Coronavirus (COVID-19) - StatPearls - NCBI Bookshelf. StatPearls - NCBI Bookshelf. <https://www.ncbi.nlm.nih.gov/books/NBK554776/>
- Chaturvedi, R., Lui, B., Aaronson, J.A., White, R.S. & Samuels, J.D. (2022), “COVID-19 complications in males and females: recent developments”, *Journal of Comparative Effectiveness Research*, Vol. 11 No. 9, pp. 689–698.
- Chowdhury, S. (2022). Epidemiology of COVID-19. *SpringerBriefs in Public Health*, 19–43. https://doi.org/10.1007/978-3-030-88766-7_2

- Eschborn, S., & Weitkamp, J. H. (2019). Procalcitonin versus C-reactive protein: review of kinetics and performance for diagnosis of neonatal sepsis. In *Journal of Perinatology* (Vol. 39, Issue 7, pp. 893–903). Nature Publishing Group. <https://doi.org/10.1038/s41372-019-0363-4>
- Farshbafnadi, M., Kamali Zonouzi, S., Sabahi, M., Dolatshahi, M. & Aarabi, M.H. (2021), “Aging & COVID-19 susceptibility, disease severity, and clinical outcomes: The role of entangled risk factors.”, *Experimental Gerontology*, England, Vol. 154, p. 111507.
- Forget, P., Khalifa, C., Defour, J.P., Latinne, D., Van Pel, M.C. & De Kock, M. (2017), “What is the normal value of the neutrophil-to-lymphocyte ratio?”, *BMC Research Notes*, BioMed Central, Vol. 10 No. 1, pp. 1–4.
- Gupta, A., Madhavan, M. V., Sehgal, K., Nair, N., Mahajan, S., Sehrawat, T.S., Bikdeli, B., et al. (2020), “Extrapulmonary manifestations of COVID-19”, *Nature Medicine*, Springer US, Vol. 26 No. 7, pp. 1017–1032.
- Huang, C., Wang, Y., Li, X., Ren, L., Zhao, J., Hu, Y., Zhang, L., Fan, G., Xu, J., Gu, X., Cheng, Z., Yu, T., Xia, J., Wei, Y., Wu, W., Xie, X., Yin, W., Li, H., Liu, M., ... Cao, B. (2020). Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *The Lancet*, 395(10223), 497–506. [https://doi.org/10.1016/S0140-6736\(20\)30183-5](https://doi.org/10.1016/S0140-6736(20)30183-5)
- Huang, Z., Fu, Z., Huang, W., & Huang, K. (2020). Prognostic value of neutrophil-to-lymphocyte ratio in sepsis: A meta-analysis. In *American Journal of Emergency Medicine* (Vol. 38, Issue 3, pp. 641–647). W.B. Saunders. <https://doi.org/10.1016/j.ajem.2019.10.023>
- Johns, I., Moschonas, K. E., Medina, J., Ossei-Gerning, N., Kassianos, G., & Halcox, J. P. (2018). Risk classification in primary prevention of CVD according to QRISK2 and JBS3 - ‘heart age’, and prevalence of elevated high-sensitivity C reactive protein in the UK cohort of the EURIKA study. *Open Heart*, 5(2). <https://doi.org/10.1136/openhrt-2018-000849>
- Karyono, D. R., & Wicaksana, A. L. (2020). Current prevalence, characteristics, and comorbidities of patients with COVID-19 in Indonesia. *Journal of Community Empowerment for Health*, 3(2), 77. <https://doi.org/10.22146/jcoemph.57325>
- Komazawa, O., Suriastini, N., ... I. W.-J. P. B., & 2021, U. (2021). Older People and COVID-19 in Indonesia. In *ERIA*.
- Lee, Y., McKechnie, T., Doumouras, A. G., Handler, C., Eskicioglu, C., Gmora, S., Anvari, M., & Hong, D. (2019). Diagnostic Value of C-Reactive Protein Levels in Postoperative Infectious Complications After Bariatric Surgery: a Systematic Review and Meta-Analysis. In *Obesity Surgery*. Springer New York LLC. <https://doi.org/10.1007/s11695-019-03832-5>
- Li, F., He, M., Zhou, M., Lai, Y., Zhu, Y., Liu, Z., Wang, Y., et al. (2023), “Association of C-reactive protein with mortality in Covid-19 patients: a secondary analysis of a cohort study”, *Scientific Reports*, Nature Publishing Group UK, Vol. 13 No. 1, pp. 1–10.
- Li, X., Li, T., & Wang, H. (2020). Treatment and prognosis of COVID 19: Current scenario and prospects (Review). *Experimental and Therapeutic Medicine*, 20(6), 1–1. <https://doi.org/10.3892/etm.2020.9435>
- Liu, T., Liang, W., Zhong, H., He, J., Chen, Z., He, G., Song, T., Chen, S., Wang, P., Li, J., Lan, Y., Cheng, M., Huang, J., Niu, J., Xia, L., Xiao, J., Hu, J., Lin, L., Huang, Q., ... Ma, W. (2020). Risk factors associated with COVID-19 infection: a retrospective cohort study based on contacts tracing. *Emerging Microbes and Infections*, 1546–1553. <https://doi.org/10.1080/22221751.2020.1787799>
- Man, M. A., Rajnoveanu, R. M., Motoc, N. S., Bondor, C. I., Chis, A. F., Lesan, A., Puiu, R., Lucaciu, S. R., Dantes, E., Gergely-Domokos, B., & Fira-Mladinescu, O. (2021).

- Neutrophil-to-lymphocyte ratio, platelets-to-lymphocyte ratio, and eosinophils correlation with high-resolution computer tomography severity score in COVID-19 patients. *PLoS ONE*, 16(6 June). <https://doi.org/10.1371/journal.pone.0252599>
- Molins, B., Figueras-roca, M., Valero, O., Sibila, O., Romero-va, S. & Ada, A. (2023), “C-reactive protein isoforms as prognostic markers of COVID-19 severity”, Vol. 2019 No. January, pp. 1–8.
- Mousavi-Nasab, S. D., Mardani, R., Azadani, H. N., Zali, F., Vasmehjani, A. A., Sabeti, S., Darazam, I. A., & Ahmadi, N. (2020). Neutrophil to lymphocyte ratio and C-reactive protein level as prognostic markers in mild versus severe COVID-19 patients. *Gastroenterology and Hepatology from Bed to Bench*, 13(4), 361–366. <https://doi.org/10.22037/ghfbb.v13i4.2109>
- Nehring, S. M., Goyal, A., Bhupendra, ;, & Affiliations, C. P. (2023). C Reactive Protein. <https://www.ncbi.nlm.nih.gov/books/NBK441843/?report=printable>
- Nori, W. (2022). C-Reactive protein role in assessing COVID-19 deceased geriatrics and survivors of severe and critical illness. *World Journal of Clinical Cases*, 10(30), 11210–11213. <https://doi.org/10.12998/wjcc.v10.i30.11210>
- Ortona, E., Pierdominici, M., & Rider, V. (2019). Editorial: Sex hormones and gender differences in immune responses. In *Frontiers in Immunology* (Vol. 10, Issue MAY). Frontiers Media S.A. <https://doi.org/10.3389/fimmu.2019.01076>
- Pakos, I.S., Lo, K.B., Salacup, G., Pelayo, J., Bhargav, R., Peterson, E., Gul, F., et al. (2020), “Characteristics of peripheral blood differential counts in hospitalized patients with COVID-19.”, *European Journal of Haematology*, England, Vol. 105 No. 6, pp. 773–778.
- Parasher, A. (2021). COVID-19: Current understanding of its Pathophysiology, Clinical presentation and Treatment. *Postgraduate Medical Journal*, 97(1147), 312–320. <https://doi.org/10.1136/postgradmedj-2020-138577>
- Peckham, H., de Gruijter, N.M., Raine, C., Radziszewska, A., Ciurtin, C., Wedderburn, L.R., Rosser, E.C., et al. (2020), “Male sex identified by global COVID-19 meta-analysis as a risk factor for death and ITU admission”, *Nature Communications*, Springer US, Vol. 11 No. 1, pp. 1–10.
- Perrotta, F., Corbi, G., Mazzeo, G., Boccia, M., Aronne, L., D’Agnano, V., Komici, K., Mazzarella, G., Parrella, R., & Bianco, A. (2020). COVID-19 and the elderly: insights into pathogenesis and clinical decision-making. *Aging Clinical and Experimental Research*, 32(8), 1599–1608. <https://doi.org/10.1007/s40520-020-01631-y>
- Pramono, A., Setiawan, Y. B., & Maryani, N. (2022). Risk Factors for Mortality in Indonesian COVID-19 Patients. *Open Access Macedonian Journal of Medical Sciences*, 9(T5), 181–184. <https://doi.org/10.3889/oamjms.2021.7826>
- Rotty, L., Kurube, J., Harijanto, P. N., Wantania, F., Haroen, H., Hendratta, C., Lasut, P., Kawengian, C., & Adiwinata, R. (2022). The Correlation between Neutrophil-to-Lymphocyte Ratio with C-reactive Protein and D-dimer Level among Indonesian COVID-19 Cases. *Open Access Macedonian Journal of Medical Sciences*, 10, 335–338. <https://doi.org/10.3889/oamjms.2022.8545>
- Sarkar, S., Khanna, P. & Singh, A.K. (2022), “The Impact of Neutrophil-Lymphocyte Count Ratio in COVID-19: A Systematic Review and Meta-Analysis.”, *Journal of Intensive Care Medicine*, United States, Vol. 37 No. 7, pp. 857–869.
- Setiadi, W., Rozi, I. E., Safari, D., Daningrat, W. O. D., Johar, E., Yohan, B., Yudhaputri, F. A., Lestari, K. D., Oktavianthi, S., Myint, K. S. A., Malik, S. G., & Soebandrio, A. (2022). Prevalence and epidemiological characteristics of COVID-19 after one year of pandemic in

- Jakarta and neighbouring areas, Indonesia: A single center study. *PLoS ONE*, 17(5 May), 1–9. <https://doi.org/10.1371/journal.pone.0268241>
- Setiati, S., & Azwar, M. K. (2020). COVID-19 and Indonesia. *Acta Medica Indonesiana*, 52(1), 84–89.
- Singhal, S., Kumar, P., Singh, S., Saha, S., & Dey, A. B. (2021). Clinical features and outcomes of COVID-19 in older adults : a systematic review and meta-analysis. 1–9.
- Singhal, T. (2020). Review on COVID-19. *The Indian Journal of Pediatrics*, 87(April), 281–286.
- Song, M., Graubard, B. I., Rabkin, C. S., & Engels, E. A. (2021). Neutrophil-to-lymphocyte ratio and mortality in the United States general population. *Scientific Reports*, 11(1). <https://doi.org/10.1038/s41598-020-79431-7>
- Sutaryono, S., Andasari, S. D., & Kasjono, H. S. (2020). Diagnosis and epidemiology of Coronavirus (COVID-19) outbreak in Indonesia. *Jurnal Teknologi Laboratorium*, 9(1), 49–57. <https://doi.org/10.29238/teknolabjournal.v9i1.222>
- Toori, K.U., Qureshi, M.A., Chaudhry, A. & Safdar, M.F. (2021), “Neutrophil to lymphocyte ratio (Nlr) in covid-19: A cheap prognostic marker in a resource constraint setting”, *Pakistan Journal of Medical Sciences*, Vol. 37 No. 5, pp. 1435–1439.
- Villoteau, A., Asfar, M., Otekpo, M., Loison, J., Gautier, J., & Annweiler, C. (2021). Elevated C-reactive protein in early COVID-19 predicts worse survival among hospitalized geriatric patients. *PLoS ONE*, 16(9 September). <https://doi.org/10.1371/journal.pone.0256931>
- WHO. (2021), “Living Guidance for clinical management of COVID-19”, World Health Organization, No. November, p. 63.
- Wolff, D., Nee, S., Hickey, N. S., & Marschollek, M. (2021). Risk factors for Covid-19 severity and fatality: a structured literature review. *Infection*, 49(1), 15–28. <https://doi.org/10.1007/s15010-020-01509-1>
- World Health Organization. (2023). Indonesia: WHO Coronavirus Disease (COVID-19) Dashboard With Vaccination Data | WHO Coronavirus (COVID-19) Dashboard With Vaccination Data. <https://covid19.who.int/region/searo/country/id>
- Yuki, K., Fujiogi, M., & Koutsogiannaki, S. (2020). COVID-19 pathophysiology: A review. *Clinical Immunology*, 215(April). <https://doi.org/10.1016/j.clim.2020.10842>
- Zahorec, R. (2021). Neutrophil-to-lymphocyte ratio, past, present and future perspectives. *Bratislava Medical Journal*, 122(7), 474–488. https://doi.org/10.4149/BLL_2021_078
- Zakianis, Adzania, F. H., Fauzia, S., Aryati, G. P., & Mahkota, R. (2021). Sociodemographic and environmental health risk factor of COVID-19 in Jakarta, Indonesia: An ecological study. *One Health*, 13(May), 100303. <https://doi.org/10.1016/j.onehlt.2021.100303>