

DAFTAR PUSTAKA

- Agarwal, A., & Nath, K. A. (2019). Pathophysiology of Chronic Kidney Disease Progression: Organ and Cellular Considerations. In *Chronic Renal Disease*. Elsevier Inc. <https://doi.org/10.1016/B978-0-12-815876-0.00018-8>
- Aina, F., Amin, Z., Sabri, S., Ismail, M., Chan, K. W., Ismail, N., Esa, N. M., Azmi, M., & Lila, M. (2020). Probiotic Properties of Bacillus Strains Isolated from Stingless Bee (*Heterotrigona itama*) Honey Collected across Malaysia. *International Journal of Environmental Research and Public Health*, 17(278), 1–15.
- Banoglu, E., Jha, G. G., & King, R. S. (2001). Hepatic microsomal metabolism of indole to indoxyl, a precursor of indoxyl sulfate. *European Journal of Drug Metabolism and Pharmacokinetics*, 26(4), 235–240. <https://doi.org/10.1007/BF03226377>
- Bao, N., Chen, F., & Dai, D. (2020). The Regulation of Host Intestinal Microbiota by Polyphenols in the Development and Prevention of Chronic Kidney Disease. *Frontiers in Immunology*, 10(January), 1–13. <https://doi.org/10.3389/fimmu.2019.02981>
- Baragetti I, De Simone I, Biazzi C, Buzzi L, Ferrario F, Luise MC, et al. The low-protein diet for chronic kidney disease: 8 years of clinical experience in a nephrology ward. *Clin Kidney J*. 2020;13(5):253-260
- Biluca FC, da Silva B, Caon T, Mohr ETB, Vieira GN, Gonzaga LV, Vitali L, Micke G, Fett R, Dalmarco EM, Costa ACO. (2020). Investigation of phenolic compounds, antioxidant and anti-inflammatory activities in stingless bee honey (Meliponinae). *Food Res Int*. 129:108756. doi: 10.1016/j.foodres.2019.108756.
- Black AP, Anjos JS, Cardozo L, Carmo FL, Dolenga CJ, Nakao LS, de Carvalho Ferreira D, Rosado A, Carraro Eduardo JC, Mafra D. (2018). Does Low-Protein Diet Influence the Uremic Toxin Serum Levels From the Gut Microbiota in Nondialysis Chronic Kidney Disease Patients? *J Ren Nutr*, 28(3):208-214. doi: 10.1053/j.jrn.2017.11.007.
- Bogdanov, S., Jurendic, T., Sieber, R., & Gallmann, P. (2008). Honey for nutrition and health: A review. *Journal of the American College of Nutrition*, 27(6), 677–689. <https://doi.org/10.1080/07315724.2008.10719745>
- Borges, N. A., Barros, A. F., Nakao, L. S., Dolenga, C. J., Fouque, D., & Mafra, D. (2016). Protein-Bound Uremic Toxins from Gut Microbiota and Inflammatory Markers in Chronic Kidney Disease. *Journal of Renal Nutrition*, 26(6), 396–400. <https://doi.org/10.1053/j.jrn.2016.07.005>
- Buketov, A. V., & Krasnen'Ky, V. M. (2013). Influence of the range of

particle sizes of the filler on the adhesion strength and residual stresses in epoxy composites. *Materials Science*, 48(4), 521–530. <https://doi.org/10.1007/s11003-013-9533-2>

Cano, R.; Bermúdez, V.; Galban, N.; Garrido, B.; Santeliz, R.; Gotera, M.P.; Duran, P.; Boscan, A.; Carbonell-Zabaleta, A.-K.; Durán-Agüero, S.; et al. Dietary Polyphenols and Gut Microbiota Cross-Talk: Molecular and Therapeutic Perspectives for Cardiometabolic Disease: A Narrative Review. *Int. J. Mol. Sci.* 2024, 25, 9118. <https://doi.org/10.3390/ijms25169118>

Catalkaya, G., Venema, K., Lucini, L., Rocchetti, G., Delmas, D., Daglia, M., De Filippis, A., Xiao, H., Quiles, J. L., Xiao, J., & Capanoglu, E. (2020). Interaction of dietary polyphenols and gut microbiota: Microbial metabolism of polyphenols, influence on the gut microbiota, and implications on host health. *Food Frontiers*, 1(2), 109–133. <https://doi.org/10.1002/fft2.25>

Chen, T. K., Knicely, D. H., & Grams, M. E. (2019). Chronic Kidney Disease Diagnosis and Management: A Review. *JAMA - Journal of the American Medical Association*, 322(13), 1294–1304. <https://doi.org/10.1001/jama.2019.14745>

Cheng, T. H., Ma, M. C., Liao, M. T., Zheng, C. M., Lu, K. C., Liao, C. H., Hou, Y. C., Liu, W. C., & Lu, C. L. (2020). Indoxyl Sulfate, a Tubular Toxin, Contributes to the Development of Chronic Kidney Disease. *Toxins*, 12(11), 1–15. <https://doi.org/10.3390/toxins12110684>

Duda-Chodak A, Tarko T, Satora P, et al. Interaction of dietary compounds, especially polyphenols, with the intestinal microbiota: a review. *Eur J Nutr.* 2015;54(3):325– 341. doi: 10.1007/s00394-015-0852-y

Felizardo, R. J. F., Mizuno Watanabe, I. K., Dardi, P., Venturini Rossoni, L., & Olsen Saraiva Câmara, N. (2019). The interplay among gut microbiota, hypertension and kidney diseases: The role of short-chain fatty acids. *Pharmacological Research*, 141, 366–377. <https://doi.org/10.1016/j.phrs.2019.01.019>

Ferreira, E., Cendoroglo, M., Manfredi, S. R., Canziani, M. E., Marie, B., Quinto, R., Grabulosa, C. C., Guimarães-souza, N. K., Peres, A. T., Tarcísio, J., Carvalho, G. De, Batista, M. C., & Dalboni, M. A. (2014). *Effect of Indoxyl Sulfate on Oxidative Stress, Apoptosis, and Monocyte Chemoattractant Protein-1 in Leukocytes.* 2014, 26–28.

Firrman J, Liu L, Zhang L, et al. The effect of quercetin on genetic expression of the commensal gut microbes *Bifidobacterium catenulatum*, *Enterococcus caccae* and *Ruminococcus gauvreauii*. *Anaerobe.* 2016;42:130–141. doi: 10.1016/j.anaerobe.2016.10.004

- Goldstein-Fuchs J, LaPierre AF. Nutrition and Kidney Disease. In: Gilbert SJ, Weiner DE, editors. National Kidney Foundation Primer on Kidney Diseases. 6th ed. Philadelphia: W.B. Saunders; 2014. p. 467-75.
- Guarino, M. P. L., Altomare, A., Emerenziani, S., Di Rosa, C., Ribolsi, M., Balestrieri, P., Iovino, P., Rocchi, G., & Cicala, M. (2020). Mechanisms of action of prebiotics and their effects on gastro-intestinal disorders in adults. *Nutrients*, 12(4), 1–24. <https://doi.org/10.3390/nu12041037>
- Hashim, K.-N., Chin, K.-Y., & Ahmad, F. (2021). The Mechanism of Honey in Reversing Metabolic Syndrome. *Molecules*, 26(4), 808. <https://doi.org/10.3390/molecules26040808>
- Hassan, M. S., Ahmed, Y. S., Sarhaan, E. I., Mehanna, N. S., Madbouli, N. N., & Abdelgawad, M. A. (2022). Effect of dietary synbiotic supplementation on serum indoxyl sulfate in prevalent hemodialysis patients. *The Egyptian Journal of Internal Medicine*, 34(1), 1–7. <https://doi.org/10.1186/s43162-021-00096-3>
- Hobby, G. P., Karaduta, O., Dusio, G. F., Singh, M., Zybaïlov, B. L., & Arthur, J. M. (2019). Chronic kidney disease and the gut microbiome. *American Journal of Physiology - Renal Physiology*, 316(6), F1211–F1217. <https://doi.org/10.1152/ajprenal.00298.2018>
- Holle, J., Kirchner, M., Okun, J., Bayazit, A. K., Obrycki, L., Canpolat, N., Bulut, I. K., Azukaitis, K., Duzova, A., Ranchin, B., Shroff, R., Candan, C., Oh, J., Klaus, G., Lugani, F., Gimpel, C., Buscher, R., Yilmaz, A., Baskin, E., ... Schaefer, F. (2020). Serum indoxyl sulfate concentrations associate with progression of chronic kidney disease in children. *PLoS ONE*, 15(10 October), 1–17. <https://doi.org/10.1371/journal.pone.0240446>
- Hung, S. C., Kuo, K. L., Wu, C. C., & Tarng, D. C. (2017). Indoxyl sulfate: A novel cardiovascular risk factor in chronic kidney disease. *Journal of the American Heart Association*, 6(2), 1–8. <https://doi.org/10.1161/JAHA.116.005022>
- Ismail, M. M. (2017). Honey...Prebiotic and Antibiotic. *EC Microbiology*, 85, 243–245.
- Jan Mei, S., Mohd Nordin, M. S., & Norrakiah, A. S. (2010). Fructooligosaccharides in honey and effects of honey on growth of *Bifidobacterium longum* BB 536. *International Food Research Journal*, 17(3), 557–561.
- Jia, L., Jia, Q., Yang, J., Jia, R., & Zhang, H. (2018). Efficacy of probiotics supplementation on chronic kidney disease: A systematic review and meta-analysis. *Kidney and Blood Pressure Research*, 43(5), 1623–1635. <https://doi.org/10.1159/000494677>

- KDIGO 2024 Clinical Practice Guideline for the Evaluation and Management of Chronic Kidney Disease. *Kidney Int* 2024;105:A1.
- Kemenkes RI. (2023). Survei Kesehatan Indonesia Tahun 2023. *Kementrian Kesehatan RI*, 53(9), 1689–1699.
- Kovesdy. (2022). Epidemiology of chronic kidney disease: an update 2022. *Kidney international Supplements*, (2022)12,7-11. <https://doi.org/10.1016/j.kisu.2021.11.003>
- Lashani, E., Davoodabadi, A., & Dallal, M. M. S. (2020). Some probiotic properties of lactobacillus species isolated from honey and their antimicrobial activity against foodborne pathogens. *Veterinary Research Forum*, 11(2), 121–126. <https://doi.org/10.30466/vrf.2018.90418.2188>
- Lau, W. L., Savoj, J., Nakata, M. B., & Vaziri, N. D. (2018). Altered microbiome in chronic kidney disease: Systemic effects of gut-derived uremic toxins. *Clinical Science*, 132(5), 509–522. <https://doi.org/10.1042/CS20171107>
- Liang Y, Meng Z, Ding XL, Jiang M. Effects of proton pump inhibitors on inflammatory bowel disease: An updated review. *World J Gastroenterol* 2024; 30(21): 2751-2762 [PMID: [38899331](https://pubmed.ncbi.nlm.nih.gov/38899331/)] DOI: [10.3748/wjg.v30.i21.2751](https://doi.org/10.3748/wjg.v30.i21.2751)]
- Li, D. Y., & Tang, W. H. W. (2018). Contributory Role of Gut Microbiota and Their Metabolites Toward Cardiovascular Complications in Chronic Kidney Disease. *Seminars in Nephrology*, 38(2), 193–205. <https://doi.org/10.1016/j.semnephrol.2018.01.008>
- Li, F. X., Wang, M. H., Wang, J. P., Li, R. S., & Zhang, Y. Q. (2019). Alterations to the gut microbiota and their correlation with inflammatory factors in chronic kidney disease. *Frontiers in Cellular and Infection Microbiology*, 9(JUN). <https://doi.org/10.3389/fcimb.2019.00206>
- Liu, W. C., Tomino, Y., & Lu, K. C. (2018). Impacts of indoxyl sulfate and p-Cresol sulfate on chronic kidney disease and mitigating effects of AST-120. *Toxins*, 10(9), 1–22. <https://doi.org/10.3390/toxins10090367>
- Li Q, Van de Wiele T. Gut microbiota as a driver of the interindividual variability of cardiometabolic effects from tea polyphenols. *Crit Rev Food Sci Nutr*. 2023;63(11):1500–1526. doi: 10.1080/10408398.2021.1965536.
- Ma, G., & Chen, Y. (2020). Polyphenol supplementation benefits human health via gut microbiota: A systematic review via meta-analysis. *Journal of Functional Foods*, 66, 103829.
- Mafra, D., Borges, N., Alvarenga, L., Esgalhado, M., Cardozo, L., Lindholm, B., & Stenvinkel, P. (2019). Dietary components that may influence the

- disturbed gut microbiota in chronic kidney disease. *Nutrients*, 11(3), 1–23. <https://doi.org/10.3390/nu11030496>
- Magliocca, G.; Mone, P.; Di Iorio, B.R.; Heidland, A.; Marzocco, S. Short-Chain Fatty Acids in Chronic Kidney Disease: Focus on Inflammation and Oxidative Stress Regulation. *Int. J. Mol. Sci.* **2022**, 23, 5354. <https://doi.org/10.3390/ijms23105354>
- Mahmoodpoor, F., Rahbar Saadat, Y., Barzegari, A., Ardalan, M., & V Zununi Vahed, S. (2017). The impact of gut microbiota on kidney function and pathogenesis. *Biomedicine and Pharmacotherapy*, 93, 412–419. <https://doi.org/10.1016/j.biopha.2017.06.066>
- Marzocco S, Dal Piaz F, Di Micco L, Torraca S, Sirico ML, Tartaglia D, Autore G, Di Iorio B. (2020). Very low protein diet reduces indoxyl sulfate levels in chronic kidney disease. *Blood Purif.* 35(1-3):196-201. doi: 10.1159/000346628.
- Meijers, B., Evenepoel, P., & Anders, H. J. (2019). Intestinal microbiome and fitness in kidney disease. *Nature Reviews Nephrology*, 15(9), 531–545. <https://doi.org/10.1038/s41581-019-0172-1>
- Meijers, B., Farré, R., Dejongh, S., Vicario, M., & Evenepoel, P. (2018). Intestinal barrier function in chronic kidney disease. *Toxins*, 10(7), 1–11. <https://doi.org/10.3390/toxins10070298>
- Miguel, M. G., Antunes, M. D., & Faleiro, M. L. (2017). Honey as a complementary medicine. *Integrative Medicine Insights*, 12, 1–15. <https://doi.org/10.1177/1178633717702869>
- Milik, A., & Hryniewicz, E. (2014). On translation of LD, IL and SFC given according to IEC-61131 for hardware synthesis of reconfigurable logic controller. *IFAC Proceedings Volumes (IFAC-PapersOnline)*, 19(1), 4477–4483. <https://doi.org/10.3182/20140824-6-za-1003.01333>
- Milovanova L, Fomin V, Moiseev S, Taranova M, Milovanov Y, Lysenko Kozlovskaya L, et al. Effect of essential amino acid ketoanalogues and protein restriction diet on morphogenetic proteins (FGF-23 and Klotho) in 3b-4 stages chronic kidney disease patients: a randomized pilot study. *Clin Exp Nephrol.* 2018; 22: 1351-9.
- Mitrović M, Stanković-Popović V, Tolinački M, Golić N, Soković Bajić S, Veljović K, Nastasijević B, Soldatović I, Svorcan P, Dimković N. The Impact of Synbiotic Treatment on the Levels of Gut-Derived Uremic Toxins, Inflammation, and Gut Microbiome of Chronic Kidney Disease Patients-A Randomized Trial. (2023). *J Ren Nut* ;33(2):278-288. doi: 10.1053/j.jrn.2022.07.008
- Mohan, A., Quek, S.-Y., Gutierrez-Maddox, N., Gao, Y., & Shu, Q. (2017a). Effect of honey in improving the gut microbial balance. *Food Quality*

- and Safety*, 1(May), 107–115. <https://doi.org/10.1093/fqs/fyx015>
- Moradi, H., Sica, D. A., & Kalantar-Zadeh, K. (2013). Cardiovascular burden associated with uremic toxins in patients with chronic kidney disease. *American Journal of Nephrology*, 38(2), 136–148. <https://doi.org/10.1159/000351758>
- Nishiyama K, Aono K, Fujimoto Y, Kuwamura M, Okada T, Tokumoto H, Izawa T, Okano R, Nakajima H, Takeuchi T, Azuma YT. (2019). Chronic kidney disease after 5/6 nephrectomy disturbs the intestinal microbiota and alters intestinal motility. *J Cell Physiol*. 234(5):6667-6678. doi: 10.1002/jcp.27408.
- Orenes-piñero, E., Ondrussek-sekac, M., Navas-carrillo, D., Orenes-piñero, E., Navas-carrillo, D., & Orenes-, E. (2020). Intestinal microbiota alterations in chronic kidney disease and the influence Intestinal microbiota alterations in chronic kidney disease and the influence of dietary components. *Critical Reviews in Food Science and Nutrition*, 0(0), 1–13. <https://doi.org/10.1080/10408398.2020.1761771>
- Pan, W., & Kang, Y. (2018). Gut microbiota and chronic kidney disease: implications for novel mechanistic insights and therapeutic strategies. *International Urology and Nephrology*, 50(2), 289–299. <https://doi.org/10.1007/s11255-017-1689-5>
- Plata, C., Cruz, C., Cervantes, L. G., & Ramírez, V. (2019). The gut microbiota and its relationship with chronic kidney disease. *International Urology and Nephrology*, 51(12), 2209–2226. <https://doi.org/10.1007/s11255-019-02291-2>
- Ramezani, A., Massy, Z. A., Meijers, B., Evenepoel, P., Vanholder, R., & Raj, D. S. (2016). Role of the gut microbiome in Uremia: A potential therapeutic target. *American Journal of Kidney Diseases*, 67(3), 483–498. <https://doi.org/10.1053/j.ajkd.2015.09.027>
- Ranneh Y, Akim AM, Hamid HA, Khazaai H, Fadel A, Zakaria ZA, Albujja M, Bakar MFA. (2021). Honey and its nutritional and anti-inflammatory value. *BMC Complement Med Ther*. 21(1):30. doi: 10.1186/s12906-020-03170-5.
- Rao, P. V., Krishnan, K. T., Salleh, N., & Gan, S. H. (2016). Biological and therapeutic effects of honey produced by honey bees and stingless bees: A comparative review. *Revista Brasileira de Farmacognosia*, 26(5), 657–664. <https://doi.org/10.1016/j.bjp.2016.01.012>
- Rossi M, Campbell KL, Johnson DW, Stanton T, Vesey DA, Coombes JS, Weston KS, Hawley CM, McWhinney BC, Ungerer JP, Isbel N. (2014). Protein-bound uremic toxins, inflammation and oxidative stress: a cross-sectional study in stage 3-4 chronic kidney disease. *Arch Med Res*; 45(4):309-17. doi: 10.1016/j.arcmed.2014.04.002.

- Rukavina Mikusic, N. L., Kouyoumdzian, N. M., & Choi, M. R. (2020). Gut microbiota and chronic kidney disease: evidences and mechanisms that mediate a new communication in the gastrointestinal-renal axis. *Pflugers Archiv European Journal of Physiology*, 472(3), 303–320. <https://doi.org/10.1007/s00424-020-02352-x>
- Sabatino, A., Regolisti, G., Brusasco, I., Cabassi, A., Morabito, S., & Fiaccadori, E. (2015). Alterations of intestinal barrier and microbiota in chronic kidney disease. *Nephrology Dialysis Transplantation*, 30(6), 924–933. <https://doi.org/10.1093/ndt/gfu287>
- Sampaio-Maia, B., Simões-Silva, L., Pestana, M., Araujo, R., & Soares-Silva, I. J. (2016). The Role of the Gut Microbiome on Chronic Kidney Disease. *Advances in Applied Microbiology*, 96, 65–94. <https://doi.org/10.1016/bs.aambs.2016.06.002>
- shin, H.-S., & Ustunol, Z. (2005). Carbohydrate composition of honey from different floral sources and their influence on growth of selected intestinal bacteria: An in vitro comparison. *Food Research International*, 38(6), 721–728.
- Sircana, A., De Michieli, F., Parente, R., Framarin, L., Leone, N., Berrutti, M., Paschetta, E., Bongiovanni, D., & Musso, G. (2019). Gut microbiota, hypertension and chronic kidney disease: Recent advances. *Pharmacological Research*, 144, 390–408. <https://doi.org/10.1016/j.phrs.2018.01.013>
- Sirich, T. L., Plummer, N. S., Gardner, C. D., Hostetter, T. H., & Meyer, T. W. (2014). Effect of increasing dietary fiber on plasma levels of colon-derived solutes in hemodialysis patients. *Clinical Journal of the American Society of Nephrology*, 9(9), 1603–1610. <https://doi.org/10.2215/CJN.00490114>
- So D, Whelan K, Rossi M, Morrison M, Holtmann G, Kelly JT, Shanahan ER, Staudacher HM, Campbell KL. (2018). Dietary fiber intervention on gut microbiota composition in healthy adults: a systematic review and meta-analysis. *Am J Clin Nutr*, 107(6):965-983. doi: 10.1093/ajcn/nqy041.
- Sumida, K., & Kovesdy, C. P. (2019). The gut-kidney-heart axis in chronic kidney disease. *Physiology International*, 106(3), 195–206. <https://doi.org/10.1556/2060.106.2019.19>
- Syamsul, T. D., Natzir, R., & As, S. (2021). Phenolic Profile and Antioksidant Activity of Indonesia. *Medico-Legal Update*, 21(2), 1199–1203.
- Tan, X., Cao, X., Zou, J., Shen, B., Zhang, X., Liu, Z., Lv, W., Teng, J., & Ding, X. (2017). Indoxyl sulfate, a valuable biomarker in chronic kidney disease and dialysis. *Hemodialysis International*, 21(2), 161–167. <https://doi.org/10.1111/hdi.12483>

- Tayebi-Khosroshahi, H.; Habibzadeh, A.; Niknafs, B.; Ghotaslou, R.; Yeganeh Sefidan, F.; Ghojatzadeh, M.; Moghaddaszadeh, M.; Parkhide, S. The effect of lactulose supplementation on fecal microflora of patients with chronic kidney disease; a randomized clinical trial. *J. Ren. Inj. Prev.* 2016, 5, 162–167. <https://doi.org/10.15171/jrip.2016.34>.
- Tian, N., Li, L., Ng, J. K. C., & Li, P. K. T. (2022). The Potential Benefits and Controversies of Probiotics Use in Patients at Different Stages of Chronic Kidney Disease. *Nutrients*, 14(19), 1–12. <https://doi.org/10.3390/nu14194044>
- Vanholder, R., Pletinck, A., & Schepers, E. (2018). *Biochemical and Clinical Impact of Organic Uremic Retention Solutes: A Comprehensive Update*. 1–57. <https://doi.org/10.3390/toxins10010033>
- Venereol, I. D., Rold, R., Puig, S., & Malveyh, J. (2012). *in C Er Ig E C Er*. 147(2), 149–159.
- Vit, P., Roubik, D. W., & Pedro, S. R. M. (2012). Pot-Honey: A legacy of stingless bees. *Pot-Honey: A Legacy of Stingless Bees, January 2013*, 1–654. <https://doi.org/10.1007/978-1-4614-4960-7>
- Wang, F., Jiang, H., Shi, K., Ren, Y. I., Zhang, P. A. N., & Cheng, S. (2012). *Gut bacterial translocation is associated with microinflammation in end-stage renal disease patients Correspondence : 17*(March 2010), 733–738. <https://doi.org/10.1111/j.1440-1797.2012.01647.x>
- Wong, J., Piceno, M., & Desantis, T. Z. (2014). *Expansion of Urease- and Uricase-Containing , Indole- and p-Cresol-Forming and Contraction of Short-Chain Fatty Acid-Producing Intestinal Microbiota in ESRD*. 230–237. <https://doi.org/10.1159/000360010>
- Wu, M., Cai, X., Lin, J., Zhang, X., Scott, E. M., & Li, X. (2019). Association between fibre intake and indoxyl sulphate/P-cresyl sulphate in patients with chronic kidney disease: Meta-analysis and systematic review of experimental studies. *Clinical Nutrition*, 38(5), 2016–2022. <https://doi.org/10.1016/j.clnu.2018.09.015>
- Yang, C. Y., & Tarng, D. C. (2018). Diet, gut microbiome and indoxyl sulphate in chronic kidney disease patients. *Nephrology*, 23, 16–20. <https://doi.org/10.1111/nep.13452>
- Zhang, L. S., & Davies, S. S. (2016). Microbial metabolism of dietary components to bioactive metabolites: Opportunities for new therapeutic interventions. *Genome Medicine*, 8(1), 1–18. <https://doi.org/10.1186/s13073-016-0296-x>