

Daftar Pustaka

- Alves, R. & Grimalt, R., 2018. A review of platelet-rich plasma: history, biology, mechanism of action, and classification. *Skin Appendage Disorders*, 4(1), pp.18–24.
- Bayat, M., et al., 2023. Comparison of dextrose proloterapi and triamcinolone intraarticular injection on pain and function in patients with knee osteoarthritis: A randomised clinical trial. *Anesthesiology and Pain Medicine*.
- Benazzoug, Y., et al., 1998. Effect of high glucose concentrations on the expression of collagens by fibroblasts. *Connective Tissue Research*, 38(1–4), pp.xxx–xxx.
- Bertrand, H., Reeves, K. D., Bennett, C. J., Bicknell, S., & Cheng, A. L. (2016). Dextrose prolotherapy versus control injections in painful rotator cuff tendinopathy: A systematic review and meta-analysis. *PM&R*, 8(11), 1127–1136. <https://doi.org/10.1016/j.pmrj.2016.04.001>
- Blau, H.M., Chiu, C.P. & Webster, C., 1983. Cytoplasmic activation of human nuclear genes in stable heterocaryons. *Cell*, 32(4), pp.1171–1180.
- Boyden, S., 1962. The chemotactic effect of mixtures of antibody and antigen on polymorphonuclear leucocytes. *Journal of Experimental Medicine*, 115(3), pp.453–466.
- Burattini, S., Ferri, P., Battistelli, M., Curci, R., Luchetti, F. & Falcieri, E. (2004). C2C12 murine myoblasts as a model of skeletal muscle development: morpho-functional characterization. *European Journal of Histochemistry*, 48(3), 223–233. <https://doi.org/10.4081/1184>
- Chen, M., Zheng, H., Wei, T., Wang, D., Xia, H., Zhao, L., Ji, J., & Gao, H. (2016). High Glucose- Induced PC12 Cell Death by Increasing Glutamate Production and Decreasing Methyl Group Metabolism. *BioMed research international*, 2016, 4125731. <https://doi.org/10.1155/2016/4125731>
- Coppinger, J.A., Cagney, G., Toomey, S., Kislinger, T., Belton, O., McRedmond, J.P., et al., 2004. Characterization of the proteins released from activated platelets leads to localization of novel platelet proteins in human atherosclerotic lesions. *Blood*, 103(6), pp.2096–2104.
- buCowley, S., Paterson, H., Kemp, P., & Marshall, C.J. (1994). Activation of MAP kinase kinase is necessary and sufficient for PC12 differentiation and for transformation of NIH 3T3 cells.

Cell, 77(6), 841–852.

DeChellis, D.M. & Cortazzo, M.H., 2011. Regenerative medicine in the field of pain medicine: Proloterapi, platelet-rich plasma therapy, and stem cell therapy—Theory and evidence. *Techniques in Regional Anesthesia and Pain Management*, 15(2), pp.74–80. <https://doi.org/10.1053/j.trap.2011.05.002>

Dohl, J., et al., 2018. Acclimation of C2C12 myoblasts to physiological glucose: implications for MAPK/ERK signaling. *ScienceDirect*.

Gavrieli, Y., Sherman, Y. & Ben-Sasson, S.A., 1992. Identification of programmed cell death in situ via specific labeling of nuclear DNA fragmentation. *Journal of Cell Biology*, 119(3), pp.493–501.

Gerdes, J., Schwab, U., Lemke, H. & Stein, H., 1984. Production of a mouse monoclonal antibody reactive with a human nuclear antigen associated with cell proliferation. *International Journal of Cancer*, 31(1), pp.13–20.

Gleberzon, B., et al., 2011. Proloterapi in the management of musculoskeletal pain. *The Journal of Musculoskeletal Medicine*, 28(10), pp.115–121.

Gratzner, H.G., 1982. Monoclonal antibody to 5-bromo- and 5-iododeoxyuridine: A new reagent for detection of DNA replication. *Science*, 218(4571), pp.474–475.

Greene, L.A. & Tischler, A.S., 1976. Establishment of a noradrenergic clonal line of rat adrenal pheochromocytoma cells which respond to nerve growth factor. *Proceedings of the National Academy of Sciences*, 73(7), pp.2424–2428.

Güran, Ş., et al., 2018. Dextrose solution used for proloterapi decreases cell viability and alters expression of growth and apoptotic factors in human fibroblast culture. *Gulhane Medical Journal*, 60, pp.xxx–xxx.

Harting, B. R., Fox, S. K., Foster, A. D., Motyka, T. M., & Hinkelman, A. N. (2024). In vitro model exploring the mechanisms of dextrose prolotherapy: Fibroblasts exposed to clinical concentrations of dextrose exhibit significant rebound effects 48 hours after exposure.

Heijnen, H.F., Debili, N., Vainchencker, W., Breton-Gorius, J., Geuze, H.J. & Sixma, J.J., 1998. Multivesicular bodies are an intermediate stage in the formation of platelet alpha-granules.

Blood, 91(7), p.2313.

Karasimav, Ö., 2023. Dextrose concentration for prolotherapy: A study on human neuroblastoma cells. *Gulhane Medical Journal*.

Katsetos, C.D., Legido, A., Perentes, E. & Mörk, S.J., 2003. Class III β -tubulin isotype: a key cytoskeletal protein at the crossroads of developmental neurobiology and tumor neuropathology. *Journal of Child Neurology*, 18(12), pp.851–866.

Kjaer, M., 2004. Role of extracellular matrix in adaptation of tendon and skeletal muscle to mechanical loading. *Physiological Reviews*, 84(2), pp.649–698.

Li, Y. T., Chen, S. R., Shen, Y. P., & Wu, Y. T. (2021). Ultrasound-guided perineural injection with dextrose for carpal tunnel syndrome: A randomized controlled trial. *Mayo Clinic Proceedings*, 96(2), 350–359. <https://doi.org/10.1016/j.mayocp.2020.06.031>

Liang, C.C., Park, A.Y. & Guan, J.L., 2007. In vitro scratch assay: a convenient and inexpensive method for analysis of cell migration in vitro. *Nature Protocols*, 2(2), pp.329–333.

Lin, J. A., Chang, K. V., Wu, W. T., & Özçakar, L. (2020). Comparative effectiveness of ultrasound-guided dextrose injection vs corticosteroid for carpal tunnel syndrome: A systematic review and meta-analysis. *Archives of Physical Medicine and Rehabilitation*, 101(5), 950–961. <https://doi.org/10.1016/j.apmr.2019.11.005>

Luo, W., Ai, L., Wang, B. F., & Zhou, Y. (2019). High glucose inhibits myogenesis and induces insulin resistance by down-regulating AKT signaling. *Biomedicine & pharmacotherapy = Biomedecine & pharmacotherapie*, 120, 109498. <https://doi.org/10.1016/j.biopha.2019.109498>

Lyftogt, J., 2005. Prolotherapy and Achilles tendinopathy: a prospective pilot study of an ultrasound- guided injection technique. *British Journal of Sports Medicine*.

Lyftogt, J. (2007). Subcutaneous prolotherapy treatment of refractory knee, shoulder, and lateral elbow pain. *Australasian Musculoskeletal Medicine*, 12(2), 110–112.

Lyons, A.B. & Parish, C.R., 1994. Determination of lymphocyte division by flow cytometry. *Journal of Immunological Methods*, 171(1), pp.131–137.

- Milano, G., Sánchez, M., Jo, C.H., Saccomanno, M.F., Thampatty, B.P. & Wang, J.H., 2019. Platelet-rich plasma in orthopaedic sports medicine: state of the art. *J ISAKOS: Joint Disorders Orthop Sports Med*, 4(4), pp.188.
- Mollinari, C., et al. (2002). Expression of class III β -tubulin in proliferating and differentiating rat brain neuroblasts in vitro. *European Journal of Neuroscience*, 16(11), 2094–2102.
- Mosmann, T., 1983. Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays. *Journal of Immunological Methods*, 65(1–2), pp.55–63.
- Nicholson, D.W., et al., 1995. Identification and inhibition of the ICE/CED-3 protease necessary for mammalian apoptosis. *Nature*, 376(6535), pp.37–43.
- Oh, J. H., Chung, S. W., Kim, J. Y., & Kim, S. H. (2019). Proloterapi for the treatment of chronic musculoskeletal pain: A systematic review and meta-analysis. *Clinical Rehabilitation*, 33(11), 1744–1757. <https://doi.org/10.1177/0269215519863494>
- Oh, S., et al., 2019. Proloterapi injections and physiotherapy used singly and in combination in the management of lateral epicondylalgia: a single-blind randomized clinical trial. *BMC Musculoskeletal Disorders*.
- Reeves, K. D., Sit, R. W. S., Rabago, D., & Yelland, M. (2015). Dextrose proloterapi: A narrative review of basic science, clinical research, and best treatment practices. *British Journal of Sports Medicine*, 50(20), 1207–1217. <https://doi.org/10.1136/bjsports-2015-095186>
- Sharifi, A.M., Mousavi, S.H., Farhadi, M. & Larijani, B., 2007. Study of high glucose-induced apoptosis in PC12 cells: role of Bax protein. *Journal of Pharmacological Sciences*, 104(3), pp.258–262.
- Sharma, H.M. & Neeraj, R., 2014. Use of proloterapi in the treatment of chronic musculoskeletal pain: An overview. *Journal of Pain Research*, 7, pp.287–292.
- Strober, W., 2001. Trypan blue exclusion test of cell viability. *Current Protocols in Immunology*, Appendix 3, pp.A3B.1–A3B.2.
- Subramaniam, S., & Unsicker, K. (2010). ERK and cell death: ERK1/2 in neuronal death. *FEBS*

Journal, 277(1), 22–29.

Sweatt, J.D., 2001. The neuronal MAP kinase cascade: a biochemical signal integration system subserving synaptic plasticity and memory. *Journal of Neurochemistry*, 76(1), pp.1–10.

Tischler, A. S. & Greene, L. A. (1980). Morphologic and cytochemical properties of a clonal line of rat adrenal pheochromocytoma cells (PC12) with neuronal properties. *Laboratory Investigation*, 42(6), 511–518.

Uberoi, R., Srikandarajah, N., & Saifuddin, A. (2019). The role of imaging in peripheral nerve disorders and interventions. *Clinical Radiology*, 74(9), 665–676. <https://doi.org/10.1016/j.crad.2019.05.012>

Van Pelt, R.S., 2010. Proloterapi injection technique of the elbow. *Journal of Proloterapi*.

Vermes, I., Haanen, C., Steffens-Nakken, H. & Reutelingsperger, C., 1995. A novel assay for apoptosis: flow cytometric detection of phosphatidylserine expression on early apoptotic cells using fluorescein labelled Annexin V. *Journal of Immunological Methods*, 184(1), pp.39–51.

Woo, M.S., et al., 2016. The proper concentrations of dextrose and lidocaine in regenerative injection therapy: An in vitro study. *Journal of Pain Research*, 9, pp.283–289. <https://doi.org/10.2147/JPR.S107122>

Woo, P. W., Park, J. H., & Lee, S. Y. (2016). Effects of dextrose concentrations on fibroblast proliferation and growth factor expression. *Annals of Rehabilitation Medicine*, 40(1), 11–18. <https://doi.org/10.5535/arm.2016.40.1.11>

Woo, M.S., et al., 2021. The proper concentrations of dextrose and lidocaine in fibroblast culture and implications for proloterapi. *Korean Journal of Pain / ePain*, 34(1), pp.19–27.

Wu, Y. T., Chen, S. R., Li, T. Y., Ho, T. Y., Shen, Y. P., Tsai, C. K., & Chen, L. C. (2017). Nerve hydrodissection for carpal tunnel syndrome: A prospective, randomized, double-blind, controlled trial. *Muscle & Nerve*, 56(5), 869–876. <https://doi.org/10.1002/mus.25693>

Wu, Y. T., Ke, M. J., Ho, T. Y., Li, T. Y., Shen, Y. P., Chen, L. C., & Chen, S. R. (2018).

Randomized double-blinded clinical trial of 5% dextrose versus corticosteroid injections for carpal tunnel syndrome. *Annals of Neurology*, 84(4), 601–610. <https://doi.org/10.1002/ana.25332>

Yaffe, D. & Saxel, O., 1977. Serial passaging and differentiation of myogenic cells isolated from dystrophic mouse muscle. *Nature*, 270(5639), pp.725–727.

Zhang, X., et al., 2012. A simple, sensitive, and rapid cell viability assay based on the reduction of WST-8. *Journal of Immunological Methods*, 379(1–2), pp.1–7.

Zhao, A.T., et al., 2022. A comprehensive update of prolotherapy in the management of musculoskeletal disorders. *PMC Review*.

Zhu, J., Jiang, Y., Xu, Y., & Zhao, J. (2020). Effectiveness of dextrose prolotherapy for lateral epicondylitis: A meta-analysis of randomized controlled trials. *American Journal of Physical Medicine & Rehabilitation*, 99(8), 701–708. <https://doi.org/10.1097/PHM.0000000000001444>