

DAFTAR PUSTAKA

- Abdelaziz, T. A., Mohamed, M. A. A., Ismail, W. I., Idreis, I. A. I., & Baz, E. G. (2023). A prospective cohort study of severe sepsis-induced dyslipidemia and changes in D-dimer levels in children: do they affect the prognosis? *Egyptian Pediatric Association Gazette*, 71(1), 93.
- Al Jalbout N, Balhara KS, Hamade B, Hsieh YH, Kelen GD, Bayram JD. Shock index as a predictor of hospital admission and inpatient mortality in a US national database of emergency departments. *Emerg Med J*. 2019;36(5):293–7. pmid:30910912
- Avci, M., & Doganay, F. (2022). Prognostic Performance of Shock Index, Diastolic Shock Index, Age Shock Index, and Modified Shock Index in COVID-19 Pneumonia. *Disaster Medicine and Public Health Preparedness*, 17, e189. <https://doi.org/10.1017/dmp.2022.110>
- Berger, T., Green, J., Horeczko, T., Hagar, Y., Garg, N., Suarez, A., Panacek, E., & Shapiro, N. (2013). Shock index and early recognition of sepsis in the emergency department: pilot study. *The Western Journal of Emergency Medicine*, 14(2), 168–174. <https://doi.org/10.5811/westjem.2012.8.11546>
- Chung JY, Hsu CC, Chen JH, Chen WL, Lin HJ, Guo HR, et al. Shock index predicted mortality in geriatric patients with influenza in the emergency department. *Am J Emerg Med*. 2019;37(3):391–4. pmid:29866414
- Do SN, Dao CX, Nguyen TA, et al. Sequential Organ Failure Assessment (SOFA) Score for predicting mortality in patients with sepsis in Vietnamese intensive care units: a multicentre, cross-sectional study. *BMJ Open* 2023;13:e064870. doi:10.1136/bmjopen-2022-064870
- Evans, L., Rhodes, A., Alhazzani, W., Antonelli, M., Coopersmith, C. M., French, C., Machado, F. R., McIntyre, L., Ostermann, M., & Prescott, H. C. (2021). Surviving sepsis campaign: international guidelines for management of sepsis and septic shock 2021. *Critical Care Medicine*, 49(11), e1063–e1143.
- Fernández, J. J. D., Castañeda-González, J. P., Zambrano, J. I. T., Martínez, F. E. D., & Ortiz, M. Á. S. (2024). Assessment of the shock index in septic shock: A systematic review. *Medicina Intensiva (English Edition)*.
- Font, M. D., Thyagarajan, B., & Khanna, A. K. (2020). Sepsis and Septic Shock—Basics of diagnosis, pathophysiology and clinical decision making. *Medical Clinics*, 104(4), 573–585.
- Gregoriano, C., Heilmann, E., Molitor, A., & Schuetz, P. (2020). Role of procalcitonin use in the management of sepsis. *Journal of Thoracic Disease*, 12(Suppl 1), S5–S15. <https://doi.org/10.21037/jtd.2019.11.63>
- Guarino, M., Perna, B., Cesaro, A. E., Maritati, M., Spampinato, M. D., Contini, C., & De Giorgio, R. (2023). 2023 update on sepsis and septic shock in adult patients: management in the emergency department. *Journal of Clinical Medicine*, 12(9), 3188.
- Gupta, S., & Tomar, D. S. (2019). Septic Shock and Troponin I: Are They in a Relationship? In *Indian journal of critical care medicine : peer-reviewed, official publication of Indian Society of Critical Care Medicine* (Vol. 23, Issue 7, p. 294). <https://doi.org/10.5005/jp-journals-10071-23201>

- Gupta, T., Puskarich, M. A., DeVos, E., Javed, A., Smotherman, C., Sterling, S. A., Wang, H. E., Moore, F. A., Jones, A. E., & Guirgis, F. W. (2020). Sequential organ failure assessment component score prediction of in-hospital mortality from sepsis. *Journal of Intensive Care Medicine*, 35(8), 810–817.
- Jain, S., Sinha, S., Sharma, S. K., Samantaray, J. C., Aggrawal, P., Vikram, N. K., Biswas, A., Sood, S., Goel, M., Das, M., Vishnubhatla, S., & Khan, N. (2014). Procalcitonin as a prognostic marker for sepsis: a prospective observational study. *BMC Research Notes*, 7, 458. <https://doi.org/10.1186/1756-0500-7-458>
- Jendoubi, A., Jerbi, S., Maamar, E., Abbess, A., Samoud, Z., Kanzari, L., Boutiba, I., Ghedira, S., & Houissa, M. (2019). Prognostic Value of High-sensitivity Troponin I in Patients with Septic Shock: A Prospective Observational Study. *Indian Journal of Critical Care Medicine: Peer-Reviewed, Official Publication of Indian Society of Critical Care Medicine*, 23(7), 320–325. <https://doi.org/10.5005/jp-journals-10071-23206>
- Jentzer JC, van Diepen S, Barsness GW, Henry TD, Menon V, Rihal CS, et al. Cardiogenic Shock Classification to Predict Mortality in the Cardiac Intensive Care Unit. *J Am Coll Cardiol*. 2019;74(17):2117–28.
- Jeon Y, Kim S, Ahn S, Park JH, Cho H, Moon S, Lee S. Predicting septic shock in patients with sepsis at emergency department triage using systolic and diastolic shock index. *Am J Emerg Med*. 2024 Apr;78:196-201. doi: 10.1016/j.ajem.2024.01.029. Epub 2024 Jan 28. PMID: 38301370.
- Jeon, Y., Kim, S., Ahn, S., Park, J.-H., Cho, H., Moon, S., & Lee, S. (2024). Predicting septic shock in patients with sepsis at emergency department triage level using systolic and diastolic shock index. *The American Journal of Emergency Medicine*, 78, 196–201.
- Jouffroy, R., Gille, S., Gilbert, B., Travers, S., Bloch-Laine, E., Ecollan, P., Boularan, J., Bounes, V., Vivien, B., & Gueye, P. (2024). Relationship between shock index, modified shock index, and age shock index and 28-day mortality among patients with prehospital septic shock. *The Journal of Emergency Medicine*, 66(2), 144–153.
- Kenzaka, T., Okayama, M., Kuroki, S., Fukui, M., Yahata, S., Hayashi, H., Kitao, A., Sugiyama, D., Kajii, E., & Hashimoto, M., 2012. Importance of vital signs to the early diagnosis and severity of sepsis: association between vital signs and sequential organ failure assessment score in patients with sepsis.. *Internal medicine*, 51 8, pp. 871-6 . <https://doi.org/10.2169/internalmedicine.51.6951>.
- Kilinc Toker A, Kose S, Turken M. Comparison of SOFA Score, SIRS, qSOFA, and qSOFA + L Criteria in the Diagnosis and Prognosis of Sepsis. *Eurasian J Med*. 2021;53(1):40-47. doi:10.5152/eurasianjmed.2021.20081
- Kim DS, Park JE, Hwang SY, et al. Prediction of vasopressor requirement among hypotensive patients with suspected infection: usefulness of diastolic shock index and lactate. *Clin Exp Emerg Med*. 2022;9:176–86. doi: 10.15441/ceem.22.324
- Kim, T. H., Jeong, D., Park, J. E., Hwang, S. Y., Suh, G. J., Choi, S.-H., Chung, S. P., Kim, W. Y., Lee, G. T., & Shin, T. G. (2023). Prognostic accuracy of initial and 24-h maximum SOFA scores of septic shock patients in the emergency department. *Heliyon*, 9(9).

- Koch, E., Lovett, S., Nghiem, T., Riggs, R. A., & Rech, M. A. (2019). Shock index in the emergency department: utility and limitations. *Open Access Emergency Medicine : OAEM*, 11, 179–199. <https://doi.org/10.2147/OAEM.S178358>
- La Via, L., Sangiorgio, G., Stefani, S., Marino, A., Nunnari, G., Cocuzza, S., La Mantia, I., Cacopardo, B., Stracquadanio, S., & Spampinato, S. (2024). The global burden of sepsis and septic shock. *Epidemiologia*, 5(3), 456–478.
- Lanspa, M. J., Brown, S. M., Hirshberg, E. L., Jones, J. P., & Grissom, C. K. (2012). Central venous pressure and shock index predict lack of hemodynamic response to volume expansion in septic shock: a prospective, observational study. *Journal of Critical Care*, 27(6), 609–615. <https://doi.org/10.1016/j.jcrc.2012.07.021>
- Lee, K. J., Kim, Y. K., Jeon, K., Ko, R.-E., Suh, G. Y., Oh, D. K., Lim, S. Y., Lee, Y. J., Lee, S. Y., & Park, M.-H. (2024). Shock indices are associated with in-hospital mortality among patients with septic shock and normal left ventricular ejection fraction. *Plos One*, 19(3), e0298617.
- Lee, S. M., & An, W. S. (2016). New clinical criteria for septic shock: serum lactate level as new emerging vital sign. *Journal of Thoracic Disease*, 8(7), 1388–1390. <https://doi.org/10.21037/jtd.2016.05.55>
- Li D, Wei Y, Zhang C, et al. Value of SOFA score, APACHE II score, and WBC count for mortality risk assessment in septic patients: A retrospective study. *Medicine (Baltimore)*. 2025;104(20):e42464. doi:10.1097/MD.00000000000042464
- Liu, C., Suo, S., Luo, L., Chen, X., Ling, C., & Cao, S. (2022). SOFA Score in relation to Sepsis: Clinical Implications in Diagnosis, Treatment, and Prognostic Assessment. *Computational and Mathematical Methods in Medicine*, 2022, 7870434. <https://doi.org/10.1155/2022/7870434>
- Mahashabde ML, Bhimani YR, Bhavsar HM. The Correlation Between the Lactate/Albumin Ratio and Sequential Organ Failure Assessment (SOFA) Score in Patients With Sepsis and Septic Shock. *Cureus*. 2024;16(7):e65616. Published 2024 Jul 29. doi:10.7759/cureus.65616
- Marshall, J. C., Bosco, L., Adhikari, N. K., Connolly, B., Diaz, J. V, Dorman, T., Fowler, R. A., Meyfroidt, G., Nakagawa, S., & Pelosi, P. (2017). What is an intensive care unit? A report of the task force of the World Federation of Societies of Intensive and Critical Care Medicine. *Journal of Critical Care*, 37, 270–276.
- Matsunaga, N., Yoshioka, Y., & Fukuta, Y. (2021). Extremely high troponin levels induced by septic shock: a case report. *Journal of Medical Case Reports*, 15, 1–4.
- Naderpour, Z., Momeni, M., Vahidi, E., Safavi, J., & Saeedi, M. (2019). Procalcitonin and D-dimer for predicting 28-day-mortality rate and sepsis severity based on SOFA score; a cross-sectional study. *Bulletin of Emergency & Trauma*, 7(4), 361.
- Naidu SS, Baran DA, Jentzer JC, Hollenberg SM, van Diepen S, Basir MB, et al. SCAI SHOCK Stage Classification Expert Consensus Update: A Review and Incorporation of Validation Studies: This statement was endorsed by the American College of Cardiology (ACC), American College of Emergency Physicians (ACEP), American Heart Association (AHA), European Society of Cardiology (ESC) Association for Acute Cardiovascular Care (ACVC), International Society for Heart and Lung Transplantation (ISHLT), Society of Critical Care Medicine (SCCM), and Society of Thoracic Surgeons (STS) in December 2021. *J Am Coll Cardiol*. 2022;79(9):933–46.

- Nguyen, T. T. M., Vu, M. P., Nguyen, T. T., Nguyen, T. T. M., Vu, H., Duong, H. Y., Pham, P. T., Hoang, T. H., & Kieu, T. V. O. (2025). Elevated plasma D-dimer level and the presence of septic shock in patients with sepsis. *European Journal of Inflammation*, 23, 1721727X251330015.
- Ospina-Tascon GA, Teboul JL, Hernandez G, Alvarez I, Sanchez-Ortiz AI, Calderon-Tapia LE, et al. Diastolic shock index and clinical outcomes in patients with septic shock. *Ann Intensive Care*. 2020;10:41. doi: 10.1186/s13613-020-00658-8.
- Ospina-Tascón GA, García-Gallardo G, Orozco N. Using the diastolic shock index to determine when to promptly administer vasopressors in patients with septic shock. *Clin Exp Emerg Med*. 2022;9(4):367-369. doi:10.15441/ceem.22.401
- Ospina-Tascón GA, Teboul JL, Hernandez G, et al. Diastolic shock index and clinical outcomes in patients with septic shock. *Ann Intensive Care*. 2020;10(1):41. Published 2020 Apr 16. doi:10.1186/s13613-020-00658-8
- Ospina-Tascón, G. A., Teboul, J.-L., Hernandez, G., Alvarez, I., Sánchez-Ortiz, A. I., Calderón-Tapia, L. E., Manzano-Nunez, R., Quiñones, E., Madriñan-Navia, H. J., Ruiz, J. E., Aldana, J. L., & Bakker, J. (2020). Diastolic shock index and clinical outcomes in patients with septic shock. *Annals of Intensive Care*, 10(1), 41. <https://doi.org/10.1186/s13613-020-00658-8>
- Padkins M, Kashani K, Tabi M, Gajic O, Jentzer JC (2024) Association between the shock index on admission and in-hospital mortality in the cardiac intensive care unit. *PLOS ONE* 19(4): e0298327. <https://doi.org/10.1371/journal.pone.0298327>
- Park, H., Lee, J., Oh, D.K. et al. Serial evaluation of the serum lactate level with the SOFA score to predict mortality in patients with sepsis. *Sci Rep* 13, 6351 (2023). <https://doi.org/10.1038/s41598-023-33227-7>
- Putra, I. M. P. (2018). Pendekatan Sepsis dengan Skor SOFA. *Cermin Dunia Kedokteran*, 45(8), 400599.
- Reinhart K, Bauer M, Riedemann NC, Hartog CS. New approaches to sepsis: molecular diagnostics and biomarkers. *Clin Microbiol Rev*. 2012;25:609–634. doi: 10.1128/CMR.00016-12
- Rodelo, J. R., De la Rosa, G., Valencia, M. L., Ospina, S., Arango, C. M., Gómez, C. I., García, A., Nuñez, E., & Jaimes, F. A. (2012). D-dimer is a significant prognostic factor in patients with suspected infection and sepsis. *The American Journal of Emergency Medicine*, 30(9), 1991–1999.
- Schmitt, F. C. F., Manolov, V., Morgenstern, J., Fleming, T., Heitmeier, S., Uhle, F., Al-Saeedi, M., Hackert, T., Bruckner, T., & Schöchl, H. (2019). Acute fibrinolysis shutdown occurs early in septic shock and is associated with increased morbidity and mortality: results of an observational pilot study. *Annals of Intensive Care*, 9, 1–15.
- Sharma, A., Sikka, M., Gomber, S., & Sharma, S. (2018). Plasma fibrinogen and D-dimer in children with sepsis: a single-center experience. *Iranian Journal of Pathology*, 13(2), 272.
- Singer, M., Deutschman, C. S., Seymour, C. W., Shankar-Hari, M., Annane, D., Bauer, M., Bellomo, R., Bernard, G. R., Chiche, J.-D., Coopersmith, C. M., Hotchkiss, R. S., Levy, M. M., Marshall, J. C., Martin, G. S., Opal, S. M., Rubenfeld, G. D., van der Poll, T., Vincent, J.-L., & Angus, D. C. (2016). The Third International Consensus Definitions

- for Sepsis and Septic Shock (Sepsis-3). *JAMA*, 315(8), 801–810. <https://doi.org/10.1001/jama.2016.0287>
- Stretch, B., & Shepherd, S. J. (2021). Criteria for intensive care unit admission and severity of illness. *Surgery (Oxford)*, 39(1), 22–28.
- Tang, J., Yuan, H., Wu, Y. L., Fu, S., & Pan, X. Y. (2023). The predictive value of heparin-binding protein and D-dimer in patients with sepsis. *International Journal of General Medicine*, 2295–2303.
- Thakur R, Naga Rohith V, Arora JK. Mean SOFA Score in Comparison With APACHE II Score in Predicting Mortality in Surgical Patients With Sepsis. *Cureus*. 2023;15(3):e36653. Published 2023 Mar 24. doi:10.7759/cureus.36653
- Wang, G., Liu, J., Xu, R., & Liu, X. (2022). Elevated plasma D-dimer levels are associated with the poor prognosis of critically ill children. *Frontiers in Pediatrics*, 10, 1001893.
- Wira, C. R., Francis, M. W., Bhat, S., Ehrman, R., Conner, D., & Siegel, M. (2014). The shock index as a predictor of vasopressor use in emergency department patients with severe sepsis. *The Western Journal of Emergency Medicine*, 15(1), 60–66. <https://doi.org/10.5811/westjem.2013.7.18472>
- World Health Organization. (2017). Clinical Criteria for the Determination of Death. September.
- Yeo HJ, Lee YS, Kim TH, et al. Vasopressor initiation within 1 hour of fluid loading is associated with increased mortality in septic shock patients: analysis of National Registry Data. *Crit Care Med*. 2022;50:e351–60. doi:10.1097/CCM.0000000000005363
- Yu, G. et al. Utility of the early lactate area score as a prognostic marker for septic shock patients in the emergency department. *Acute Crit. Care* 34, 126–132 (2019).
- Zhang TN, Hao PH, Gao SY, Liu CF, Yang N. Evaluation of SI, MSI and DSI for very early (3-day) mortality in patients with septic shock. *Eur J Med Res*. 2022;27(1):227. Published 2022 Nov 3. doi:10.1186/s40001-022-00857-y
- Zhang, T.-N., Hao, P.-H., Gao, S.-Y., Liu, C.-F., & Yang, N. (2022). Evaluation of SI, MSI and DSI for very early (3-day) mortality in patients with septic shock. *European Journal of Medical Research*, 27(1), 227.

