

DAFTAR PUSTAKA

1. Qiuru Wang, Jian Hu, Ting Ma. Comparison of Different Concentrations of Ropivacaine Used for Ultrasound-Guided *Adductor Canal Block* þ *IPACK Block* in *Total knee arthroplasty*, Thieme Medical Publishers 2022.
2. Wylde V, Rooker J, Halliday L, Blom A. Acute postoperative pain at rest after hip and knee arthroplasty: severity, sensory qualities and impact on sleep. *Orthop Traumatol Surg Res* 2011; 97 (02) 139-144.
3. Grosu I, Lavand'homme P, Thienpont E. Pain after knee arthroplasty: an unresolved issue. *Knee Surg Sports Traumatol Arthrosc* 2014; 22 (08) 1744-1758.
4. Burns LC, Ritvo SE, Ferguson MK, Clarke H, Seltzer Z, Katz J. Pain catastrophizing as a risk factor for chronic pain after *Total knee arthroplasty*: a systematic review. *J Pain Res* 2015; 8: 21-32.
5. Lovald ST, Ong KL, Lau EC, Joshi GP, Kurtz SM, Malkani AL. Readmission and complications for catheter and injection femoral nerve block administration after *Total knee arthroplasty* in the medicare population. *J Arthroplasty* 2015; 30 (12) 2076-2081.
6. Di Francesco A, Flamini S, Pizzoferrato R, Fusco P, Paglia A. Continuous intraarticular and periarticular levobupivacaine for management of pain relief after *Total knee arthroplasty*: a prospective randomized, double-blind pilot study. *J Orthop* 2016; 13 (03) 119-122.
7. Stowers MD, Manuopangai L, Hill AG, Gray JR, Coleman B, Munro JT. Enhanced recovery after surgery in elective hip and knee arthroplasty reduces length of hospital stay. *ANZ J Surg* 2016; 86 (06) 475-479.
8. Domagalska, M.; Reysner, T.; Kowalski, G.; Daroszewski, P.; Mularski, A.; Wiczorowska-Tobis, K. Pain Management, Functional Recovery, and Stress Response Expressed by NLR and PLR after the *IPACK Block* Combined with *Adductor Canal Block* for *Total knee arthroplasty*—A Prospective, Randomised, Double-Blinded Clinical Trial. *J. Clin. Med.* 2023, 12, 7088.
9. Wang, A.C.; Than, K.D.; Etame, A.B.; La Marca, F.; Park, P. Impact of Anesthesia on Transcranial Electric Motor Evoked Potential Monitoring during Spine Surgery: A Review of the Literature. *Neurosurg. Focus* 2009, 2.
10. Zhang, Z.; Wang, Y.; Liu, Y. Effectiveness of Continuous *Adductor Canal Block* versus Continuous Femoral Nerve Block in Patients with *Total knee arthroplasty*: A PRISMA Guided Systematic Review and Meta-Analysis. *Medicine* 2019.
11. Huber, J.; Lisi ński, P.; Kłoskowska, P.; Gronek, A.; Lisiewicz, E.; Trzeciak,

- T. Meniscus Suture Provides Better Clinical and Biomechanical Results at 1-Year Follow-up than Meniscectomy. *Arch. Orthop. Trauma Surg.* 2013.
12. Domagalska, M.; Reysner, T. Pain Management in *Total knee arthroplasty*. A Comprehensive Review. *Chir. Narz  ad w Ruchu I Ortop. Pol.* 2022, 87, 173–180.
 13. Fr d ric Martin, Val ria Martinez, Jean Xavier Mazoit, Didier Bouhassira, Kamel Cherif, Marc Edouard Gentili, Philippe Piriou, Marcel Chauvin, Dominique Fletcher; Antiinflammatory Effect of Peripheral Nerve Blocks after Knee Surgery: Clinical and Biologic Evaluation. *Anesthesiology* 2008; 109:484–490 .
 14. Bartfai T: Immunology: Telling the brain about pain. *Nature* 2001; 410:425, 427.
 15. Ek M, Engblom D, Saha S, Blomqvist A, Jakobsson PJ, Ericsson-Dahlstrand A: Inflammatory response: Pathway across the blood-brain barrier. *Nature* 2001; 410:430–1.
 16. Gentili ME, Mazoit JX, Samii KK, Fletcher D: The effect of a sciatic nerve block on the development of inflammation in carrageenan injected rats. *Anesth Analg* 1999; 89:979–84.
 17. Bagry H, de la Cuadra Fontaine JC, Asenjo JF, Bracco D, Carli F: Effect of a continuous peripheral nerve block on the inflammatory response in knee arthroplasty. *Reg Anesth Pain Med* 2008; 33:17–23.
 18. Bain, C.R., Myles, P.S., Corcoran, T. and Dieleman, J.M. (2023), Postoperative systemic inflammatory dysregulation and corticosteroids: a narrative review. *Anaesthesia*, 78: 356-370.
 19. D cke W-D, Randow F, Syrbe U, et al. Monocyte deactivation in septic patients: restoration by IFN-  treatment. *Nature Medicine* 1997; 3: 678–81.
 20. Williams LM, Ricchetti G, Sarma U, Smallie T, Foxwell BMJ. Interleukin-10 suppression of myeloid cell activation—a continuing puzzle. *Immunology* 2004; 113: 281–92.
 21. Lund J, Jenstrup MT, Jaeger P, S rensen AM, Dahl JB. Continuous adductor-canal-blockade for adjuvant post-operative analgesia after major knee surgery: preliminary results. *Acta Anaesthesiol Scand* 2011; 55 (01) 14-19.
 22. Grevstad U, Mathiesen O, Valentiner LS, Jaeger P, Hilsted KL, Dahl JB. Effect of *Adductor Canal Block* versus femoral nerve block on quadriceps strength, mobilization, and pain after *Total knee arthroplasty*: a randomized, blinded study. *Reg Anesth Pain Med* 2015; 40 (01) 3-10.
 23. Li D, Alqwbani M, Wang Q, Liao R, Yang J, Kang P. Efficacy of *Adductor Canal Block* combined with additional analgesic methods for postoperative analgesia in *Total knee arthroplasty*: a prospective, double-blind, randomized controlled study. *J Arthroplasty* 2020; 35 (12) 3554-3562.

24. Ilfeld BM, McCartney CJL. Searching for the optimal pain management technique after knee arthroplasty: analgesia is just the tip of the iceberg. *Anesthesiology* 2017; 126 (05) 768-770.
25. Kampitak W, Tanavalee A, Ngarmukos S, Amarase C. Opioid-sparing analgesia and enhanced recovery after *Total knee arthroplasty* using combined triple nerve blocks with local infiltration analgesia. *J Arthroplasty* 2019; 34 (02) 295-302.
26. Elliott CEMT, Soberon JR. The *Adductor Canal Block* combined with *IPACK* improves physical therapy performance and reduces hospital length of stay. Paper presented at: 40th Annual Regional Anesthesiology and Acute Pain Medicine Meeting (ASRA); Las Vegas, NV, 2015.
27. Wang Q, Hu J, Zeng Y, Li D, Yang J, Kang P. Efficacy of two unique combinations of nerve blocks on postoperative pain and functional outcome after *Total knee arthroplasty*: a prospective, double-blind, randomized controlled study. *J Arthroplasty* 2021; 36 (10) 3421-3431.
28. Sankineani SR, Reddy ARC, Eachempati KK, Jangale A, Gurava Reddy AV. Comparison of *Adductor Canal Block* and *IPACK* block (interspace between the popliteal artery and the capsule of the posterior knee) with *Adductor Canal Block* alone after *Total knee arthroplasty*: a prospective control trial on pain and knee function in immediate postoperative period. *Eur J Orthop Surg Traumatol* 2018; 28 (07) 1391-1395.
29. Ochroch J, Qi V, Badiola I. et al. Analgesic efficacy of adding the *IPACK* block to a multimodal analgesia protocol for primary *Total knee arthroplasty*. *Reg Anesth Pain Med* 2020; 45 (10) 799-804.
30. Li, J.; Ma, Y.; Xiao, L. Postoperative Pain Management in *Total knee arthroplasty*. *Orthop. Surg.* 2019, 11, 755–761.
31. Gao, J., Xing, D., Dong, S. et al. The primary *Total knee arthroplasty*: a global analysis. *J Orthop Surg Res* 15, 190 (2020).
32. Ghosh A, Chatterji U. An evidence-based review of enhanced recovery after surgery in total knee replacement surgery. *J Perioper Pract.* 2019;29(9):281–90.
33. Guo EW, Sayeed Z, Padela MT, Qazi M, Zekaj M, Schaefer P, Darwiche HF. Improving total joint replacement with continuous quality improvement methods and tools. *Orthop Clin North Am.* 2018.
34. Hsu H, Siwiec RM. Knee Arthroplasty. 2023 Jul 24. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2024 Jan–. PMID: 29939691.
35. Davies PS, Graham SM, Maqungo S, Harrison WJ. Total joint replacement in sub-Saharan Africa: a systematic review. *Trop Doct.* 2019 Apr;49(2):120-128.
36. Adie S, Harris I, Chuan A, Lewis P, Naylor JM. Selecting and optimising patients for *Total knee arthroplasty*. *Med J Aust.* 2019 Feb;210(3):135-141.

37. Raja SN, Carr DB, Cohen M, Finnerup NB, Flor H, Gibson S, et al. The revised International Association for the Study of Pain definition of pain: concepts, challenges, and compromises. *Pain*. 2020;1:161(9):1976-82.
38. European Society of Regional Anesthesia and Pain Therapy. Postoperative pain management good clinical practice.pdf [Internet]. Tersedia dari: <https://www.fpmx.com.au/resources/clinicians/postoperative-pain-management-good-clinical-practice.pdf>.
39. Corke P. Postoperative pain management. Tersedia dari: <https://www.nps.org.au/australian-prescriber/articles/postoperative-pain-management-1>.
40. Rao M. Acute Post Operative Pain. *Indian J Anaesth*. 2006;50(5):340.
41. Council (U.S.) NP, Organizations JC on A of H. Pain: Current Understanding of Assessment, Management, and Treatments. National Pharmaceutical Council, Incorporated; 2001.
42. Dureja GP, Iyer RN, Das G, Ahdal J, Narang P. Evidence and consensus recommendations for the pharmacological management of pain in India. *J Pain Res*. 2017;10:709-36.
43. Swieboda P, Filip R, Prystupa A, Drozd M. Assessment of pain: types, mechanism and treatment. *Ann Agric Environ Med AAEM*. 2013;Spec no. 1:2-7.
44. Breivik H, Borchgrevink PC, Allen SM, Rosseland LA, Romundstad L, Hals EKB, et al. Assessment of pain. *Br J Anaesth*. 2008;101(1):17-24.
45. Ong KS, Seymour RA. Pain measurement in humans. *Surg J R Coll Surg Edinb Irel*. 2004;2(1):15-27.
46. Chan, E.; Howle, R.; Onwochei, D.; Desai, N. Infiltration between the Popliteal Artery and the Capsule of the Knee (*IPACK*) Block in Knee Surgery: A Narrative Review. *Reg. Anesth. Pain Med*. 2021, 46, 784–805.
47. Grosso, M.J.; Murtaugh, T.; Lakra, A.; Brown, A.R.; Maniker, R.B.; Cooper, H.J.; Macaulay, W.; Shah, R.P.; Geller, J.A. *Adductor Canal Block Compared with Periarticular Bupivacaine Injection for Total knee arthroplasty: A Prospective Randomized Trial*. *JBJS* 2018, 100, 1141–1146.
48. Chan, E.; Howle, R.; Onwochei, D.; Desai, N. Infiltration between the Popliteal Artery and the Capsule of the Knee (*IPACK*) Block in Knee Surgery: A Narrative Review. *Reg. Anesth. Pain Med*. 2021, 46, 784–805.
49. Hussain, N.; Brull, R.; Vannabouathong, C.; Robinson, C.; Zhou, S.; D'Souza, R.; Sawyer, T.; Terkawi, A.S.; Abdallah, F.W. The Analgesic Effectiveness of Motor-Sparing Nerve Blocks for *Total knee arthroplasty: A Network Meta-Analysis*. *Anesthesiology* 2023, 139, 444–461.
50. Albrecht, E.; Wegrzyn, J.; Dabetic, A.; El-Boghdadly, K. The Analgesic Efficacy of *IPACK* after Knee Surgery: A Systematic Review and Meta-Analysis with Trial Sequential Analysis. *J. Clin. Anesth*. 2021, 72, 110305.

51. Et T, Korkusuz M, Basaran B, Yarimoğlu R, Toprak H, Bilge A, Kumru N, Dedeli İ. Comparison of *IPACK* and periarticular block with adductor block alone after *Total knee arthroplasty*: a randomized clinical trial. *J Anesth*. 2022 Apr;36(2):276-286. doi: 10.1007/s00540-022-03047-6. Epub 2022 Feb 14. PMID: 35157136; PMCID: PMC8853355.
52. Elmallah RK, Chughtai M, Khlopas A, Newman JM, Stearns KL, Roche M, Kelly MA, Harwin SF, Mont MA. Pain control in *Total knee arthroplasty*. *J Knee Surg*. 2018;31:504–513.
53. Kuang MJ, Ma JX, Fu L, He WW, Zhao J, Ma XL. Is *Adductor Canal Block* better than femoral nerve block in primary *Total knee arthroplasty*? A GRADE analysis of the evidence through a systematic review and meta-analysis. *J Arthroplasty*. 2017;32:3238–3248.
54. Wang Q, Hu J, Ma T, Zhao D, Yang J, Kang P. Comparison of Different Concentrations of Ropivacaine Used for Ultrasound-Guided Adductor Canal Block + *IPACK* Block in *Total knee arthroplasty*. *J Knee Surg*. 2023 Oct;36(12):1273-1282.
55. McMahon SE, Doran E, O'Brien S, Cassidy RS, Boldt JG, Beverland DE. Seventeen to Twenty Years of Follow-Up of the Low Contact Stress Rotating-Platform *Total knee arthroplasty* With a Cementless Tibia in All Cases. *J Arthroplasty*. 2019.
56. George AM, Liu M. Ropivacaine. 2023 Jul 31. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2024.
57. Aberg G. Toxicological and local anaesthetic effects of optically active isomers of two local anaesthetic compounds. *Acta Pharmacol Toxicol (Copenh)* 1972;31(4):273-86.
58. Graf BM, Abraham I, Eberbach N, Kunst G, Stowe DF, Martin E. Differences in cardiotoxicity of bupivacaine and ropivacaine are the result of physicochemical and stereoselective properties. *Anesthesiology*. 2002.
59. Alazawi W, Pirmadjid N, Lahiri R, Bhattacharya S. Inflammatory and immune responses to surgery and their clinical impact. *Annals of Surgery* 2016; 264: 73–80.
60. Takeuchi O, Akira S. Pattern recognition receptors and inflammation. *Cell* 2010; 140: 805–20.
61. O'Dwyer MJ, Owen HC, Torrance HDT. The perioperative immune response. *Current Opinion in Critical Care* 2015; 21: 342–2.
62. Konishi I. Basic Principle and Step-by-Step Procedure of Abdominal Hysterectomy: Part 2. *Surg J (N Y)*. 2018 Dec 26;5(Suppl 1):S11-S21.
63. Tohme S, Simmons RL, Tsung A. Surgery for cancer: a trigger for metastases. *Cancer Res* 2017.
64. Hou BJ, Du Y, Gu SX, Fan J, Wang R, Deng H, Guo DX, Wang L, Wang YY. General anesthesia combined with epidural anesthesia maintaining

appropriate anesthesia depth may protect excessive production of inflammatory cytokines and stress hormones in colon cancer patients during and after surgery. *Medicine (Baltimore)*. 2019.

65. Abrar Mohammad. Cardiovascular response during induction, laryngoscopy, intubation and surgery in normotensive and controlled hypertensive patients undergoing abdominal surgeries under general anaesthesia. *International Journal of Contemporary Medical Research* 2019.
66. Chen WK, Miao CH. The effect of anesthetic technique on survival in human cancers: a meta-analysis of retrospective and prospective studies. *PLoS One* 2013.
67. Buonacera A, Stancanelli B, Colaci M, Malatino L. Neutrophil to Lymphocyte Ratio: An Emerging Marker of the Relationships between the Immune System and Diseases. *Int J Mol Sci*. 2022 Mar 26;23(7).
68. He YM, Liu X, Zhong SY, Fu QH. Neutrophil-to-lymphocyte ratio in relation to trauma severity as prognosis factors in patients with multiple injuries complicated by multiple organ dysfunction syndrome: A retrospective analysis. *Immun Inflamm Dis*. 2023.
69. Heffernan DS, Monaghan SF, Thakkar RK, Machan JT, Cioffi WG, Ayala A. Failure to normalize lymphopenia following trauma is associated with increased mortality, independent of the leukocytosis pattern. *Crit Care*. 2012.
70. Zou P, Yang E, Li Z. Neutrophil-to-lymphocyte ratio is an independent predictor for survival outcomes in cervical cancer: a systematic review and meta-analysis. *Sci Rep*. 2020.
71. Cristian Bartoc; Robert J. Frumento; Maya Jalbout; Elliott Bennett-Guerrero; Evelyn Du; Ervant Nishanian (2006). A Randomized, Double-Blind, Placebo-Controlled Study Assessing the Anti-inflammatory Effects of Ketamine in Cardiac Surgical Patients. , 20(2), 0–222. doi:10.1053/j.jvca.2005.12.005.
72. Cleland, J., Koppenhaver, S., Su, J., & Netter, F. H. (2013). *Netter's orthopaedic clinical examination: an evidence-based approach* (Second edition). Elsevier.
73. Kandarian B, Indelli PF, Sinha S, Hunter OO, Wang RR, Kim TE, et al. Implementation of the IPACK (Infiltration between the Popliteal Artery and Capsule of the Knee) block into a multimodal analgesic pathway for total knee replacement. *Korean J Anesthesiol*. 2019;72(3):238-244.
74. NYSORA. iPACK block: infiltration between the popliteal artery and capsule of the knee (tips/technique). NYSORA [Internet]. [cited 2026 Feb 4].
75. Mou P, Wang X, Zhang X, Zhang Y, Li J, Zhang J. Adductor Canal Block Combined With IPACK Block for Postoperative Analgesia and Function Recovery Following Total Knee Arthroplasty: A Prospective Double-Blind Randomized Controlled Study. *J Arthroplasty*. 2022;37

76. Kertkiatkachorn W, Kampitak W, Tanavalee A, Ngarmukos S, Kerr SJ, et al. Adductor Canal Block Combined With iPACK (Interspace between the Popliteal Artery and Posterior Capsule of the Knee) Block vs periarticular injection (noninferiority) in total knee arthroplasty. (Acta Orthop/Anesth-related RCT). 2021.
77. Kruidenier J, Dingemans SA, Van Dieren S, De Jong VM, Goslings JC, Schepers T. C-reactive protein kinetics and its predictive value in orthopedic (trauma) surgery: a systematic review. *Acta Orthop Belg*. 2018;397-406.
78. Huang ZY, et al. Normal trajectory of interleukin-6 and C-reactive protein in the immediate hours and days after primary total knee arthroplasty under an ERAS protocol. 2020.
79. Schmidt F, Ward M, Repanos C. Postoperative serum C-reactive protein dynamics after pharyngolaryngectomy with jejunal free-flap reconstruction. *Ann R Coll Surg Engl*. 2023.
80. Agarwal N, Kay R, Duckworth AD, Clement ND, Griffith DM. Adductor canal block in total knee arthroplasty: a scoping review of the literature. *BJA Open*. 2025;14:100381.
81. Albrecht E, Wegrzyn J, Dabetic A, El-Boghdadly K. The analgesic efficacy of iPACK after knee surgery: a systematic review and meta-analysis with trial sequential analysis. *J Clin Anesth*. 2021;72:110305.
82. .Tang X, Lai Y, Wang S, et al. Analgesic efficacy of adding the IPACK block to multimodal analgesia programs for total knee arthroplasty: a systematic review and meta-analysis of randomized controlled trials. *J Orthop Surg Res*. 2022;17.
83. Zheng FY, Chen XY, et al. The impact of IPACK combined with adductor canal block under ultrasound guidance on early motor function after total knee arthroplasty. *Braz J Anesthesiol*. 2022;72(1):110-114.
84. Sung CH, et al. Comparative analysis of adductor canal block combined with iPACK (functional recovery/motor recovery). 2025.
85. Pryambodho P, et al. Comparison of proximal adductor canal block plus iPACK (total knee arthroplasty). *J Anesth Pain Intensive Care*. 2024;28(1):1-8.
86. Reysner T, et al. iPACK block with adductor canal block vs lumbar erector spinae plane block in pain management following total knee arthroplasty. (Randomized trial). 2025.